	1
	-
Shortest Path Tool	
NAACCR Town Hall Webinar	
Presented by the NAACCR GIS Committee 10/27/11	-
NAACCR	
]
Questions	
Submit all questions concerning content through the Q&A panel	-
If you experience technical difficulties, let us	
know via the chat panel or call us at 217-698	
0800 x 111	
² NAACCR	
	1
Agenda	
• Overview	-
Recinda Sherman, MPH, CTR Florida Cancer Data Systems	
Proximity to Treatment and Likelihood of	
Mastectomy	
– Christopher Johnson, MPH Cancer Data Registry of Idaho	
Iddito	
3 NAÁCCR	

Agenda

- Breast Cancer Stage at Diagnosis: Is Travel Time Important?
 - Kevin A Henry, Ph.D. University of Utah, Department of Geography
- Shortest Path Finder Tool Demonstration
 - Dan Goldberg, Ph.D. University of Southern California

4

NAACCR

NAACCR Shortest Path Tool Town Hall Meeting: Overview

Recinda Sherman, MPH CTR Senior Research Associate Florida Cancer Data Systems Sylvester Comprehensive Cancer Center Miller Medical School, University of Miami

NAACCR

The Issue

- Distance between residence and treatment facility impacts
 - Treatment type, adherence
 - Screening type, adherence
 - Prognosis
 - stage at diagnosis, recurrence, survival
- Measurement of distance and time
 - Complex
 - Approximates
 - Straight "crow flies" line

10/27/2011 **Shortest Pathfinder**

The Early Days

- Great Circle Distance Tool
- NAACCR GIS Committee
 - Chris Johnson
- 2005, SAS, available on website
- Estimates distance between residence and up to 5 case-level facilities
- "Crow Flies", curvature of earth



A Better Measure



Shortest Path Project

- Develop software to process batches of patient records
 Calculate driving distance and time using current road network
- Funding
 - Susan G. Komen For the Cure
 - 3 year, \$300,000 Grant, Oct 2007
- Collaboration

 - NAACCR GIS Committee
 Frank Boscoe, Kevin Henry, Chris Johnson, David Stinchcomb
 - University of Southern California
 - Dan Goldberg, Myles Cockburn, Kaveh Shahabi, John Wilson
 - NAACCR Office
 - · Josh Whitley, Charlie Blackburn

		_	=	$\overline{}$
ATA	Æ.	n	r	ח
IVA	ΑІ	ш		ĸ

Shortest Path Tool

- Calculates Shortest Travel Distance & Time
 - Between patient residence and treatment facility
 - Using actual street network
 - Utilizes comprehensive, current street level data
- Requires geocoded cases & facilities
- Geocoded facilities available on GIS Website
- Free to NAACCR Members
- Designed for cancer registry data
- MyNAACCR
 - Upload files to secure, encrypted server



Use of the NAACCR Shortest Path Tool in Health Services Research

NAACCR Shortest Path Town Hall Meeting October 27, 2011

Christopher Johnson, MPH, Cancer Data Registry of Idaho Kevin Henry, PhD, University of Utah Francis Boscoe, PhD, New York State Cancer Registry



Use of the NAACCR Shortest Path Tool in Health Services Research

- 1. Geographic proximity to treatment for early stage breast cancer and likelihood of mastectomy.
 - "Treatment Paper"
- 2. Breast Cancer Stage at Diagnosis: Is Travel Time Important?
 - "Stage Paper"
- 3. Ideas for future research

Komen GIS Project Protocol

- Participating registries were asked to prepare two data files:
- Latitude and longitude coordinates of cases and facilities that reported them were submitted to a secure file server at the University of Southern California GIS Lab for road network calculation purposes.
- Cancer registry data fields plus the distance measures; submitted to NAACCR and used to conduct the research projects resulting in the two papers.

13

NAACCR

Case Selection Criteria - Source Records

- Female in situ and invasive primary breast cancer cases diagnosed during 2004-2006.
- Reportable cases (i.e. residents of geographic areas covered by respective registries at time of diagnosis).
- Because we used distances between patient location and facility location(s), we needed to use <u>source</u> <u>records</u> for this project, not consolidated records.
 - Data files contained one record per case <u>per reporting</u> <u>facility</u>, with one unique identifying number per individual.

14

NAACCI



Shortest Pathfinder

Geographic proximity to treatment for early stage breast cancer and likelihood of mastectomy.

> Boscoe FP, Johnson CJ, Henry KA, Goldberg DW, Shahabi K, Elkin EB, Ballas LK, Cockburn M. Breast. 2011 Aug;20(4):324-8.

NAACCR Shortest Path Town Hall Meeting October 27, 2011 Presented by: Christopher Johnson, MPH Cancer Data Registry of Idaho

NAACCR

Treatment Paper – Introduction

- Women diagnosed with early stage invasive breast cancer in the United States are typically presented with a choice of mastectomy or breast conserving surgery (BCS) with radiation therapy (RT).
 - Two options have long been established as equally effective in terms of survival.
 - Following the National Institutes of Health's 1990 consensus statement that BCS with RT was the preferred treatment for most women, national mastectomy rates fell steadily through 2006.

17

NAACCR

Treatment Paper - Methods

- Early stage breast cancer cases
 - Collaborative Stage-derived SEER Summary Stage 2000 localized cases
 - Women diagnosed between 2004 and 2006.
- Cases were limited to women who had either:
 - Breast conserving surgery (FORDS codes 20-24)
 - Mastectomy (FORDS codes 30-80)
- Radiation treatment data were also available, but because of differential reporting by state were determined not to be of sufficient quality to use in the analysis.

18

Treatment Paper - Methods

- Two distance measures were calculated for each patient:
- 1. Travel distance via roads from the patient's residence to the location of surgical treatment.
 - Realized access to cancer care.
 - For patients whose diagnosis and treatment spanned multiple facilities, the minimum distance from among the candidate facilities was used.
- 2. Travel distance via roads from the road nearest the centroid of the patient's census tract to the nearest radiation treatment location.
 - Potential access to RT.

19



Treatment Paper - Methods

- The likelihood of receiving breast conserving surgery versus mastectomy was modeled using multilevel logistic regression using PROC GLIMMIX in SAS Version 9.2.
- Two models were developed, one using each distance measure.
- · State random effect.
- Fixed effects in the model included:
 - age
- census tract-level
- year of diagnosis
- household poverty rate
- Race
- rural/urban status
- Ethnicity
- whether the tumor was a person's first primary
- •

tumor.



Treatment Paper – Methods


Bypass Patients


- We were further interested in assessing the probability of mastectomy for patients who traveled more than 60 km for surgery but who lived within 30 km of a radiation treatment center.
 - Seek care at locations outside of their own community, typically at large, urban, nationally recognized hospitals.

21



Shortest Pathfinder





Treatment Paper - Results

Of the 2678 patients defined as <u>bypass patients</u> (>60 km from surgery and <30 km from nearest RT),
 42.2% received mastectomy (OR=1.27).

24

Treatment Paper - Conclusions

- In real estate, it is: **LOCATION, LOCATION, LOCATION**.
- In early stage breast cancer surgery, it is:
 - LOCATION (Distance to Treatment)
 - **LOCATION** (Regional/State Differences)
 - And some other things.
- Geographic distance is a fundamental element of patterns of cancer care.

25

NAACCR

Breast Cancer Stage at Diagnosis: Is Travel Time Important?

Henry KA, Boscoe FP, Johnson CJ, Goldberg DW, Sherman R, Cockburn M. Journal of Community Health. 2011, in press

Presented by: Kevin A Henry, Ph.D.

University of Utah, Department of Geography

NAACCR

Introduction

- Screening mammography has increased early detection and reduced breast cancer mortality (Taplin et al 2004)
- Breast cancer stage at diagnosis is related to the utilization of early detection procedures and following recommended guidelines.
- Many factors associated with stage at diagnosis:

Age Health Insurance
Facility Capacity Race/ethnicity
Obesity History of mammography
Area Poverty Geographic Accessibility

NAACCR

Introduction

- Geographic barriers (e.g., lengthy travel) to mammography could discourage screening
- Previous Study Results

		A	ssociation of proximit	:y		
			to nearest			
	Study		mammography and	Pub		
Geographic location	Period	N	stage at diagnosis	Year	Author	
Los Angeles	1992-1996	24,993	Yes	2006	Gumpertz	
Northern England	1994-2002	28,002	No	2008	Jones	
Metropolitan Chicago	1996-1998	4,936	No	2008	Tarlov	
Western Washington State	1996-1999	1,012	No	2011	Onega	
New Hampshire	1998-2004	6,305	No	2009	Celaya	
Detroit Metropolitan Area	1998-2002	12,413	Yes	2011	Dai	
Illinois	1998-2000	31,511	No	2010	Wang	
Kentucky	1999-2003	12,322	Yes	2009	Huang	an
Virginia	2000-2001	8,170	No	2009	Schroen	CR

Study Objectives:

Investigate whether geographic access to mammography screening is associated with the stage of breast cancer at diagnosis through two measures of accessibility:

- 1. Travel time to location of diagnosing facility
- 2. Travel time to location of nearest mammography

29

NAACCR

Study Population and Data

 Female breast cancer cases from ten different state cancer registries (2004-2006)

Arkansas	California	Idaho	Iowa	North Carolina
Kentucky	New Hampshire	New Jersey	New York	Oregon

- Histologically confirmed first primary in situ or invasive tumor of the breast (ICD-O3 C500-C509; excluding histology codes 9590–9989(N=174,609)
- SEER Summary Stage 2000
- · Aged 40 and older

30

Methods

- Two travel distance measures were calculated for each patient:
- 1. Realized (Actual) access: From the patient's residence to the <u>actual</u> location of diagnosis
- 2. Potential access: From the patient's census tract to the nearest mammography facility.
 - (Geocoded 2005 FDA Mammography facilities from contiguous United States N=8,770)


31

NAACCR


Methods

- $\hfill\Box$ Outcome: Odds of late stage (regional, distant) $\underline{vs}.$ early stage (in situ, localized)
- ☐ Two multilevel logistic regression models each using one travel time measure (facility and mammography)
 - ☐ Fixed Effects: travel time, age, race/ethnicity, diagnosis year, insurance, area poverty, rural/urban
- ☐ Random Effects: State
- ☐ PROC GLIMMIX, SAS Version 9.2

32



Shortest Pathfinder



Results

Significant factors associated with late stage breast cancer at diagnosis

Odds of Late Stage	Adjusted OR (95% CI)
Uninsured vs. Insured	1.79(1.64, 1.95)
Medicaid vs. Insured	1.74 (1.67, 1.83)
Age <50 vs. 65+	1.27(1.24, 1.31)
Age 50-64 vs. 65+	1.15 (1.11, 1.19)
Non-Hispanic Black vs. Non-Hispanic	1.31 (1.26, 1.37)
Hispanic vs. Non-Hispanic White	1.15 (1.09, 1.21)
High poverty census tracts (>20%) vs.	
low poverty tracts (<5%)	1.35 (1.30, 1.41)

N/

Conclusions

- Travel time to diagnosing facility or nearest mammography facility was not a determinant of late stage breast cancer at diagnosis
- Better geographic proximity did not assure more favorable stage distributions.
- Other factors were more important predictors of stage at diagnosis (e.g., insurance, census poverty, race/ethnicity)
- Additional measures of geographic accessibility should be examined.

 NAACCR

Possible Research Uses of the Shortest Path Tool

Francis Boscoe, PhD, New York State Cancer Registry

3


NAACCR


Possible research uses for the Shortest Path Tool

1. Identify underserved areas

38


NAACCI







Possible research uses for the Shortest Path Tool

2. Identify disparities that may be attributable to distance





Possible research uses for the Shortest Path Tool 3. Measure the impact that hospital closures have on travel time and whether this translates to changes in outcomes NAACCR



Possible research uses for the Shortest Path Tool

4. Identify and characterize patients who do not seek treatment at his or her closest location ("bypass patients")

46

NAACCR



Volume 20, Number 3, August 2009 E-ISSN: 1548-6869 Print ISSN: 1049-2089

Do Patients Bypass Rural Hospitals?

locé 1 Escarce MD 9hD and Vanika Vanur 9hD

Using data from California hospitals, the paper shows that patients were more likely to choose nearby hospitals, larger hospitals, and hospitals that offered more services and technologies. However, even after adjusting for these factors, patients had a propensity to bypass rural hospitals in favor of large urban hospitals. Offering additional services and technologies would increase the share of rural residents choosing rural hospitals only slightly.



Possible research uses for the Shortest Path Tool

5. Patient navigation: given facility locations and addresses of patients and volunteers, what is the most efficient way of pairing patients and volunteers?

48

OUESTIONS		
QUESTIONS?		
49	NAACCR	
	. u s 1001t	-
Shortest Path Finder Tool		
Dan Goldberg, PhD		
Research Assistant Professor of		
Spatial Science at the University of Southern California	V4 / 200	
50	B (B -6 /7/717)	
•	NAACCR	
	NAACCK	
	NAACCK	
	NAACCR	
OUESTIONS?	NAACCR	
QUESTIONS?	NAACCR	
QUESTIONS?	NAACCR	

	7
	-
Send questions or comments to Jim Hofferkamp inhofferkamp@naaccr.org A recording of this webinar will be posted to the NAACCR website	
THANK YOU!!!	
NAACCR NAACCR	RI