Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates[1] and 95% Confidence Intervals[2] for All Sites
Selected Areas in the United States[3], 1995-2000, Hispanic/Latino, All Races



1. Rates are per 100,000 and are adjusted to the 2000 U.S. population standard. Rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the rates for the U.S. combined.
2. See http://seer.cancer.gov/seerstat/WebHelp/seerstat.htm for the methodology used in the calculation of the age-adjusted 95% confidence intervals.
3. See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.

Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates[1] and 95% Confidence Intervals[2] for Bladder
Selected Areas in the United States[3], 1995-2000, Hispanic/Latino, All Races



1. Rates are per 100,000 and are adjusted to the 2000 U.S. population standard. Rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the rates for the U.S. combined.
2. See http://seer.cancer.gov/seerstat/WebHelp/seerstat.htm for the methodology used in the calculation of the age-adjusted 95% confidence intervals.
3. See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.

Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates[1] and 95% Confidence Intervals[2] for Brain and Other Nervous System
Selected Areas in the United States[3], 1995-2000, Hispanic/Latino, All Races



1. Rates are per 100,000 and are adjusted to the 2000 U.S. population standard. Rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the rates for the U.S. combined.
2. See http://seer.cancer.gov/seerstat/WebHelp/seerstat.htm for the methodology used in the calculation of the age-adjusted 95% confidence intervals.
3. See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.

Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates[1] and 95% Confidence Intervals[2] for Female Breast
Selected Areas in the United States[3], 1995-2000, Hispanic/Latina, All Races



1. Rates are per 100,000 and are adjusted to the 2000 U.S. population standard. Rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the rates for the U.S. combined.
2. See http://seer.cancer.gov/seerstat/WebHelp/seerstat.htm for the methodology used in the calculation of the age-adjusted 95% confidence intervals.
3. See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.

Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates[1] and 95% Confidence Intervals[2] for Buccal Cavity and Pharynx
Selected Areas in the United States[3], 1995-2000, Hispanic/Latino, All Races



1. Rates are per 100,000 and are adjusted to the 2000 U.S. population standard. Rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the rates for the U.S. combined.
2. See http://seer.cancer.gov/seerstat/WebHelp/seerstat.htm for the methodology used in the calculation of the age-adjusted 95% confidence intervals.
3. See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.

Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates[1] and 95% Confidence Intervals[2] for Cervix Uteri
Selected Areas in the United States[3], 1995-2000, Hispanic/Latina, All Races



1. Rates are per 100,000 and are adjusted to the 2000 U.S. population standard. Rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the rates for the U.S. combined.
2. See http://seer.cancer.gov/seerstat/WebHelp/seerstat.htm for the methodology used in the calculation of the age-adjusted 95% confidence intervals.
3. See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.

Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates[1] and 95% Confidence Intervals[2] for Colon and Rectum
Selected Areas in the United States[3], 1995-2000, Hispanic/Latino, All Races



1. Rates are per 100,000 and are adjusted to the 2000 U.S. population standard. Rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the rates for the U.S. combined.
2. See http://seer.cancer.gov/seerstat/WebHelp/seerstat.htm for the methodology used in the calculation of the age-adjusted 95% confidence intervals.
3. See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.

Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates[1] and 95% Confidence Intervals[2] for Corpus Uteri and Uterus, NOS
Selected Areas in the United States[3], 1995-2000, Hispanic/Latina, All Races



1. Rates are per 100,000 and are adjusted to the 2000 U.S. population standard. Rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the rates for the U.S. combined.
2. See http://seer.cancer.gov/seerstat/WebHelp/seerstat.htm for the methodology used in the calculation of the age-adjusted 95% confidence intervals.
3. See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.

Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates[1] and 95% Confidence Intervals[2] for Gallbladder
Selected Areas in the United States[3], 1995-2000, Hispanic/Latino, All Races



1. Rates are per 100,000 and are adjusted to the 2000 U.S. population standard. Rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the rates for the U.S. combined.
2. See http://seer.cancer.gov/seerstat/WebHelp/seerstat.htm for the methodology used in the calculation of the age-adjusted 95% confidence intervals.
3. See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.

Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates[1] and 95% Confidence Intervals[2] for Kidney and Renal Pelvis
Selected Areas in the United States[3], 1995-2000, Hispanic/Latino, All Races



1. Rates are per 100,000 and are adjusted to the 2000 U.S. population standard. Rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the rates for the U.S. combined.
2. See http://seer.cancer.gov/seerstat/WebHelp/seerstat.htm for the methodology used in the calculation of the age-adjusted 95% confidence intervals.
3. See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.

Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates[1] and 95% Confidence Intervals[2] for Leukemias
Selected Areas in the United States[3], 1995-2000, Hispanic/Latino, All Races



1. Rates are per 100,000 and are adjusted to the 2000 U.S. population standard. Rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the rates for the U.S. combined.
2. See http://seer.cancer.gov/seerstat/WebHelp/seerstat.htm for the methodology used in the calculation of the age-adjusted 95% confidence intervals.
3. See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.

Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates[1] and 95% Confidence Intervals[2] for Liver
Selected Areas in the United States[3], 1995-2000, Hispanic/Latino, All Races



1. Rates are per 100,000 and are adjusted to the 2000 U.S. population standard. Rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the rates for the U.S. combined.
2. See http://seer.cancer.gov/seerstat/WebHelp/seerstat.htm for the methodology used in the calculation of the age-adjusted 95% confidence intervals.
3. See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.

Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates[1] and 95% Confidence Intervals[2] for Lung and Bronchus
Selected Areas in the United States[3], 1995-2000, Hispanic/Latino, All Races



1. Rates are per 100,000 and are adjusted to the 2000 U.S. population standard. Rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the rates for the U.S. combined.
2. See http://seer.cancer.gov/seerstat/WebHelp/seerstat.htm for the methodology used in the calculation of the age-adjusted 95% confidence intervals.
3. See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.

Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates[1] and 95% Confidence Intervals[2] for Lymphomas
Selected Areas in the United States[3], 1995-2000, Hispanic/Latino, All Races



1. Rates are per 100,000 and are adjusted to the 2000 U.S. population standard. Rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the rates for the U.S. combined.
2. See http://seer.cancer.gov/seerstat/WebHelp/seerstat.htm for the methodology used in the calculation of the age-adjusted 95% confidence intervals.
3. See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.

Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates[1] and 95% Confidence Intervals[2] for Ovary
Selected Areas in the United States[3], 1995-2000, Hispanic/Latina, All Races



1. Rates are per 100,000 and are adjusted to the 2000 U.S. population standard. Rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the rates for the U.S. combined.
2. See http://seer.cancer.gov/seerstat/WebHelp/seerstat.htm for the methodology used in the calculation of the age-adjusted 95% confidence intervals.
3. See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.

Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates[1] and 95% Confidence Intervals[2] for Pancreas
Selected Areas in the United States[3], 1995-2000, Hispanic/Latino, All Races



1. Rates are per 100,000 and are adjusted to the 2000 U.S. population standard. Rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the rates for the U.S. combined.
2. See http://seer.cancer.gov/seerstat/WebHelp/seerstat.htm for the methodology used in the calculation of the age-adjusted 95% confidence intervals.
3. See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.

Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates[1] and 95% Confidence Intervals[2] for Prostate
Selected Areas in the United States[3], 1995-2000, Hispanic/Latino, All Races



1. Rates are per 100,000 and are adjusted to the 2000 U.S. population standard. Rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the rates for the U.S. combined.
2. See http://seer.cancer.gov/seerstat/WebHelp/seerstat.htm for the methodology used in the calculation of the age-adjusted 95% confidence intervals.
3. See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.

Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates[1] and 95% Confidence Intervals[2] for Stomach
Selected Areas in the United States[3], 1995-2000, Hispanic/Latino, All Races



1. Rates are per 100,000 and are adjusted to the 2000 U.S. population standard. Rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the rates for the U.S. combined.
2. See http://seer.cancer.gov/seerstat/WebHelp/seerstat.htm for the methodology used in the calculation of the age-adjusted 95% confidence intervals.
3. See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.

Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates[1] and 95% Confidence Intervals[2] for Testis
Selected Areas in the United States[3], 1995-2000, Hispanic/Latino, All Races



1. Rates are per 100,000 and are adjusted to the 2000 U.S. population standard. Rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the rates for the U.S. combined.
2. See http://seer.cancer.gov/seerstat/WebHelp/seerstat.htm for the methodology used in the calculation of the age-adjusted 95% confidence intervals.
3. See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.

Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates[1] and 95% Confidence Intervals[2] for Thyroid
Selected Areas in the United States[3], 1995-2000, Hispanic/Latino, All Races



1. Rates are per 100,000 and are adjusted to the 2000 U.S. population standard. Rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the rates for the U.S. combined.
2. See http://seer.cancer.gov/seerstat/WebHelp/seerstat.htm for the methodology used in the calculation of the age-adjusted 95% confidence intervals.
3. See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.

This Page Intentionally Left Blank

Average Annual Age-adjusted (2000 U.S. Standard) U.S. Cancer Incidence Rates (NAACCR Combined)[1] for Selected Cancer Sites[2] by Race/Ethnicity, Males, 1995-2000



1. Rates are per 100,000 and are adjusted to the 2000 U.S. population standard. Rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the rates for the U.S. combined.
2. See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.

Average Annual Age-adjusted (2000 U.S. Standard) U.S. Cancer Incidence Rates (NAACCR Combined)[1] for Selected Cancer Sites[2] by Race/Ethnicity, Females, 1995-2000



1. Rates are per 100,000 and are adjusted to the 2000 U.S. population standard. Rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the rates for the U.S. combined.
2. See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.