# Section V

## Comparative Graphs of Average-annual Age-adjusted Cancer Incidence Rates, by Registry, Among Hispanic Latino Populations, Selected Sites

This Page Intentionally Left Blank

**Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates[1] and 95% Confidence Intervals[2]**
**Selected Areas in the United States[3], 1999-2003, Hispanic/Latino, All Races, Males**




1 Rates are per 100,000 and are adjusted to the 2000 U.S. population standard. Rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the rates for the U.S. combined.

2. See http://seer.cancer.gov/seerstat/WebHelp/seerstat.htm for the methodology used in the calculation of the age-adjusted 95% confidence intervals.

3. See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.

N.B. In areas with small Latino populations, methods to indirectly identify Latinos (like NHIA) can overestimate the ethnicity-specific counts of cancer cases. Also, even small errors in Latino population estimates can affect the magnitude of the cancer rates.

**Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates[1] and 95% Confidence Intervals[2]**
**Selected Areas in the United States[3], 1999-2003, Hispanic/Latina, All Races, Females**




[1] Rates are per 100,000 and are adjusted to the 2000 U.S. population standard. Rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the rates for the U.S. combined.

[2] See http://seer.cancer.gov/seerstat/WebHelp/seerstat.htm for the methodology used in the calculation of the age-adjusted 95% confidence intervals.

[3] See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.

N.B. In areas with small Latino populations, methods to indirectly identify Latinos (like NHIA) can overestimate the ethnicity-specific counts of cancer cases. Also, even small errors in Latino population estimates can affect the magnitude of the cancer rates.

**Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates[1] and 95% Confidence Intervals[2]**
**Selected Areas in the United States[3], 1999-2003, Hispanic/Latino, All Races, Males**




[1] Rates are per 100,000 and are adjusted to the 2000 U.S. population standard. Rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the rates for the U.S. combined.

[2] See http://seer.cancer.gov/seerstat/WebHelp/seerstat.htm for the methodology used in the calculation of the age-adjusted 95% confidence intervals.

[3] See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics

N.B. In areas with small Latino populations, methods to indirectly identify Latinos (like NHIA) can overestimate the ethnicity-specific counts of cancer cases. Also, even small errors in Latino population estimates can affect the magnitude of the cancer rates.

**Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates[1] and 95% Confidence Intervals[2]**
**Selected Areas in the United States[3], 1999-2003, Hispanic/Latina, All Races, Females**

**Bladder**




[1] Rates are per 100,000 and are adjusted to the 2000 U.S. population standard. Rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the rates for the U.S. combined.

[2] See http://seer.cancer.gov/seerstat/WebHelp/seerstat.htm for the methodology used in the calculation of the age-adjusted 95% confidence intervals.

[3] See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics

N.B. In areas with small Latino populations, methods to indirectly identify Latinos (like NHIA) can overestimate the ethnicity-specific counts of cancer cases. Also, even small errors in Latino population estimates can affect the magnitude of the cancer rates.

**Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates[1] and 95% Confidence Intervals[2]**
**Selected Areas in the United States[3], 1999-2003, Hispanic/Latino, All Races, Males**

**Brain and Other Nervous System**




[1] Rates are per 100,000 and are adjusted to the 2000 U.S. population standard. Rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the rates for the U.S. combined.

[2] See http://seer.cancer.gov/seerstat/WebHelp/seerstat.htm for the methodology used in the calculation of the age-adjusted 95% confidence intervals.

[3] See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics

N.B. In areas with small Latino populations, methods to indirectly identify Latinos (like NHIA) can overestimate the ethnicity-specific counts of cancer cases. Also, even small errors in Latino population estimates can affect the magnitude of the cancer rates.

**Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates[1] and 95% Confidence Intervals[2]**
**Selected Areas in the United States[3], 1999-2003, Hispanic/Latina, All Races, Females**

**Brain and Other Nervous System**




1 Rates are per 100,000 and are adjusted to the 2000 U.S. population standard. Rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the rates for the U.S. combined.

2. See http://seer.cancer.gov/seerstat/WebHelp/seerstat.htm for the methodology used in the calculation of the age-adjusted 95% confidence intervals.

3. See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics

N.B. In areas with small Latino populations, methods to indirectly identify Latinos (like NHIA) can overestimate the ethnicity-specific counts of cancer cases. Also, even small errors in Latino population estimates can affect the magnitude of the cancer rates.

This Page Intentionally Left Blank

**Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates[1] and 95% Confidence Intervals[2]**
**Selected Areas in the United States[3], 1999-2003, Hispanic/Latina, All Races, Females**




1 Rates are per 100,000 and are adjusted to the 2000 U.S. population standard. Rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the rates for the U.S. combined.

2. See http://seer.cancer.gov/seerstat/WebHelp/seerstat.htm for the methodology used in the calculation of the age-adjusted 95% confidence intervals.

3. See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics

N.B. In areas with small Latino populations, methods to indirectly identify Latinos (like NHIA) can overestimate the ethnicity-specific counts of cancer cases. Also, even small errors in Latino population estimates can affect the magnitude of the cancer rates.

**Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates[1] and 95% Confidence Intervals[2]**
**Selected Areas in the United States[3], 1999-2003, Hispanic/Latino, All Races, Males**

**Buccal Cavity and Pharynx**




[1] Rates are per 100,000 and are adjusted to the 2000 U.S. population standard. Rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the rates for the U.S. combined.

[2] See http://seer.cancer.gov/seerstat/WebHelp/seerstat.htm for the methodology used in the calculation of the age-adjusted 95% confidence intervals.

[3] See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.

N.B. In areas with small Latino populations, methods to indirectly identify Latinos (like NHIA) can overestimate the ethnicity-specific counts of cancer cases. Also, even small errors in Latino population estimates can affect the magnitude of the cancer rates.

**Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates[1] and 95% Confidence Intervals[2]**
**Selected Areas in the United States[3], 1999-2003, Hispanic/Latina, All Races, Females**

**Buccal Cavity and Pharynx**




1 Rates are per 100,000 and are adjusted to the 2000 U.S. population standard. Rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the rates for the U.S. combined.

2. See http://seer.cancer.gov/seerstat/WebHelp/seerstat.htm for the methodology used in the calculation of the age-adjusted 95% confidence intervals.

3. See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.

N.B. In areas with small Latino populations, methods to indirectly identify Latinos (like NHIA) can overestimate the ethnicity-specific counts of cancer cases. Also, even small errors in Latino population estimates can affect the magnitude of the cancer rates.

This Page Intentionally Left Blank

**Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates[1] and 95% Confidence Intervals[2]**
**Selected Areas in the United States[3], 1999-2003, Hispanic/Latina, All Races, Females**

**Cervix Uteri**




1 Rates are per 100,000 and are adjusted to the 2000 U.S. population standard. Rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the rates for the U.S. combined.

2. See http://seer.cancer.gov/seerstat/WebHelp/seerstat.htm for the methodology used in the calculation of the age-adjusted 95% confidence intervals.

3. See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.

N.B. In areas with small Latino populations, methods to indirectly identify Latinos (like NHIA) can overestimate the ethnicity-specific counts of cancer cases. Also, even small errors in Latino population estimates can affect the magnitude of the cancer rates.

**Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates[1] and 95% Confidence Intervals[2]**
**Selected Areas in the United States[3], 1999-2003, Hispanic/Latino, All Races, Males**

**Colon and Rectum**




1 Rates are per 100,000 and are adjusted to the 2000 U.S. population standard. Rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the rates for the U.S. combined.

2. See http://seer.cancer.gov/seerstat/WebHelp/seerstat.htm for the methodology used in the calculation of the age-adjusted 95% confidence intervals.

3. See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.

N.B. In areas with small Latino populations, methods to indirectly identify Latinos (like NHIA) can overestimate the ethnicity-specific counts of cancer cases. Also, even small errors in Latino population estimates can affect the magnitude of the cancer rates.

**Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates[1] and 95% Confidence Intervals[2]**
**Selected Areas in the United States[3], 1999-2003, Hispanic/Latina, All Races, Females**

**Colon and Rectum**




[1] Rates are per 100,000 and are adjusted to the 2000 U.S. population standard. Rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the rates for the U.S. combined.

[2] See http://seer.cancer.gov/seerstat/WebHelp/seerstat.htm for the methodology used in the calculation of the age-adjusted 95% confidence intervals.

[3] See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.

N.B. In areas with small Latino populations, methods to indirectly identify Latinos (like NHIA) can overestimate the ethnicity-specific counts of cancer cases. Also, even small errors in Latino population estimates can affect the magnitude of the cancer rates.

This Page Intentionally Left Blank

**Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates[1] and 95% Confidence Intervals[2]**
**Selected Areas in the United States[3], 1999-2003, Hispanic/Latina, All Races, Females**

**Corpus Uteri and Uterus, NOS**




1 Rates are per 100,000 and are adjusted to the 2000 U.S. population standard. Rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the rates for the U.S. combined.

2. See http://seer.cancer.gov/seerstat/WebHelp/seerstat.htm for the methodology used in the calculation of the age-adjusted 95% confidence intervals.

3. See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.

N.B. In areas with small Latino populations, methods to indirectly identify Latinos (like NHIA) can overestimate the ethnicity-specific counts of cancer cases. Also, even small errors in Latino population estimates can affect the magnitude of the cancer rates.

**Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates[1] and 95% Confidence Intervals[2]**
**Selected Areas in the United States[3], 1999-2003, Hispanic/Latino, All Races, Males**

**Gallbladder**




1 Rates are per 100,000 and are adjusted to the 2000 U.S. population standard. Rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the rates for the U.S. combined.

2. See http://seer.cancer.gov/seerstat/WebHelp/seerstat.htm for the methodology used in the calculation of the age-adjusted 95% confidence intervals.

3. See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.

N.B. In areas with small Latino populations, methods to indirectly identify Latinos (like NHIA) can overestimate the ethnicity-specific counts of cancer cases. Also, even small errors in Latino population estimates can affect the magnitude of the cancer rates.

**Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates[1] and 95% Confidence Intervals[2]**
**Selected Areas in the United States[3], 1999-2003, Hispanic/Latina, All Races, Females**




1 Rates are per 100,000 and are adjusted to the 2000 U.S. population standard. Rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the rates for the U.S. combined.

2. See http://seer.cancer.gov/seerstat/WebHelp/seerstat.htm for the methodology used in the calculation of the age-adjusted 95% confidence intervals.

3. See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.

N.B. In areas with small Latino populations, methods to indirectly identify Latinos (like NHIA) can overestimate the ethnicity-specific counts of cancer cases. Also, even small errors in Latino population estimates can affect the magnitude of the cancer rates.

**Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates[1] and 95% Confidence Intervals[2]**
**Selected Areas in the United States[3], 1999-2003, Hispanic/Latino, All Races, Males**

**Kidney and Renal Pelvis**




1 Rates are per 100,000 and are adjusted to the 2000 U.S. population standard. Rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the rates for the U.S. combined.

2. See http://seer.cancer.gov/seerstat/WebHelp/seerstat.htm for the methodology used in the calculation of the age-adjusted 95% confidence intervals.

3. See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.

N.B. In areas with small Latino populations, methods to indirectly identify Latinos (like NHIA) can overestimate the ethnicity-specific counts of cancer cases. Also, even small errors in Latino population estimates can affect the magnitude of the cancer rates.

**Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates[1] and 95% Confidence Intervals[2]**
**Selected Areas in the United States[3], 1999-2003, Hispanic/Latina, All Races, Females**

**Kidney and Renal Pelvis**




1 Rates are per 100,000 and are adjusted to the 2000 U.S. population standard. Rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the rates for the U.S. combined.

2. See http://seer.cancer.gov/seerstat/WebHelp/seerstat.htm for the methodology used in the calculation of the age-adjusted 95% confidence intervals.

3. See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.

N.B. In areas with small Latino populations, methods to indirectly identify Latinos (like NHIA) can overestimate the ethnicity-specific counts of cancer cases. Also, even small errors in Latino population estimates can affect the magnitude of the cancer rates.

**Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates[1] and 95% Confidence Intervals[2]**
**Selected Areas in the United States[3], 1999-2003, Hispanic/Latino, All Races, Males**

**Leukemias**




1 Rates are per 100,000 and are adjusted to the 2000 U.S. population standard. Rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the rates for the U.S. combined.

2. See http://seer.cancer.gov/seerstat/WebHelp/seerstat.htm for the methodology used in the calculation of the age-adjusted 95% confidence intervals.

3. See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.

N.B. In areas with small Latino populations, methods to indirectly identify Latinos (like NHIA) can overestimate the ethnicity-specific counts of cancer cases. Also, even small errors in Latino population estimates can affect the magnitude of the cancer rates.

**Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates[1] and 95% Confidence Intervals[2]**
**Selected Areas in the United States[3], 1999-2003, Hispanic/Latina, All Races, Females**




1 Rates are per 100,000 and are adjusted to the 2000 U.S. population standard. Rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the rates for the U.S. combined.

2. See http://seer.cancer.gov/seerstat/WebHelp/seerstat.htm for the methodology used in the calculation of the age-adjusted 95% confidence intervals.

3. See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.

N.B. In areas with small Latino populations, methods to indirectly identify Latinos (like NHIA) can overestimate the ethnicity-specific counts of cancer cases. Also, even small errors in Latino population estimates can affect the magnitude of the cancer rates.

**Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates[1] and 95% Confidence Intervals[2]**
**Selected Areas in the United States[3], 1999-2003, Hispanic/Latino, All Races, Males**




1 Rates are per 100,000 and are adjusted to the 2000 U.S. population standard. Rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the rates for the U.S. combined.

2. See http://seer.cancer.gov/seerstat/WebHelp/seerstat.htm for the methodology used in the calculation of the age-adjusted 95% confidence intervals.

3. See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.

N.B. In areas with small Latino populations, methods to indirectly identify Latinos (like NHIA) can overestimate the ethnicity-specific counts of cancer cases. Also, even small errors in Latino population estimates can affect the magnitude of the cancer rates.

**Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates[1] and 95% Confidence Intervals[2]**
**Selected Areas in the United States[3], 1999-2003, Hispanic/Latina, All Races, Females**

**Liver**




1 Rates are per 100,000 and are adjusted to the 2000 U.S. population standard. Rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the rates for the U.S. combined.

2. See http://seer.cancer.gov/seerstat/WebHelp/seerstat.htm for the methodology used in the calculation of the age-adjusted 95% confidence intervals.

3. See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.

N.B. In areas with small Latino populations, methods to indirectly identify Latinos (like NHIA) can overestimate the ethnicity-specific counts of cancer cases. Also, even small errors in Latino population estimates can affect the magnitude of the cancer rates.

**Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates[1] and 95% Confidence Intervals[2]**
**Selected Areas in the United States[3], 1999-2003, Hispanic/Latino, All Races, Males**

**Lung and Bronchus**




1 Rates are per 100,000 and are adjusted to the 2000 U.S. population standard. Rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the rates for the U.S. combined.

2. See http://seer.cancer.gov/seerstat/WebHelp/seerstat.htm for the methodology used in the calculation of the age-adjusted 95% confidence intervals.

3. See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.

N.B. In areas with small Latino populations, methods to indirectly identify Latinos (like NHIA) can overestimate the ethnicity-specific counts of cancer cases. Also, even small errors in Latino population estimates can affect the magnitude of the cancer rates.

**Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates[1] and 95% Confidence Intervals[2]**
**Selected Areas in the United States[3], 1999-2003, Hispanic/Latina, All Races, Females**

**Lung and Bronchus**




1 Rates are per 100,000 and are adjusted to the 2000 U.S. population standard. Rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the rates for the U.S. combined.

2. See http://seer.cancer.gov/seerstat/WebHelp/seerstat.htm for the methodology used in the calculation of the age-adjusted 95% confidence intervals.

3. See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.

N.B. In areas with small Latino populations, methods to indirectly identify Latinos (like NHIA) can overestimate the ethnicity-specific counts of cancer cases. Also, even small errors in Latino population estimates can affect the magnitude of the cancer rates.

**Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates[1] and 95% Confidence Intervals[2]**
**Selected Areas in the United States[3], 1999-2003, Hispanic/Latino, All Races, Males**

**Lymphomas**




1 Rates are per 100,000 and are adjusted to the 2000 U.S. population standard. Rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the rates for the U.S. combined.

2. See http://seer.cancer.gov/seerstat/WebHelp/seerstat.htm for the methodology used in the calculation of the age-adjusted 95% confidence intervals.

3. See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.

N.B. In areas with small Latino populations, methods to indirectly identify Latinos (like NHIA) can overestimate the ethnicity-specific counts of cancer cases. Also, even small errors in Latino population estimates can affect the magnitude of the cancer rates.

**Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates[1] and 95% Confidence Intervals[2]**
**Selected Areas in the United States[3], 1999-2003, Hispanic/Latina, All Races, Females**

**Lymphomas**




1 Rates are per 100,000 and are adjusted to the 2000 U.S. population standard. Rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the rates for the U.S. combined.

2. See http://seer.cancer.gov/seerstat/WebHelp/seerstat.htm for the methodology used in the calculation of the age-adjusted 95% confidence intervals.

3. See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.

N.B. In areas with small Latino populations, methods to indirectly identify Latinos (like NHIA) can overestimate the ethnicity-specific counts of cancer cases. Also, even small errors in Latino population estimates can affect the magnitude of the cancer rates.

This Page Intentionally Left Blank

**Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates[1] and 95% Confidence Intervals[2]**
**Selected Areas in the United States[3], 1999-2003, Hispanic/Latina, All Races, Females**




1 Rates are per 100,000 and are adjusted to the 2000 U.S. population standard. Rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the rates for the U.S. combined.

2. See http://seer.cancer.gov/seerstat/WebHelp/seerstat.htm for the methodology used in the calculation of the age-adjusted 95% confidence intervals.

3. See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.

N.B. In areas with small Latino populations, methods to indirectly identify Latinos (like NHIA) can overestimate the ethnicity-specific counts of cancer cases. Also, even small errors in Latino population estimates can affect the magnitude of the cancer rates.

**Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates[1] and 95% Confidence Intervals[2]**
**Selected Areas in the United States[3], 1999-2003, Hispanic/Latino, All Races, Males**




1 Rates are per 100,000 and are adjusted to the 2000 U.S. population standard. Rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the rates for the U.S. combined.

2. See http://seer.cancer.gov/seerstat/WebHelp/seerstat.htm for the methodology used in the calculation of the age-adjusted 95% confidence intervals.

3. See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.

N.B. In areas with small Latino populations, methods to indirectly identify Latinos (like NHIA) can overestimate the ethnicity-specific counts of cancer cases. Also, even small errors in Latino population estimates can affect the magnitude of the cancer rates.

**Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates[1] and 95% Confidence Intervals[2]**
**Selected Areas in the United States[3], 1999-2003, Hispanic/Latina, All Races, Females**




1 Rates are per 100,000 and are adjusted to the 2000 U.S. population standard. Rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the rates for the U.S. combined.

2. See http://seer.cancer.gov/seerstat/WebHelp/seerstat.htm for the methodology used in the calculation of the age-adjusted 95% confidence intervals.

3. See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.

N.B. In areas with small Latino populations, methods to indirectly identify Latinos (like NHIA) can overestimate the ethnicity-specific counts of cancer cases. Also, even small errors in Latino population estimates can affect the magnitude of the cancer rates.

**Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates[1] and 95% Confidence Intervals[2]**
**Selected Areas in the United States[3], 1999-2003, Hispanic/Latino, All Races, Males**




[1] Rates are per 100,000 and are adjusted to the 2000 U.S. population standard. Rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the rates for the U.S. combined.

[2] See http://seer.cancer.gov/seerstat/WebHelp/seerstat.htm for the methodology used in the calculation of the age-adjusted 95% confidence intervals.

[3] See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.

N.B. In areas with small Latino populations, methods to indirectly identify Latinos (like NHIA) can overestimate the ethnicity-specific counts of cancer cases. Also, even small errors in Latino population estimates can affect the magnitude of the cancer rates.

This Page Intentionally Left Blank

**Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates[1] and 95% Confidence Intervals[2]**
**Selected Areas in the United States[3], 1999-2003, Hispanic/Latino, All Races, Males**




1 Rates are per 100,000 and are adjusted to the 2000 U.S. population standard. Rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the rates for the U.S. combined.

2. See http://seer.cancer.gov/seerstat/WebHelp/seerstat.htm for the methodology used in the calculation of the age-adjusted 95% confidence intervals.

3. See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.

N.B. In areas with small Latino populations, methods to indirectly identify Latinos (like NHIA) can overestimate the ethnicity-specific counts of cancer cases. Also, even small errors in Latino population estimates can affect the magnitude of the cancer rates.

**Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates[1] and 95% Confidence Intervals[2]**
**Selected Areas in the United States[3], 1999-2003, Hispanic/Latina, All Races, Females**




1 Rates are per 100,000 and are adjusted to the 2000 U.S. population standard. Rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the rates for the U.S. combined.

2. See http://seer.cancer.gov/seerstat/WebHelp/seerstat.htm for the methodology used in the calculation of the age-adjusted 95% confidence intervals.

3. See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.

N.B. In areas with small Latino populations, methods to indirectly identify Latinos (like NHIA) can overestimate the ethnicity-specific counts of cancer cases. Also, even small errors in Latino population estimates can affect the magnitude of the cancer rates.

**Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates[1] and 95% Confidence Intervals[2]**
**Selected Areas in the United States[3], 1999-2003, Hispanic/Latino, All Races, Males**




1 Rates are per 100,000 and are adjusted to the 2000 U.S. population standard. Rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the rates for the U.S. combined.

2. See http://seer.cancer.gov/seerstat/WebHelp/seerstat.htm for the methodology used in the calculation of the age-adjusted 95% confidence intervals.

3. See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.

N.B. In areas with small Latino populations, methods to indirectly identify Latinos (like NHIA) can overestimate the ethnicity-specific counts of cancer cases. Also, even small errors in Latino population estimates can affect the magnitude of the cancer rates.

This Page Intentionally Left Blank

**Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates[1] and 95% Confidence Intervals[2]**
**Selected Areas in the United States[3], 1999-2003, Hispanic/Latino, All Races, Males**

**Thyroid**




1 Rates are per 100,000 and are adjusted to the 2000 U.S. population standard. Rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the rates for the U.S. combined.

2. See http://seer.cancer.gov/seerstat/WebHelp/seerstat.htm for the methodology used in the calculation of the age-adjusted 95% confidence intervals.

3. See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.

N.B. In areas with small Latino populations, methods to indirectly identify Latinos (like NHIA) can overestimate the ethnicity-specific counts of cancer cases. Also, even small errors in Latino population estimates can affect the magnitude of the cancer rates.

**Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates[1] and 95% Confidence Intervals[2]**
**Selected Areas in the United States[3], 1999-2003, Hispanic/Latina, All Races, Females**

**Thyroid**




1 Rates are per 100,000 and are adjusted to the 2000 U.S. population standard. Rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the rates for the U.S. combined.

2. See http://seer.cancer.gov/seerstat/WebHelp/seerstat.htm for the methodology used in the calculation of the age-adjusted 95% confidence intervals.

3. See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.

N.B. In areas with small Latino populations, methods to indirectly identify Latinos (like NHIA) can overestimate the ethnicity-specific counts of cancer cases. Also, even small errors in Latino population estimates can affect the magnitude of the cancer rates.

**Average Annual Age-adjusted (2000 U.S. Standard) U.S. Cancer Incidence Rates[1] (NAACCR Combined[2]) for Selected Cancer Sites by Race/Ethnicity, Males, 1999-2003**




1. Rates are per 100,000 population and were age-adjusted by five-year age groups to the 2000 U.S. standard population based on single years of age. Rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the rates for the U.S. combined.
2. See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.

N.B. In areas with small Latino populations, methods to indirectly identify Latinos (like NHIA) can overestimate the ethnicity-specific counts of cancer cases. Also, even small errors in Latino population estimates can affect the magnitude of the cancer rates.

**Average Annual Age-adjusted (2000 U.S. Standard) U.S. Cancer Incidence Rates[1] (NAACCR Combined[2]) for Selected Cancer Sites by Race/Ethnicity, Females, 1999-2003**




1. Rates are per 100,000 population and were age-adjusted by five-year age groups to the 2000 U.S. standard population based on single years of age. Rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the rates for the U.S. combined.
2. See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.

N.B. In areas with small Latino populations, methods to indirectly identify Latinos (like NHIA) can overestimate the ethnicity-specific counts of cancer cases. Also, even small errors in Latino population estimates can affect the magnitude of the cancer rates.