Section III

Average-annual Age-adjusted Cancer Incidence for the NAACCR (U.S.) Combined by Sex and Hispanic/Latino Ethnicity



Population Indicators:

Combined Population, 1998-2002:

Hispanic/Latino	135,708,450
Non-Hispanic White	772,317,547
Non-Hispanic Black	120,754,669

Number of Cases in the Combined Cancer Statistics:

Hispanic/Latino	261,827
Non-Hispanic White	4,313,275
Non-Hispanic Black	436,644

Percentage of U.S. Population Reflected in the Combined Cancer Statistics:

Hispanic/Latino	76.5%
Non-Hispanic White	78.2%
Non-Hispanic Black	68.7%

Demographic Information

Hispanic/Latino Population 2000: 26,979,057

% Distribution of Hispanic/Latino Population 2000 Census Data



% Hispanic/Latino of Total Population



Site-Specific Microscopic Confirmation (1998-2002)

Cancer Site	% Micro. Confirm	Cancer Site	% Micro. Confirm
All Sites	93.6	Ovary	91.9
Buccal Cavity & Pharynx	97.6	Prostate	96.6
Esophagus	94.5	Testis	99.2
Stomach	95.5	Bladder (incl. *in situ*)	98.2
Colon & Rectum	96.4	Kidney & Renal Pelvis	90.4
Liver	73.3	Brain & Other Nervous System	85.5
Pancreas	74.3	Thyroid Gland	99.2
Lung & Bronchus	88.0	Hodgkin Disease	98.8
Melanomas of the Skin	98.8	Non-Hodgkin Lymphoma	96.6
Female Breast (excl. *in situ*)	98.0	Multiple Myeloma	88.0
Uterus	98.2	Leukemias	87.8
Cervix uteri	97.6	Breast *in situ*	100.0
Corpus uteri & Uterus, NOS	98.4		

Registries Included in the NAACCR U.S. Combined Incidence 1998-2002



POPULATIONS FOR NAACCR (U.S.) COMBINED REGISTRIES

1998

Age	Males Hispanic	Males Non-Hisp White	Males Non-Hisp Black	Females Hispanic	Females Non-Hisp White	Females Non-Hisp Black
< 1	276464	915731	192914	262941	867107	187739
1-4	1090460	3785161	833483	1041178	3594546	808853
5-9	1306080	5182783	1174544	1251838	4907908	1137193
10-14	1111992	5205874	1048312	1063477	4923823	1017371
15-19	1177970	5172332	1014172	1045390	4923774	991456
20-24	1250907	4521304	839980	1063869	4414738	883801
25-29	1305280	5023938	845110	1106846	4975508	932188
30-34	1222228	5573996	868836	1077600	5499916	968502
35-39	1042363	6421027	905968	967044	6386284	1022135
40-44	814501	6304804	823024	791456	6322255	940349
45-49	602092	5654075	665970	613737	5721752	782781
50-54	445570	4857217	504141	474108	4981549	603665
55-59	321267	3797259	385097	356002	3995507	477165
60-64	249637	3118001	311949	289012	3384019	398170
65-69	198063	2904608	264092	242844	3336732	354338
70-74	145671	2649979	197343	191949	3350300	295911
75-79	91665	2069627	142307	132597	2942720	231763
80-84	46863	1260866	73652	80950	2138836	150461
85+	33954	814180	53568	71463	2038605	143383
Total	12733027	75232762	11144462	12124301	78705879	12327224

2001

Age	Males Hispanic	Males Non-Hisp White	Males Non-Hisp Black	Females Hispanic	Females Non-Hisp White	Females Non-Hisp Black
< 1	320865	915838	223533	308103	873480	217337
1-4	1162069	3639773	819424	1110969	3457382	790645
5-9	1453965	4907193	1130381	1389753	4646779	1092024
10-14	1312520	5274401	1152309	1255355	4992075	1116641
15-19	1260780	5197535	1037534	1137560	4933776	1007160
20-24	1483148	4794043	928628	1209328	4700838	968851
25-29	1452540	4532923	819495	1229450	4453423	899461
30-34	1351783	5275699	859795	1179871	5199847	954036
35-39	1200225	5952871	904320	1091934	5924636	1012214
40-44	971172	6408558	887240	927147	6419652	1000994
45-49	735945	6020498	748800	737755	6079596	866943
50-54	552412	5460823	610785	579289	5580443	723789
55-59	386245	4243700	429233	426048	4440493	529013
60-64	287699	3286780	334605	330527	3553815	427730
65-69	218541	2803950	268648	269734	3179865	362740
70-74	169635	2590895	203357	221302	3187576	302557
75-79	113147	2107586	147488	159403	2961969	243856
80-84	60173	1365915	84057	97977	2240065	162305
85+	42243	930573	58763	86831	2189929	156285
Total	14535107	75709554	11648395	13748336	79015639	12834581

1999

Age	Males Hispanic	Males Non-Hisp White	Males Non-Hisp Black	Females Hispanic	Females Non-Hisp White	Females Non-Hisp Black
< 1	290705	904774	198468	276802	859319	192237
1-4	1112637	3728901	827418	1064312	3540274	801240
5-9	1377669	5127711	1175784	1318525	4855331	1137332
10-14	1173589	5244311	1089246	1122686	4959848	1056511
15-19	1227611	5186896	1022507	1087188	4946148	1000038
20-24	1336014	4573312	858929	1118126	4473170	907134
25-29	1343350	4866698	842294	1149396	4806579	930311
30-34	1255372	5400806	857485	1102664	5328422	951915
35-39	1092927	6317375	912976	1012182	6288535	1025574
40-44	861922	6357334	849899	833510	6368389	966294
45-49	640531	5775615	696196	651063	5838430	810942
50-54	479307	5072618	539613	507746	5192874	642380
55-59	336573	3937039	398071	373402	4136658	491877
60-64	262330	3187573	321666	304220	3453386	410354
65-69	201822	2848779	262636	248879	3251927	352926
70-74	152213	2635834	201976	201646	3303779	301038
75-79	98507	2098637	143283	139927	2973861	235782
80-84	49221	1291561	76750	83759	2166432	153595
85+	35509	848011	55025	74854	2089069	148555
Total	13327809	75403785	11330222	12670887	78832431	12516035

2002

Age	Males Hispanic	Males Non-Hisp White	Males Non-Hisp Black	Females Hispanic	Females Non-Hisp White	Females Non-Hisp Black
< 1	328674	898025	221603	316614	856817	214863
1-4	1204773	3636621	835850	1152058	3457685	808484
5-9	1469140	4793259	1097736	1402611	4540032	1061007
10-14	1373095	5255571	1175627	1312332	4977264	1138895
15-19	1260299	5210844	1050558	1155279	4932846	1017507
20-24	1528568	4919360	964166	1240613	4819962	993166
25-29	1511111	4459130	816601	1261548	4376076	893928
30-34	1413260	5209186	865778	1227567	5125470	958657
35-39	1246190	5741998	890772	1124743	5711139	996841
40-44	1030201	6387195	895881	973539	6395441	1011009
45-49	785344	6112117	775241	781673	6170386	895133
50-54	587436	5519344	634526	612546	5637265	754289
55-59	416074	4475213	449355	455770	4668320	552914
60-64	302743	3432770	343542	347533	3705704	442021
65-69	228245	2800400	272703	280561	3166776	367316
70-74	176442	2552204	204141	228859	3118710	302735
75-79	120477	2103015	149043	169174	2929272	247250
80-84	66957	1409870	88277	106436	2287957	167888
85+	45861	974130	61085	93459	2238879	159524
Total	15094890	75890252	11792485	14242915	79116001	12983427

2000

Age	Males Hispanic	Males Non-Hisp White	Males Non-Hisp Black	Females Hispanic	Females Non-Hisp White	Females Non-Hisp Black
< 1	296048	913844	204719	281341	866775	196925
1-4	1146670	3680218	829164	1098646	3494287	799834
5-9	1429682	5019650	1158583	1366997	4754096	1119913
10-14	1251277	5270690	1123646	1197777	4983432	1090041
15-19	1267091	5179539	1028407	1124083	4941453	1004496
20-24	1424770	4657865	888119	1175043	4565969	936107
25-29	1395447	4688544	828882	1194968	4614961	914003
30-34	1295809	5312440	857079	1136090	5242646	949534
35-39	1144601	6136719	909726	1054352	6117327	1020287
40-44	914786	6389267	872716	880742	6399424	985507
45-49	688316	5898691	724400	694422	5959283	838501
50-54	517370	5312977	579981	545657	5431712	687209
55-59	359309	4062392	412855	397592	4263207	508651
60-64	274472	3220651	329704	318062	3485402	419059
65-69	209654	2824115	262863	258994	3214296	355443
70-74	161256	2617158	204113	212202	3244255	304800
75-79	105504	2117573	145157	149486	2994297	238763
80-84	53910	1318056	80394	89642	2188226	157817
85+	38744	886465	56782	80366	2143342	153658
Total	13974716	75506854	11497290	13256462	78904390	12680548

Total Cancer Cases and Average Annual Age-adjusted Cancer Incidence Rates*
NAACCR (U.S.) Combined, 1998-2002, Hispanic/Latino, All Races

	MALES (HISPANIC/LATINO)					FEMALES (HISPANIC/LATINA)				
SITES	*Inc.*	*Adj. Rate 2000 US*	*S.E*	*Adj. Rate World*	*S.E.*	*Inc.*	*Adj. Rate 2000 US*	*S.E.*	*Adj. Rate World*	*S.E.*
All Sites	130760	445.6	1.37	285.7	0.82	131067	328.4	0.95	225.5	0.64
Buccal Cavity & Pharynx	3849	12.0	0.21	8.3	0.14	1688	4.4	0.11	2.9	0.07
Lip	192	0.7	0.06	0.4	0.03	71	0.2	0.02	0.1	0.01
Tongue	928	2.9	0.10	2.0	0.07	420	1.1	0.06	0.7	0.04
Major salivary gland	347	1.0	0.07	0.7	0.04	320	0.7	0.04	0.5	0.03
Floor of mouth	281	0.9	0.06	0.6	0.04	88	0.3	0.03	0.2	0.02
Gum and other mouth	487	1.6	0.08	1.1	0.05	392	1.1	0.06	0.7	0.03
Nasopharynx	279	0.7	0.05	0.6	0.04	133	0.3	0.03	0.2	0.02
Oropharynx	192	0.6	0.05	0.4	0.03	40	0.1	0.02	0.1	0.01
Hypopharynx	384	1.3	0.07	0.9	0.05	50	0.1	0.02	0.1	0.01
Digestive System	30024	108.2	0.69	66.8	0.40	25629	73.4	0.47	44.5	0.29
Esophagus	1626	5.9	0.16	3.7	0.09	524	1.6	0.07	0.9	0.04
Stomach	4408	16.1	0.27	9.6	0.15	3235	9.1	0.17	5.5	0.10
Small intestine	483	1.6	0.08	1.0	0.05	455	1.2	0.06	0.8	0.04
Colon excl. rectum	9685	36.9	0.41	21.8	0.23	9872	28.7	0.30	17.0	0.18
Rectum & rectosigmoid	4987	17.1	0.27	11.1	0.16	3610	9.8	0.17	6.4	0.11
Anus	384	1.1	0.06	0.8	0.04	621	1.7	0.07	1.1	0.05
Liver	3925	12.6	0.22	8.6	0.14	1570	4.5	0.12	2.8	0.07
Gallbladder	360	1.4	0.08	0.8	0.04	1080	3.1	0.10	2.0	0.06
Pancreas	2968	11.2	0.22	6.7	0.13	3217	9.6	0.17	5.6	0.10
Respiratory System	16322	61.4	0.52	37.3	0.30	10081	28.6	0.29	18.2	0.19
Larynx	1988	6.8	0.16	4.6	0.10	380	1.0	0.05	0.7	0.04
Lung & bronchus	13843	53.4	0.49	31.9	0.28	9427	27.0	0.29	17.0	0.18
Bones & Joints	621	1.0	0.05	0.9	0.04	498	0.8	0.04	0.8	0.03
Soft Tissue (incl. heart)	1334	3.0	0.10	2.3	0.07	1244	2.5	0.08	2.0	0.06
Skin (excl. basal and sq.)	1893	5.6	0.15	3.6	0.09	2232	4.9	0.11	3.6	0.08
Melanomas of the skin	1543	4.7	0.14	3.0	0.08	1894	4.1	0.10	3.0	0.07
Breast (excl. *in situ*)	233	0.8	0.06	0.5	0.03	38844	94.5	0.50	68.3	0.36
Female Genital System	-	-	-	-	-	20592	47.9	0.35	35.0	0.25
Cervix uteri	-	-	-	-	-	7186	14.6	0.18	11.3	0.14
Corpus uteri	-	-	-	-	-	7185	17.9	0.22	13.0	0.16
Uterus, NOS	-	-	-	-	-	313	0.8	0.05	0.5	0.03
Ovary	-	-	-	-	-	4692	11.4	0.17	8.1	0.12
Male Genital System	40927	147.4	0.79	95.9	0.49	-	-	-	-	-
Prostate	37752	142.6	0.78	91.7	0.48	-	-	-	-	-
Testis	2719	3.4	0.07	3.3	0.06	-	-	-	-	-
Penis	387	1.2	0.07	0.8	0.04	-	-	-	-	-
Urinary System	10733	38.8	0.41	23.7	0.24	5453	14.8	0.21	9.6	0.13
Bladder (incl. *in situ*)	5577	22.5	0.33	12.6	0.17	2057	6.1	0.14	3.5	0.08
Kidney & renal pelvis	4944	15.5	0.24	10.6	0.16	3261	8.3	0.15	5.8	0.10
Ureter	123	0.5	0.05	0.3	0.03	95	0.3	0.03	0.2	0.02
Eye	292	0.6	0.05	0.5	0.03	257	0.5	0.03	0.4	0.03
Brain & Other Nervous System	2869	6.2	0.14	5.0	0.10	2502	5.0	0.11	4.1	0.09
Brain	2692	5.9	0.14	4.7	0.10	2293	4.6	0.10	3.8	0.08
Endocrine System	1742	3.8	0.10	3.0	0.08	6006	11.3	0.16	9.3	0.12
Thyroid gland	1413	3.2	0.10	2.5	0.07	5724	10.8	0.15	8.8	0.12
Lymphomas	8628	22.9	0.29	15.9	0.18	6767	16.3	0.21	11.3	0.14
Hodgkin's disease	1613	3.0	0.09	2.5	0.07	1159	2.0	0.06	1.7	0.05
Non-Hodgkin's lymphoma	7015	20.0	0.28	13.4	0.17	5608	14.3	0.20	9.6	0.13
Multiple Myeloma	1751	6.3	0.17	3.9	0.09	1669	4.7	0.12	3.0	0.08
Leukemias	4982	12.2	0.21	9.0	0.13	3969	8.2	0.14	6.4	0.11
Acute lymphocytic	1686	2.1	0.06	2.5	0.06	1256	1.7	0.05	2.0	0.06
Chronic lymphocytic	669	2.7	0.11	1.5	0.06	469	1.4	0.07	0.8	0.04
Acute myeloid	1395	3.8	0.12	2.6	0.07	1316	2.9	0.09	2.1	0.06
Chronic myeloid	681	1.8	0.08	1.2	0.05	467	1.1	0.05	0.7	0.04
Other	551	1.8	0.09	1.1	0.05	461	1.1	0.06	0.7	0.04
Mesothelioma	402	1.5	0.08	0.9	0.05	135	0.4	0.03	0.2	0.02
Kaposi Sarcoma	1041	1.9	0.07	1.4	0.05	111	0.3	0.03	0.2	0.02
Ill-defined & Unknown	3117	11.9	0.24	6.8	0.13	3390	9.9	0.18	5.7	0.10
Breast *in situ* (not in All Sites)	42	0.1	0.02	0.1	0.01	8037	19.4	0.22	14.6	0.17

* Rates are per 100,000 population and were age-adjusted by five-year age groups to the 2000 U.S. standard population based on single years of age and to the World population standard.

- Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for All Sites combined.

N.B. In areas with small Latino populations, methods to indirectly identify Latinos (like NHIA) can overestimate the ethnicity-specific counts of cancer cases. Also, even small errors in Latino population estimates can affect the magnitude of the cancer rates.

Total Cancer Cases and Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates[1] for Selected Cancer Sites[2]
Selected Areas in the United States[3], 1998-2002, Hispanic/Latino, All Races, Males
{Page 1 of 4}

	All Sites			Bladder			Brain & ONS[4]			Colon & Rectum			Gallbladder			Kidney & Renal Pelvis		
	Cases	*Rate*	*S.E.*[4]	*Cases*	*Rate*	*S.E.*	*Cases*	*Rate*	*S.E.*	*Cases*	*Rate*	*S.E.*	*Cases*	*Rate*	*S.E.*	*Cases*	*Rate*	*S.E.*
NAACCR U.S. Combined	**130760**	**445.6**	**1.37**	**5577**	**22.5**	**0.33**	**2869**	**6.2**	**0.14**	**14672**	**54.0**	**0.49**	**360**	**1.4**	**0.08**	**4944**	**15.5**	**0.24**
Alabama	230	325.0	24.59	-	-	-	10	4.9	1.81	22	39.7	9.26	-	-	-	13	18.4	5.79
Alaska	86	480.5	77.24	-	-	-	-	-	-	7	39.4	16.98	-	-	-	7	16.2	6.63
Arizona	4932	383.7	6.20	187	16.8	1.34	131	6.2	0.69	535	47.2	2.28	22	1.9	0.43	263	19.2	1.34
California	42091	395.6	2.18	1555	17.9	0.50	963	5.2	0.21	4410	44.7	0.75	125	1.4	0.14	1731	15.0	0.40
CA: Greater Bay Area	4872	329.5	5.30	182	15.3	1.25	98	3.9	0.46	526	39.5	1.91	13	0.8	0.25	181	11.1	0.92
CA: Los Angeles	16810	412.9	3.59	581	17.6	0.80	382	5.3	0.34	1654	43.9	1.19	47	1.3	0.21	656	14.8	0.65
Colorado	3636	456.1	8.46	127	16.7	1.62	81	6.4	0.87	465	61.7	3.14	7	1.0	0.41	162	16.7	1.42
Connecticut	1564	534.9	15.80	77	30.5	3.99	38	8.9	1.86	172	59.9	5.25	-	-	-	59	16.8	2.55
Delaware	141	462.4	47.94	6	31.9	13.97	6	14.6	7.60	12	47.7	17.56	-	-	-	-	-	-
District of Columbia	157	344.3	32.70	-	-	-	-	-	-	10	22.2	8.07	-	-	-	7	12.4	4.89
Florida	25376	504.2	3.26	1471	31.4	0.85	424	7.2	0.36	3133	65.1	1.20	56	1.2	0.16	805	15.6	0.56
GA: Metropolitan Atlanta	550	376.9	22.24	21	27.0	6.68	17	4.7	1.57	46	34.5	6.93	-	-	-	20	15.9	4.97
Hawaii	297	362.3	22.81	16	22.6	5.86	7	6.2	2.89	36	45.6	8.00	-	-	-	10	8.6	3.00
Idaho	302	454.2	33.01	13	20.4	6.58	11	7.9	3.48	39	69.4	13.14	-	-	-	12	15.0	5.72
Illinois	5315	407.3	6.83	221	20.8	1.63	155	6.8	0.75	564	47.2	2.39	15	1.8	0.54	225	14.1	1.12
Indiana	821	422.1	16.94	33	19.0	3.95	15	5.0	1.46	92	48.6	5.56	-	-	-	53	22.4	3.62
Iowa	225	381.7	29.71	9	19.1	7.27	7	5.3	2.28	20	26.1	6.35	-	-	-	14	25.6	7.89
Kentucky	331	647.8	41.89	18	39.8	10.46	7	6.8	3.93	36	83.5	16.50	-	-	-	12	23.7	7.27
Louisiana	1126	707.6	22.72	52	37.5	5.71	18	11.2	3.04	127	84.0	8.02	-	-	-	36	20.3	3.55
Maine	54	658.9	96.44	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Massachusetts	1936	520.0	13.86	81	28.4	3.55	51	7.1	1.30	175	51.1	4.53	-	-	-	57	14.7	2.31
Michigan	1295	412.2	13.02	54	20.1	3.02	40	6.6	1.27	140	48.8	4.56	7	2.5	0.96	71	20.5	2.75
MI: Metropolitan Detroit	675	518.2	22.04	35	32.8	5.96	18	7.1	1.89	64	54.3	7.35	-	-	-	27	18.8	3.86
Minnesota	348	373.4	25.66	6	6.7	3.25	14	6.3	2.37	33	46.4	9.49	-	-	-	13	9.7	3.51
Missouri	545	471.7	23.06	17	17.8	4.90	13	5.3	1.77	82	72.5	9.17	-	-	-	23	16.7	4.06

N.B. In areas with small Latino populations, methods to indirectly identify Latinos (like NHIA) can overestimate the ethnicity-specific counts of cancer cases. Also, even small errors in Latino population estimates can affect the magnitude of the cancer rates.

Total Cancer Cases and Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates[1] for Selected Cancer Sites[2]
Selected Areas in the United States[3], 1998-2002, Hispanic/Latino, All Races, Males
{Page 2 of 4}

	All Sites			Bladder			Brain & ONS[4]			Colon & Rectum			Gallbladder			Kidney & Renal Pelvis		
	Cases	*Rate*	*S.E.*[4]	*Cases*	*Rate*	*S.E.*	*Cases*	*Rate*	*S.E.*	*Cases*	*Rate*	*S.E.*	*Cases*	*Rate*	*S.E.*	*Cases*	*Rate*	*S.E.*
Montana	104	644.9	74.49	-	-	-	-	-	-	10	76.4	28.91	-	-	-	-	-	-
Nebraska	316	434.8	29.46	14	18.4	5.35	-	-	-	48	74.8	12.23	-	-	-	14	18.2	5.80
Nevada	1315	398.6	13.62	62	24.0	3.74	48	8.2	1.64	129	43.8	4.61	-	-	-	66	19.4	2.83
New Jersey	6824	532.3	7.45	327	30.9	1.94	126	6.9	0.77	743	62.2	2.62	19	1.5	0.39	198	13.9	1.14
New Mexico	5011	416.6	6.12	209	19.2	1.38	92	6.3	0.69	628	53.5	2.21	20	2.0	0.46	234	18.3	1.24
New York	17429	499.2	4.29	623	22.1	0.99	356	7.0	0.45	2045	63.1	1.57	52	1.7	0.27	510	13.3	0.66
Ohio	1062	471.5	16.21	66	35.8	4.96	20	4.7	1.31	126	57.2	5.67	-	-	-	40	17.3	3.00
Oklahoma	571	439.9	22.37	31	28.9	5.87	17	6.6	2.01	63	53.6	8.07	-	-	-	20	13.3	3.60
Oregon	769	424.8	18.91	34	25.8	5.20	25	7.5	2.29	66	39.0	5.64	-	-	-	29	13.7	3.10
Pennsylvania	2353	617.5	14.51	106	32.5	3.52	48	8.2	1.51	281	81.9	5.47	6	1.5	0.66	76	18.7	2.53
Rhode Island	475	684.0	36.89	26	43.7	9.29	16	12.9	4.44	43	61.0	10.62	-	-	-	15	16.5	5.28
South Carolina	271	363.7	27.92	11	18.5	6.66	9	4.1	1.59	32	53.6	11.63	-	-	-	7	8.7	4.38
Utah	646	399.8	19.00	14	10.2	3.27	23	8.9	2.58	63	42.2	6.25	-	-	-	33	17.7	3.69
Washington	1443	458.5	14.82	49	18.4	3.02	48	6.6	1.38	144	52.8	5.08	-	-	-	72	19.7	2.88
WA: Seattle/Puget Sound	752	474.5	21.42	24	19.1	4.35	25	5.7	1.58	72	53.6	7.23	-	-	-	43	21.1	4.14
West Virginia	65	351.5	45.76	-	-	-	-	-	-	11	64.7	20.86	-	-	-	-	-	-
Wisconsin	748	476.9	20.75	24	19.2	4.34	22	6.0	1.54	81	53.9	6.72	-	-	-	31	15.2	3.05

1. Rates are per 100,000 population and were age-adjusted by five-year age groups to the 2000 U.S. standard population based on single years of age. Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for the U.S. combined.
2. Selected cancer sites by gender among the U.S. total population and the U.S. Hispanic/Latino population.
3. See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.
4. Abbreviations: S.E. - Standard Error; ONS - Other Nervous System

N.B. In areas with small Latino populations, methods to indirectly identify Latinos (like NHIA) can overestimate the ethnicity-specific counts of cancer cases. Also, even small errors in Latino population estimates can affect the magnitude of the cancer rates.

Total Cancer Cases and Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates[1] for Selected Cancer Sites[2]
Selected Areas in the United States[3], 1998-2002, Hispanic/Latino, All Races, Males
{Page 3 of 4}

	Liver			Lung & Bronchus			NH[4] Lymphoma			Prostate			Stomach			Testis		
	Cases	*Rate*	*S.E.*[4]	*Cases*	*Rate*	*S.E.*	*Cases*	*Rate*	*S.E.*	*Cases*	*Rate*	*S.E.*	*Cases*	*Rate*	*S.E.*	*Cases*	*Rate*	*S.E.*
NAACCR U.S. Combined	**3925**	**12.6**	**0.22**	**13843**	**53.4**	**0.49**	**7015**	**20.0**	**0.28**	**37752**	**142.6**	**0.78**	**4408**	**16.1**	**0.27**	**2719**	**3.4**	**0.07**
Alabama	7	9.5	4.36	33	53.5	9.96	14	11.6	3.61	59	99.6	13.59	8	13.2	5.61	6	1.7	0.70
Alaska	-	-	-	-	-	-	6	23.9	12.71	23	200.1	53.01	-	-	-	-	-	-
Arizona	180	13.2	1.07	566	52.1	2.39	265	17.7	1.28	1192	102.6	3.21	176	15.5	1.31	112	3.2	0.34
California	1430	12.7	0.37	3764	42.0	0.74	2343	18.4	0.45	12141	130.9	1.26	1617	16.7	0.46	1155	3.6	0.11
CA: Greater Bay Area	177	10.9	0.90	472	38.6	1.90	286	17.0	1.19	1338	101.7	2.93	189	13.8	1.12	139	3.0	0.28
CA: Los Angeles	555	13.4	0.62	1349	39.2	1.15	967	19.1	0.72	5193	146.6	2.17	685	18.7	0.80	449	3.6	0.18
Colorado	123	13.9	1.36	394	60.0	3.28	190	21.1	1.81	948	133.1	4.60	111	15.2	1.56	89	3.7	0.42
Connecticut	52	16.6	2.67	174	66.1	5.60	93	22.5	2.87	421	171.7	9.21	50	17.3	2.80	23	2.2	0.48
Delaware	-	-	-	15	50.7	14.69	9	17.9	6.98	37	155.6	28.89	-	-	-	9	10.2	3.74
District of Columbia	7	16.5	8.07	16	40.0	11.08	12	22.6	9.01	52	146.7	21.78	-	-	-	-	-	-
Florida	474	9.5	0.45	3263	67.1	1.21	1136	21.3	0.66	7811	155.7	1.80	623	12.8	0.53	273	3.5	0.21
GA: Metropolitan Atlanta	6	3.8	1.91	59	51.7	8.40	33	10.5	3.09	135	122.7	12.31	17	10.0	3.43	18	1.6	0.41
Hawaii	8	10.0	3.79	42	56.8	9.29	17	16.0	4.40	73	101.8	12.49	9	11.8	4.24	6	2.2	0.92
Idaho	7	6.8	3.04	24	44.0	10.87	26	41.0	11.01	68	127.4	17.93	15	19.2	6.77	-	-	-
Illinois	182	13.3	1.17	501	46.5	2.38	325	21.3	1.54	1391	129.9	3.95	188	15.5	1.39	152	3.2	0.34
Indiana	21	10.1	2.42	87	53.8	6.32	46	17.6	2.96	201	127.0	9.58	33	19.3	3.91	29	5.7	1.69
Iowa	10	23.7	8.25	21	43.5	10.69	15	20.9	5.82	45	95.2	15.12	8	15.0	5.76	-	-	-
Kentucky	-	-	-	62	141.9	19.59	20	33.4	9.65	70	166.7	21.74	-	-	-	9	3.0	1.05
Louisiana	33	23.2	4.41	170	107.9	8.71	51	29.9	4.59	323	209.2	12.30	23	15.5	3.58	15	4.6	1.21
Maine	-	-	-	9	103.0	35.06	-	-	-	19	260.2	62.74	-	-	-	-	-	-
Massachusetts	71	16.3	2.25	192	60.0	4.88	101	20.7	2.61	577	178.1	8.13	72	20.8	2.87	38	2.9	0.51
Michigan	41	10.9	1.81	144	54.3	4.99	60	17.9	2.73	371	133.5	7.47	44	13.4	2.28	36	3.5	0.62
MI: Metropolitan Detroit	16	10.6	2.77	80	69.6	8.37	33	20.8	4.29	210	174.6	12.77	20	15.4	3.82	14	3.8	1.10
Minnesota	17	13.5	4.11	38	61.7	11.49	20	18.4	5.88	80	113.5	14.04	16	19.2	6.05	15	2.5	0.65
Missouri	10	9.1	3.09	77	74.6	9.35	39	29.9	5.71	132	126.9	11.73	19	20.5	5.29	13	4.0	1.26

N.B. In areas with small Latino populations, methods to indirectly identify Latinos (like NHIA) can overestimate the ethnicity-specific counts of cancer cases. Also, even small errors in Latino population estimates can affect the magnitude of the cancer rates.

Total Cancer Cases and Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates[1] for Selected Cancer Sites[2]
Selected Areas in the United States[3], 1998-2002, Hispanic/Latino, All Races, Males
{Page 4 of 4}

	Liver			Lung & Bronchus			NH[4] Lymphoma			Prostate			Stomach			Testis		
	Cases	*Rate*	*S.E.*[4]	*Cases*	*Rate*	*S.E.*	*Cases*	*Rate*	*S.E.*	*Cases*	*Rate*	*S.E.*	*Cases*	*Rate*	*S.E.*	*Cases*	*Rate*	*S.E.*
Montana	-	-	-	13	77.0	22.38	9	53.3	22.20	30	193.2	39.43	-	-	-	-	-	-
Nebraska	10	16.5	6.31	43	67.0	11.73	17	22.6	6.98	69	109.8	14.53	12	28.9	9.18	8	5.4	3.66
Nevada	34	9.7	2.06	160	56.4	5.27	53	14.4	2.67	340	119.9	7.44	45	15.1	2.81	34	2.5	0.47
New Jersey	153	11.0	1.03	675	58.5	2.56	384	24.8	1.52	2281	196.7	4.57	250	18.6	1.35	121	3.5	0.33
New Mexico	159	12.3	1.01	530	48.2	2.16	182	14.1	1.09	1500	128.9	3.42	178	15.7	1.22	88	4.6	0.50
New York	611	15.9	0.71	1827	56.6	1.47	1051	23.3	0.85	5113	160.9	2.47	628	19.9	0.90	237	2.9	0.20
Ohio	32	15.6	3.10	123	56.4	5.49	62	25.8	3.88	290	144.7	9.12	24	9.3	2.02	21	3.6	0.85
Oklahoma	9	4.8	1.82	73	69.3	9.19	26	15.0	3.56	139	131.5	12.54	23	19.3	4.56	25	6.0	2.01
Oregon	21	11.1	2.97	97	72.4	8.21	53	20.9	3.80	187	129.7	10.40	19	13.0	3.71	31	3.6	0.71
Pennsylvania	94	19.1	2.15	277	82.3	5.48	126	26.8	2.87	643	186.6	7.98	75	23.0	2.99	32	3.1	0.58
Rhode Island	17	24.1	6.53	50	84.2	12.94	23	18.7	4.28	143	256.0	23.42	17	24.4	7.23	8	3.5	1.30
South Carolina	8	10.2	4.77	29	47.4	10.38	14	12.0	4.20	76	129.2	16.81	9	11.4	4.06	11	2.7	0.87
Utah	18	10.9	3.11	52	42.2	6.42	41	18.6	3.87	183	149.4	12.15	24	15.2	3.47	23	3.7	0.85
Washington	37	9.2	1.90	146	64.0	6.08	104	25.7	3.33	381	151.3	8.61	36	11.9	2.42	40	2.8	0.48
WA: Seattle/Puget Sound	27	14.6	3.58	68	57.7	8.23	52	23.6	4.19	196	163.2	13.05	14	8.6	2.81	21	2.6	0.64
West Virginia	-	-	-	10	64.6	21.05	-	-	-	17	99.4	24.55	-	-	-	-	-	-
Wisconsin	32	23.4	4.83	82	65.5	8.15	43	26.8	5.33	191	145.1	11.49	23	16.3	4.38	26	3.9	0.84

1. Rates are per 100,000 population and were age-adjusted by five-year age groups to the 2000 U.S. standard population based on single years of age. Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for the U.S. combined.
2. Selected cancer sites by gender among the U.S. total population and the U.S. Hispanic/Latino population.
3. See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.
4. Abbreviations: S.E. - Standard Error; NH - Non-Hodgkin.

N.B. In areas with small Latino populations, methods to indirectly identify Latinos (like NHIA) can overestimate the ethnicity-specific counts of cancer cases. Also, even small errors in Latino population estimates can affect the magnitude of the cancer rates.

Total Cancer Cases and Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates[1] for Selected Cancer Sites[2]
Selected Areas in the United States[3], 1998-2002, Hispanic/Latina, All Races, Females
{Page 1 of 4}

	All Sites			Breast			Cervix Uteri			Colon & Rectum			Corpus uteri & Uterus, NOS [4]			Gallbladder		
	Cases	*Rate*	*S.E.*[4]	*Cases*	*Rate*	*S.E.*	*Cases*	*Rate*	*S.E.*	*Cases*	*Rate*	*S.E.*	*Cases*	*Rate*	*S.E.*	*Cases*	*Rate*	*S.E.*
NAACCR U.S. Combined	**131067**	**328.4**	**0.95**	**38844**	**94.5**	**0.50**	**7186**	**14.6**	**0.18**	**13482**	**38.6**	**0.34**	**7498**	**18.7**	**0.22**	**1080**	**3.1**	**0.10**
Alabama	272	319.0	20.45	64	75.8	9.80	19	15.7	3.78	21	25.7	5.84	14	17.0	4.71	-	-	-
Alaska	101	452.1	61.78	39	153.9	32.75	7	17.5	7.68	-	-	-	7	22.0	9.54	-	-	-
Arizona	5004	293.9	4.43	1523	87.1	2.34	254	11.8	0.79	464	31.2	1.52	240	13.3	0.90	50	3.5	0.51
California	45824	310.2	1.54	13676	89.2	0.80	3027	16.1	0.31	4023	31.9	0.53	2517	16.9	0.35	456	3.6	0.18
CA: Greater Bay Area	7194	362.9	4.46	2243	110.2	2.40	327	13.0	0.75	664	37.9	1.52	397	20.1	1.04	60	3.4	0.45
CA: Los Angeles	18104	299.0	2.36	5191	83.1	1.21	1433	19.0	0.53	1566	30.3	0.80	1031	16.6	0.54	211	4.1	0.29
Colorado	3617	355.3	6.23	1073	101.1	3.21	167	12.1	0.99	380	43.1	2.29	146	14.4	1.24	26	3.1	0.62
Connecticut	1703	386.6	10.35	516	109.4	5.24	86	14.6	1.75	160	44.6	3.79	98	20.6	2.24	19	5.2	1.28
Delaware	137	328.7	32.01	47	111.5	18.30	11	16.8	5.93	6	18.0	7.67	14	38.7	11.00	-	-	-
District of Columbia	204	307.8	22.74	57	86.5	11.90	14	18.2	5.00	22	38.1	8.50	14	22.2	6.15	-	-	-
Florida	22841	353.7	2.35	6631	103.2	1.27	881	13.4	0.45	3195	49.9	0.88	1396	21.6	0.58	123	1.9	0.17
GA: Metropolitan Atlanta	619	346.9	16.62	193	111.7	9.04	42	16.9	3.21	44	33.3	5.64	31	17.6	3.52	-	-	-
Hawaii	192	184.7	14.19	62	54.9	7.34	-	-	-	20	20.2	4.71	12	11.1	3.51	-	-	-
Idaho	338	385.6	24.69	100	111.3	12.95	22	16.3	3.93	31	46.3	9.30	18	24.5	6.79	-	-	-
Illinois	5423	311.5	4.83	1568	85.3	2.38	389	17.1	0.97	431	31.0	1.65	334	19.4	1.19	51	3.4	0.52
Indiana	853	364.9	13.71	246	99.8	6.76	55	17.1	2.51	74	41.4	5.07	62	25.2	3.47	-	-	-
Iowa	277	357.6	23.93	81	98.5	11.70	13	11.9	3.65	19	33.7	8.28	18	24.0	5.98	-	-	-
Kentucky	299	515.4	31.84	86	144.8	16.20	9	10.1	3.53	34	69.8	12.41	18	32.1	7.97	-	-	-
Louisiana	1057	484.9	15.05	303	137.9	7.97	34	14.2	2.45	116	56.3	5.26	53	23.6	3.26	9	4.1	1.38
Maine	81	704.4	82.35	31	271.0	50.78	-	-	-	-	-	-	-	-	-	-	-	-
Massachusetts	1814	329.0	8.69	504	87.4	4.31	107	15.8	1.73	167	35.5	3.01	134	23.9	2.23	19	4.2	1.04
Michigan	967	243.2	8.43	347	87.7	4.97	51	9.2	1.38	83	25.5	2.96	54	14.1	2.01	-	-	-
MI: Metropolitan Detroit	650	415.9	17.20	188	115.5	8.82	25	12.3	2.74	62	45.0	5.95	30	19.1	3.55	-	-	-
Minnesota	365	311.0	19.21	108	85.1	9.44	24	10.0	2.20	26	30.9	6.62	24	20.8	4.90	-	-	-
Missouri	576	391.5	17.12	171	114.0	9.03	24	12.7	2.71	59	45.7	6.12	27	17.7	3.50	-	-	-

N.B. In areas with small Latino populations, methods to indirectly identify Latinos (like NHIA) can overestimate the ethnicity-specific counts of cancer cases. Also, even small errors in Latino population estimates can affect the magnitude of the cancer rates.

Total Cancer Cases and Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates[1] for Selected Cancer Sites[2]
Selected Areas in the United States[3], 1998-2002, Hispanic/Latina, All Races, Females
{Page 2 of 4}

	All Sites			Breast			Cervix Uteri			Colon & Rectum			Corpus uteri & Uterus, NOS [4]			Gallbladder		
	Cases	*Rate*	*S.E.*[4]	*Cases*	*Rate*	*S.E.*	*Cases*	*Rate*	*S.E.*	*Cases*	*Rate*	*S.E.*	*Cases*	*Rate*	*S.E.*	*Cases*	*Rate*	*S.E.*
Montana	88	397.7	45.76	25	104.3	22.75	-	-	-	7	39.3	15.91	-	-	-	-	-	-
Nebraska	360	436.6	26.01	85	105.9	12.63	26	22.2	5.03	38	61.2	10.73	24	28.2	6.47	-	-	-
Nevada	1288	304.0	9.63	374	81.4	4.61	79	13.9	1.80	115	36.8	3.73	73	16.4	2.03	10	2.4	0.77
New Jersey	6425	346.2	4.61	1950	99.8	2.38	357	15.0	0.84	725	45.1	1.76	366	20.1	1.10	49	3.2	0.48
New Mexico	4357	290.0	4.44	1364	88.8	2.42	153	9.2	0.75	462	32.6	1.53	222	14.7	0.99	41	2.9	0.45
New York	17527	333.0	2.61	5049	91.9	1.33	886	14.2	0.49	1955	41.2	0.96	1117	21.0	0.65	143	3.0	0.26
Ohio	1114	386.0	12.24	328	109.1	6.31	43	11.7	1.92	110	45.7	4.51	72	24.1	2.98	8	3.8	1.39
Oklahoma	605	356.9	16.43	190	108.7	8.67	34	13.8	2.80	59	45.6	6.49	27	15.9	3.22	6	4.6	2.00
Oregon	896	399.8	15.54	263	117.0	8.17	48	15.0	2.55	68	41.9	5.50	52	20.5	3.25	-	-	-
Pennsylvania	2109	426.7	10.06	637	125.9	5.34	108	15.4	1.61	222	51.8	3.67	124	24.3	2.31	12	3.0	0.91
Rhode Island	478	451.5	22.97	138	118.7	11.18	26	18.3	4.14	49	60.9	9.22	26	25.2	5.27	-	-	-
South Carolina	285	324.6	21.21	87	102.8	11.78	22	16.3	3.83	31	37.2	7.16	14	14.5	4.43	-	-	-
Utah	652	319.8	14.14	197	99.1	7.66	42	12.8	2.29	63	37.1	4.95	28	13.4	2.88	-	-	-
Washington	1529	380.3	11.35	473	110.5	5.71	69	12.1	1.70	119	39.2	3.98	71	17.0	2.29	15	4.9	1.40
WA: Seattle/Puget Sound	834	406.8	16.26	282	125.0	8.38	36	11.8	2.30	63	40.8	5.58	32	14.5	2.92	8	5.1	1.96
West Virginia	81	373.4	41.77	27	123.9	23.95	-	-	-	11	51.3	15.56	-	-	-	-	-	-
Wisconsin	709	382.2	16.17	231	120.9	8.74	42	16.7	2.95	67	43.4	5.76	40	20.4	3.59	-	-	-

1. Rates are per 100,000 population and were age-adjusted by five-year age groups to the 2000 U.S. standard population based on single years of age. Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for the U.S. combined.
2. Selected cancer sites by gender among the U.S. total population and the U.S. Hispanic/Latino population.
3. See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.
4. Abbreviations: S.E. - Standard Error; NOS - Not Otherwise Specified.

N.B. In areas with small Latino populations, methods to indirectly identify Latinos (like NHIA) can overestimate the ethnicity-specific counts of cancer cases. Also, even small errors in Latino population estimates can affect the magnitude of the cancer rates.

Total Cancer Cases and Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates[1] for Selected Cancer Sites[2]
Selected Areas in the United States[3], 1998-2002, Hispanic/Latina, All Races, Females
{Page 3 of 4}

	Liver			Lung & Bronchus			NH[4] Lymphoma			Ovary			Stomach			Thyroid		
	Cases	*Rate*	*S.E.*[4]	*Cases*	*Rate*	*S.E.*	*Cases*	*Rate*	*S.E.*	*Cases*	*Rate*	*S.E.*	*Cases*	*Rate*	*S.E.*	*Cases*	*Rate*	*S.E.*
NAACCR U.S. Combined	**1570**	**4.5**	**0.12**	**9427**	**27.0**	**0.29**	**5608**	**14.3**	**0.20**	**4692**	**11.4**	**0.17**	**3235**	**9.1**	**0.17**	**5724**	**10.8**	**0.15**
Alabama	-	-	-	32	45.5	8.18	16	20.4	5.36	9	10.5	3.59	-	-	-	14	11.0	3.30
Alaska	-	-	-	7	45.2	22.23	-	-	-	-	-	-	-	-	-	6	19.3	11.37
Arizona	60	4.0	0.54	334	23.6	1.34	198	11.6	0.87	198	12.3	0.93	109	7.5	0.75	249	10.6	0.73
California	644	5.2	0.21	2910	23.8	0.46	2016	14.4	0.34	1749	11.4	0.29	1257	9.8	0.29	2036	9.8	0.24
CA: Greater Bay Area	101	5.9	0.60	544	31.2	1.37	353	18.7	1.04	257	12.2	0.80	178	10.1	0.78	281	10.0	0.64
CA: Los Angeles	286	5.7	0.35	994	19.8	0.65	757	13.1	0.50	748	12.1	0.47	564	10.6	0.47	814	9.9	0.37
Colorado	36	4.0	0.69	308	35.7	2.09	141	14.3	1.27	136	12.7	1.14	110	13.1	1.29	167	12.3	1.03
Connecticut	11	3.1	1.04	129	37.8	3.52	65	14.2	1.93	47	11.0	1.79	41	11.0	1.86	93	13.7	1.51
Delaware	-	-	-	7	22.8	9.26	-	-	-	6	16.4	6.90	-	-	-	10	15.3	5.72
District of Columbia	-	-	-	11	17.5	5.58	10	16.4	5.34	9	11.2	3.93	7	9.3	3.71	14	14.4	4.10
Florida	218	3.4	0.23	1824	28.4	0.67	964	14.9	0.48	771	12.0	0.43	416	6.4	0.32	818	12.2	0.43
GA: Metropolitan Atlanta	-	-	-	30	23.7	4.78	22	13.6	3.52	27	16.8	3.71	17	16.1	4.19	35	13.3	2.82
Hawaii	-	-	-	18	20.6	5.03	6	6.3	2.78	-	-	-	6	6.4	2.81	9	7.1	2.64
Idaho	-	-	-	24	36.0	7.97	11	14.4	4.83	11	8.8	3.28	-	-	-	14	10.9	3.59
Illinois	72	6.0	0.76	349	26.6	1.55	225	13.7	1.03	196	10.3	0.84	148	9.1	0.84	264	9.8	0.68
Indiana	12	6.0	1.84	68	35.9	4.57	38	19.3	3.42	28	12.1	2.57	18	10.4	2.60	37	11.3	2.11
Iowa	-	-	-	18	29.2	7.16	-	-	-	17	19.7	5.41	-	-	-	15	12.3	3.60
Kentucky	-	-	-	49	98.8	14.62	6	12.1	5.05	11	20.4	6.55	6	10.2	4.76	15	13.3	3.88
Louisiana	14	6.7	1.80	119	56.4	5.21	45	21.1	3.17	49	21.4	3.08	26	12.7	2.51	37	15.3	2.54
Maine	-	-	-	11	97.2	30.54	-	-	-	-	-	-	-	-	-	-	-	-
Massachusetts	20	4.6	1.11	121	26.2	2.58	92	16.1	1.92	51	8.3	1.29	41	9.2	1.58	76	9.3	1.19
Michigan	8	2.5	0.94	60	17.0	2.30	35	8.8	1.63	22	5.7	1.30	18	5.0	1.25	49	9.0	1.44
MI: Metropolitan Detroit	6	4.2	1.74	67	47.9	6.06	30	20.5	3.93	25	16.0	3.31	16	11.7	3.10	21	10.8	2.57
Minnesota	-	-	-	32	42.1	8.00	24	26.7	6.12	8	6.4	2.59	8	7.3	2.85	15	8.2	2.81
Missouri	10	6.7	2.27	46	34.4	5.23	25	18.2	3.76	17	11.9	3.02	19	13.4	3.23	24	11.1	2.38

N.B. In areas with small Latino populations, methods to indirectly identify Latinos (like NHIA) can overestimate the ethnicity-specific counts of cancer cases. Also, even small errors in Latino population estimates can affect the magnitude of the cancer rates.

Total Cancer Cases and Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates[1] for Selected Cancer Sites[2]
Selected Areas in the United States[3], 1998-2002, Hispanic/Latina, All Races, Females
{Page 4 of 4}

	Liver			Lung & Bronchus			NH[4] Lymphoma			Ovary			Stomach			Thyroid		
	Cases	*Rate*	*S.E.*[4]	*Cases*	*Rate*	*S.E.*	*Cases*	*Rate*	*S.E.*	*Cases*	*Rate*	*S.E.*	*Cases*	*Rate*	*S.E.*	*Cases*	*Rate*	*S.E.*
Montana	-	-	-	8	45.7	16.66	-	-	-	6	22.8	9.81	-	-	-	8	28.1	10.97
Nebraska	6	10.2	4.38	33	46.6	8.48	15	23.5	6.76	9	10.2	3.64	-	-	-	24	14.3	3.16
Nevada	16	4.5	1.28	116	35.8	3.59	56	14.8	2.28	52	10.9	1.72	22	6.6	1.58	76	10.8	1.41
New Jersey	65	4.1	0.54	384	24.2	1.29	296	15.8	0.98	210	10.5	0.78	137	8.4	0.76	313	12.3	0.73
New Mexico	55	4.0	0.54	303	21.6	1.25	145	10.0	0.83	157	10.4	0.84	121	8.5	0.78	209	12.5	0.87
New York	226	4.7	0.32	1319	27.3	0.77	808	15.1	0.55	597	11.1	0.47	505	10.7	0.49	671	10.2	0.41
Ohio	11	4.4	1.34	109	43.7	4.32	48	16.3	2.46	41	14.9	2.44	21	7.0	1.61	45	12.4	2.02
Oklahoma	-	-	-	61	46.1	6.40	20	12.2	3.01	27	14.5	3.13	10	6.8	2.28	17	6.7	1.78
Oregon	10	6.8	2.26	88	54.9	6.27	37	14.5	2.88	28	12.5	2.66	23	12.5	2.95	52	12.5	2.18
Pennsylvania	19	4.1	1.00	180	43.2	3.36	95	18.8	2.12	62	11.6	1.59	45	10.9	1.72	104	15.5	1.70
Rhode Island	-	-	-	39	43.0	7.33	19	19.1	4.89	12	12.0	3.87	17	18.0	4.83	25	16.1	3.40
South Carolina	-	-	-	22	28.0	6.20	11	13.1	4.17	12	16.0	5.06	-	-	-	7	4.6	1.99
Utah	-	-	-	42	27.9	4.62	27	13.5	3.02	25	13.3	2.84	15	9.5	2.65	40	13.5	2.33
Washington	13	3.1	1.01	152	51.1	4.50	52	11.6	1.86	62	15.8	2.28	24	7.2	1.63	105	16.0	1.91
WA: Seattle/Puget Sound	6	2.6	1.26	95	61.7	6.81	26	12.9	2.84	31	15.1	3.03	11	6.8	2.30	50	14.9	2.58
West Virginia	-	-	-	7	32.3	12.25	-	-	-	-	-	-	-	-	-	-	-	-
Wisconsin	-	-	-	57	36.3	5.14	26	14.8	3.37	21	9.3	2.32	20	12.9	3.20	28	7.6	1.65

1. Rates are per 100,000 population and were age-adjusted by five-year age groups to the 2000 U.S. standard population based on single years of age. Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for the U.S. combined.
2. Selected cancer sites by gender among the U.S. total population and the U.S. Hispanic/Latino population.
3. See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.
4. Abbreviations: S.E. - Standard Error; NH - Non-Hodgkin.

N.B. In areas with small Latino populations, methods to indirectly identify Latinos (like NHIA) can overestimate the ethnicity-specific counts of cancer cases. Also, even small errors in Latino population estimates can affect the magnitude of the cancer rates.

Total Cancer Cases and Average Annual Age-adjusted Cancer Incidence Rates*
NAACCR (U.S.) Combined, 1998-2002, Non-Hispanic White

	MALES (NON-HISPANIC WHITE)					FEMALES (NON-HISPANIC WHITE)				
SITES	*Inc.*	*Adj. Rate 2000 US*	*S.E.*	*Adj. Rate World*	*S.E.*	*Inc.*	*Adj. Rate 2000 US*	*S.E.*	*Adj. Rate World*	*S.E.*
All Sites	2209923	562.5	0.38	369.9	0.27	2103352	431.1	0.30	300.1	0.23
Buccal Cavity & Pharynx	63618	15.9	0.06	11.4	0.05	31091	6.4	0.04	4.4	0.03
Lip	6911	1.8	0.02	1.1	0.01	2026	0.4	0.01	0.2	0.01
Tongue	16082	4.0	0.03	3.0	0.02	7846	1.6	0.02	1.1	0.01
Major salivary gland	6357	1.7	0.02	1.0	0.01	4379	0.9	0.01	0.7	0.01
Floor of mouth	4713	1.2	0.02	0.9	0.01	2345	0.5	0.01	0.3	0.01
Gum and other mouth	7728	1.9	0.02	1.3	0.02	6812	1.3	0.02	0.9	0.01
Nasopharynx	2431	0.6	0.01	0.5	0.01	1193	0.3	0.01	0.2	0.01
Oropharynx	2423	0.6	0.01	0.4	0.01	1009	0.2	0.01	0.1	0.00
Hypopharynx	5333	1.3	0.02	0.9	0.01	1674	0.3	0.01	0.2	0.01
Digestive System	426594	109.6	0.17	67.7	0.11	382081	72.7	0.12	44.0	0.08
Esophagus	33969	8.6	0.05	5.7	0.03	10623	2.0	0.02	1.3	0.01
Stomach	36490	9.4	0.05	5.7	0.03	21810	4.1	0.03	2.4	0.02
Small intestine	7724	2.0	0.02	1.3	0.02	6890	1.4	0.02	0.9	0.01
Colon excl. rectum	179491	46.6	0.11	27.5	0.07	192148	36.0	0.08	21.0	0.06
Rectum & rectosigmoid	77605	19.7	0.07	12.9	0.05	59714	11.8	0.05	7.6	0.04
Anus	4617	1.2	0.02	0.8	0.01	7569	1.6	0.02	1.1	0.01
Liver	21439	5.4	0.04	3.6	0.03	9368	1.8	0.02	1.2	0.01
Gallbladder	2900	0.8	0.01	0.4	0.01	6869	1.3	0.02	0.8	0.01
Pancreas	49060	12.6	0.06	7.8	0.04	50389	9.5	0.04	5.6	0.03
Respiratory System	397228	100.5	0.16	64.6	0.11	303188	60.3	0.11	40.6	0.08
Larynx	30267	7.5	0.04	5.3	0.03	8084	1.7	0.02	1.3	0.02
Lung & bronchus	362147	91.7	0.15	58.4	0.10	291851	58.0	0.11	38.9	0.08
Bones & Joints	4196	1.1	0.02	1.0	0.02	3482	0.8	0.01	0.8	0.01
Soft Tissue (incl. heart)	13432	3.5	0.03	2.6	0.02	11227	2.5	0.02	2.0	0.02
Skin (excl. basal and sq.)	98954	25.2	0.08	17.5	0.06	74096	16.6	0.06	12.8	0.05
Melanomas of the skin	91510	23.2	0.08	16.3	0.06	68574	15.4	0.06	12.0	0.05
Breast (excl. *in situ*)	5474	1.4	0.02	0.9	0.01	647973	136.6	0.17	98.6	0.13
Female Genital System	-	-	-	-	-	246523	52.3	0.11	38.3	0.08
Cervix uteri	-	-	-	-	-	34603	8.1	0.04	6.5	0.04
Corpus uteri	-	-	-	-	-	119414	25.0	0.07	18.3	0.06
Uterus, NOS	-	-	-	-	-	3385	0.7	0.01	0.4	0.01
Ovary	-	-	-	-	-	70439	14.7	0.06	10.6	0.04
Male Genital System	652016	164.0	0.20	111.1	0.15	-	-	-	-	-
Prostate	623191	156.4	0.20	104.4	0.14	-	-	-	-	-
Testis	24757	6.5	0.04	6.0	0.04	-	-	-	-	-
Penis	3162	0.8	0.01	0.5	0.01	-	-	-	-	-
Urinary System	237154	60.9	0.13	37.6	0.08	102506	20.2	0.06	13.2	0.05
Bladder (incl. *in situ*)	163058	42.2	0.11	24.9	0.07	56152	10.8	0.05	6.6	0.03
Kidney & renal pelvis	68719	17.3	0.07	12.0	0.05	43336	8.9	0.04	6.2	0.03
Ureter	3754	1.0	0.02	0.6	0.01	2364	0.4	0.01	0.3	0.01
Eye	4126	1.1	0.02	0.8	0.01	3529	0.8	0.01	0.6	0.01
Brain & Other Nervous System	33470	8.6	0.05	7.0	0.04	27499	6.1	0.04	5.1	0.04
Brain	31760	8.1	0.05	6.6	0.04	25721	5.7	0.04	4.7	0.03
Endocrine System	19244	4.9	0.04	4.0	0.03	49408	11.9	0.05	10.1	0.05
Thyroid gland	16387	4.1	0.03	3.3	0.03	46949	11.4	0.05	9.5	0.05
Lymphomas	103719	26.6	0.08	18.6	0.06	93847	19.4	0.06	13.7	0.05
Hodgkin's disease	13139	3.4	0.03	3.1	0.03	11109	2.8	0.03	2.6	0.03
Non-Hodgkin's lymphoma	90580	23.2	0.08	15.5	0.06	82738	16.6	0.06	11.1	0.04
Multiple Myeloma	23999	6.2	0.04	3.8	0.03	20133	3.9	0.03	2.4	0.02
Leukemias	61718	16.1	0.07	11.1	0.05	47005	9.6	0.05	7.0	0.04
Acute lymphocytic	5933	1.6	0.02	2.0	0.03	4611	1.2	0.02	1.6	0.03
Chronic lymphocytic	20220	5.2	0.04	3.1	0.02	14343	2.7	0.02	1.6	0.02
Acute myeloid	18449	4.8	0.04	3.2	0.03	15358	3.2	0.03	2.2	0.02
Chronic myeloid	7433	1.9	0.02	1.2	0.02	5688	1.1	0.02	0.8	0.01
Other	9683	2.6	0.03	1.6	0.02	7005	1.3	0.02	0.8	0.01
Mesothelioma	8878	2.3	0.02	1.3	0.01	2313	0.5	0.01	0.3	0.01
Kaposi Sarcoma	2651	0.7	0.01	0.5	0.01	421	0.1	0.00	0.0	0.00
Ill-defined & Unknown	53452	14.0	0.06	8.2	0.04	57030	10.6	0.05	6.2	0.03
Breast *in situ* (not in All Sites)	700	0.2	0.01	0.1	0.00	136851	29.9	0.08	22.6	0.06

* Rates are per 100,000 population and were age-adjusted by five-year age groups to the 2000 U.S. standard population based on single years of age and to the World population standard.

- Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for All Sites combined.

N.B. In areas with small Latino populations, methods to indirectly identify Latinos (like NHIA) can overestimate the ethnicity-specific counts of cancer cases. Also, even small errors in Latino population estimates can affect the magnitude of the cancer rates.

Total Cancer Cases and Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates[1] for Selected Cancer Sites[2]
Selected Areas in the United States[3], 1998-2002, Non-Hispanic White, Males
{Page 1 of 4}

	All Sites			Bladder			Brain & ONS[4]			Colon & Rectum			Gallbladder			Kidney & Renal Pelvis		
	Cases	*Rate*	*S.E.*[4]	*Cases*	*Rate*	*S.E.*	*Cases*	*Rate*	*S.E.*	*Cases*	*Rate*	*S.E.*	*Cases*	*Rate*	*S.E.*	*Cases*	*Rate*	*S.E.*
NAACCR U.S. Combined	**2209923**	**562.5**	**0.38**	**163058**	**42.2**	**0.11**	**33470**	**8.6**	**0.05**	**257096**	**66.3**	**0.13**	**2900**	**0.8**	**0.01**	**68719**	**17.3**	**0.07**
Alabama	39849	507.6	2.60	2475	32.8	0.68	669	8.5	0.33	4619	60.0	0.90	58	0.8	0.10	1221	15.2	0.44
Alaska	4092	560.6	10.17	275	44.3	3.08	87	8.4	0.98	405	60.0	3.45	-	-	-	147	18.0	1.76
Arizona	47209	482.5	2.25	3833	39.1	0.64	716	7.8	0.30	5468	56.2	0.77	69	0.7	0.09	1521	15.7	0.41
California	241626	552.5	1.13	17686	40.9	0.31	3833	8.9	0.15	25945	59.8	0.37	231	0.5	0.04	6788	15.4	0.19
CA: Greater Bay Area	47519	569.4	2.64	3158	38.9	0.70	807	9.6	0.34	5033	61.1	0.87	30	0.4	0.07	1321	15.6	0.43
CA: Los Angeles	50071	570.4	2.56	3764	43.1	0.70	757	9.1	0.33	5581	63.7	0.86	52	0.6	0.08	1448	16.4	0.43
Colorado	35568	516.8	2.81	2470	37.5	0.78	672	8.8	0.35	3570	54.1	0.93	42	0.7	0.11	1064	15.1	0.47
Connecticut	41746	595.1	2.92	3320	47.6	0.83	634	9.2	0.37	4935	70.7	1.01	67	1.0	0.12	1293	18.2	0.51
Delaware	8525	569.9	6.27	593	40.3	1.69	108	7.3	0.71	961	65.9	2.17	8	0.5	0.18	299	19.9	1.17
District of Columbia	1810	500.0	12.05	108	31.9	3.11	32	8.8	1.75	157	44.9	3.64	-	-	-	51	14.9	2.15
Florida	207262	571.0	1.28	16509	43.9	0.35	2651	8.4	0.17	23659	64.3	0.42	279	0.7	0.04	6132	17.3	0.23
GA: Metropolitan Atlanta	16615	553.3	4.50	961	35.6	1.20	300	8.7	0.52	1582	54.7	1.45	12	0.5	0.16	518	16.5	0.76
Hawaii	4231	579.8	9.08	256	36.5	2.33	61	8.9	1.19	430	59.6	2.92	-	-	-	125	16.2	1.47
Idaho	13910	533.6	4.56	993	39.3	1.26	290	10.5	0.62	1350	52.8	1.45	10	0.4	0.14	394	14.7	0.75
Illinois	118837	573.5	1.67	8805	43.3	0.46	1661	7.9	0.19	14787	72.4	0.60	187	0.9	0.07	3898	18.5	0.30
Indiana	63593	536.4	2.15	4277	37.3	0.58	981	7.9	0.25	7938	68.8	0.78	93	0.8	0.09	2246	18.4	0.39
Iowa	38097	553.6	2.84	2649	38.6	0.75	567	8.3	0.35	5072	74.1	1.04	52	0.8	0.11	1223	17.7	0.51
Kentucky	49848	608.8	2.79	3050	38.6	0.72	667	7.7	0.30	5792	72.8	0.98	63	0.9	0.11	1589	18.8	0.48
Louisiana	38811	592.1	3.06	2449	38.8	0.80	542	8.1	0.35	4665	72.4	1.08	50	0.8	0.11	1397	20.6	0.56
Maine	18749	607.6	4.49	1398	46.2	1.25	299	9.6	0.56	2120	70.3	1.55	22	0.7	0.16	602	18.9	0.78
Massachusetts	76589	591.3	2.15	6017	47.2	0.61	1139	8.8	0.26	8995	70.3	0.75	98	0.8	0.08	2309	17.6	0.37
Michigan	110152	595.1	1.81	8155	45.5	0.51	1733	9.1	0.22	11423	63.2	0.60	129	0.7	0.06	3180	16.8	0.30
MI: Metropolitan Detroit	42104	652.1	3.22	3202	50.8	0.91	608	9.1	0.37	4217	66.5	1.04	63	1.0	0.13	1282	19.3	0.55
Minnesota	55182	549.4	2.35	3774	38.6	0.63	912	8.6	0.29	5971	60.3	0.78	80	0.8	0.09	1753	17.0	0.41
Missouri	61162	531.0	2.16	4150	37.1	0.58	1003	8.7	0.27	7790	69.0	0.79	82	0.7	0.08	2082	17.8	0.39

N.B. In areas with small Latino populations, methods to indirectly identify Latinos (like NHIA) can overestimate the ethnicity-specific counts of cancer cases. Also, even small errors in Latino population estimates can affect the magnitude of the cancer rates.

Total Cancer Cases and Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates[1] for Selected Cancer Sites[2]
Selected Areas in the United States[3], 1998-2002, Non-Hispanic White, Males
{Page 2 of 4}

	All Sites			Bladder			Brain & ONS[4]			Colon & Rectum			Gallbladder			Kidney & Renal Pelvis		
	Cases	*Rate*	*S.E.*[4]	*Cases*	*Rate*	*S.E.*	*Cases*	*Rate*	*S.E.*	*Cases*	*Rate*	*S.E.*	*Cases*	*Rate*	*S.E.*	*Cases*	*Rate*	*S.E.*
Montana	11185	533.2	5.08	768	37.3	1.36	172	8.2	0.63	1236	59.5	1.71	16	0.8	0.19	314	14.8	0.84
Nebraska	19576	533.3	3.82	1385	38.2	1.03	334	9.0	0.50	2515	69.1	1.38	20	0.5	0.12	648	17.6	0.69
Nevada	19845	541.2	4.00	1553	44.3	1.18	294	8.0	0.48	2206	61.7	1.37	15	0.4	0.11	603	15.9	0.67
New Jersey	94533	634.4	2.07	7442	50.2	0.59	1290	8.9	0.25	11324	76.6	0.72	140	0.9	0.08	2991	19.9	0.36
New Mexico	12728	529.3	4.74	846	35.5	1.24	174	7.7	0.60	1240	51.7	1.48	19	0.8	0.19	369	15.2	0.80
New York	178755	573.8	1.36	14499	46.9	0.39	2822	9.3	0.18	22802	73.7	0.49	285	0.9	0.06	6108	19.4	0.25
Ohio	120190	530.0	1.55	8988	40.7	0.44	1857	8.0	0.19	14614	65.7	0.55	183	0.8	0.06	3719	16.1	0.27
Oklahoma	36958	548.2	2.88	2267	34.4	0.73	583	8.8	0.37	4380	66.2	1.01	38	0.6	0.10	1128	16.5	0.50
Oregon	40257	547.2	2.74	3036	41.9	0.76	678	9.2	0.36	4153	57.0	0.89	37	0.5	0.09	1112	14.8	0.45
Pennsylvania	160863	580.6	1.46	12598	45.7	0.41	2174	8.2	0.18	20333	73.9	0.52	276	1.0	0.06	5009	18.0	0.26
Rhode Island	14207	632.8	5.33	1216	54.4	1.57	191	8.7	0.63	1711	76.7	1.86	22	1.0	0.21	479	21.1	0.97
South Carolina	36705	559.7	3.01	2427	38.9	0.82	522	7.8	0.35	4130	64.4	1.04	32	0.5	0.10	1220	18.2	0.53
Utah	16712	491.5	3.86	1040	32.7	1.03	348	8.8	0.48	1599	48.6	1.23	25	0.8	0.17	428	12.2	0.60
Washington	64058	577.4	2.30	4640	43.2	0.64	1108	9.6	0.29	6517	59.8	0.75	62	0.6	0.08	1934	16.9	0.39
WA: Seattle/Puget Sound	44001	596.6	2.88	3268	46.0	0.81	746	9.6	0.36	4292	59.3	0.92	34	0.5	0.09	1328	17.4	0.48
West Virginia	25916	584.2	3.69	1810	41.8	1.00	344	7.8	0.43	3145	72.0	1.31	20	0.5	0.10	797	17.6	0.63
Wisconsin	63972	559.2	2.22	4340	38.6	0.59	1022	8.8	0.28	7557	66.7	0.77	70	0.6	0.07	2028	17.4	0.39

1. Rates are per 100,000 population and were age-adjusted by five-year age groups to the 2000 U.S. standard population based on single years of age. Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for the U.S. combined.
2. Selected cancer sites by gender among the U.S. total population and the U.S. Hispanic/Latino population.
3. See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.
4. Abbreviations: S.E. - Standard Error; ONS - Other Nervous System

N.B. In areas with small Latino populations, methods to indirectly identify Latinos (like NHIA) can overestimate the ethnicity-specific counts of cancer cases. Also, even small errors in Latino population estimates can affect the magnitude of the cancer rates.

Total Cancer Cases and Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates[1] for Selected Cancer Sites[2]
Selected Areas in the United States[3], 1998-2002, Non-Hispanic White, Males
{Page 3 of 4}

	Liver			Lung & Bronchus			NH[4] Lymphoma			Prostate			Stomach			Testis		
	Cases	*Rate*	*S.E.*[4]	*Cases*	*Rate*	*S.E.*	*Cases*	*Rate*	*S.E.*	*Cases*	*Rate*	*S.E.*	*Cases*	*Rate*	*S.E.*	*Cases*	*Rate*	*S.E.*
NAACCR U.S. Combined	**21439**	**5.4**	**0.04**	**362147**	**91.7**	**0.15**	**90580**	**23.2**	**0.08**	**623191**	**156.4**	**0.20**	**36490**	**9.4**	**0.05**	**24757**	**6.5**	**0.04**
Alabama	435	5.6	0.28	8833	111.7	1.21	1570	20.1	0.52	9380	116.6	1.23	559	7.4	0.32	395	5.1	0.26
Alaska	55	6.5	1.03	563	83.2	3.98	207	26.5	2.23	1260	182.6	5.83	62	9.7	1.50	85	6.9	0.76
Arizona	467	4.9	0.23	7706	76.8	0.88	1864	19.7	0.46	12322	121.3	1.10	691	7.0	0.27	509	6.3	0.28
California	2534	5.7	0.11	34350	78.4	0.42	10446	23.9	0.24	69146	156.4	0.60	4123	9.5	0.15	3122	7.5	0.14
CA: Greater Bay Area	551	6.3	0.27	5757	70.2	0.93	2421	28.4	0.58	14852	177.8	1.47	823	10.0	0.35	681	7.8	0.30
CA: Los Angeles	551	6.2	0.27	6554	74.1	0.92	2267	26.0	0.55	14176	159.2	1.34	945	10.8	0.35	607	7.5	0.31
Colorado	277	3.9	0.24	4441	67.3	1.03	1520	21.5	0.57	10943	159.4	1.55	468	7.0	0.33	586	6.7	0.28
Connecticut	396	5.6	0.28	6054	86.0	1.11	1749	25.0	0.60	12155	171.5	1.56	868	12.5	0.42	457	7.1	0.34
Delaware	78	5.2	0.60	1462	96.7	2.57	341	22.8	1.25	2421	157.6	3.25	126	8.8	0.80	84	6.1	0.66
District of Columbia	26	7.7	1.63	214	62.4	4.32	74	20.5	2.50	576	160.1	6.77	28	8.2	1.56	40	7.2	1.19
Florida	2114	6.1	0.13	37035	99.7	0.53	7808	22.6	0.26	56381	148.3	0.63	3271	8.9	0.16	1495	6.2	0.16
GA: Metropolitan Atlanta	128	4.3	0.40	2494	89.2	1.86	732	23.2	0.91	4952	165.1	2.43	166	5.8	0.47	280	6.5	0.39
Hawaii	57	7.5	1.01	544	74.8	3.25	145	19.7	1.68	1067	148.8	4.63	74	10.8	1.27	48	6.0	0.87
Idaho	90	3.4	0.36	1867	72.5	1.69	550	20.7	0.89	4504	172.8	2.59	176	6.8	0.52	183	6.6	0.49
Illinois	1196	5.7	0.17	20113	97.0	0.69	5021	24.2	0.34	32004	153.2	0.86	2017	9.9	0.22	1287	6.1	0.17
Indiana	554	4.6	0.20	12741	107.9	0.97	2626	21.9	0.43	15416	129.8	1.06	901	7.7	0.26	765	5.9	0.21
Iowa	280	4.1	0.24	6232	90.0	1.14	1545	22.6	0.58	10593	152.8	1.49	541	7.9	0.34	449	6.9	0.33
Kentucky	401	4.9	0.25	11370	138.1	1.32	1851	22.4	0.53	12016	147.1	1.37	718	8.9	0.34	493	5.5	0.25
Louisiana	509	7.8	0.35	7206	109.4	1.31	1535	23.7	0.62	10545	159.9	1.59	564	8.9	0.38	397	5.7	0.29
Maine	162	5.3	0.43	3130	101.0	1.82	706	22.9	0.87	5312	170.3	2.36	327	10.7	0.60	206	6.9	0.48
Massachusetts	807	6.2	0.22	11280	87.2	0.82	2896	22.4	0.42	22957	175.4	1.16	1425	11.3	0.30	950	7.2	0.23
Michigan	1064	5.7	0.18	17164	93.0	0.72	4383	23.6	0.36	34357	184.4	1.00	1672	9.4	0.23	1200	6.1	0.18
MI: Metropolitan Detroit	444	6.8	0.33	6455	100.5	1.26	1644	25.3	0.63	13355	206.4	1.80	750	12.0	0.45	470	6.7	0.31
Minnesota	367	3.6	0.19	7155	71.6	0.85	2541	25.1	0.50	18363	183.9	1.36	842	8.6	0.30	835	7.6	0.26
Missouri	540	4.7	0.20	11987	103.5	0.95	2638	23.2	0.46	15369	130.6	1.06	868	7.6	0.26	612	5.3	0.22

N.B. In areas with small Latino populations, methods to indirectly identify Latinos (like NHIA) can overestimate the ethnicity-specific counts of cancer cases. Also, even small errors in Latino population estimates can affect the magnitude of the cancer rates.

Total Cancer Cases and Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates[1] for Selected Cancer Sites[2]
Selected Areas in the United States[3], 1998-2002, Non-Hispanic White, Males
{Page 4 of 4}

	Liver			Lung & Bronchus			NH[4] Lymphoma			Prostate			Stomach			Testis		
	Cases	*Rate*	*S.E.*[4]	*Cases*	*Rate*	*S.E.*	*Cases*	*Rate*	*S.E.*	*Cases*	*Rate*	*S.E.*	*Cases*	*Rate*	*S.E.*	*Cases*	*Rate*	*S.E.*
Montana	73	3.5	0.41	1703	81.5	1.99	457	22.0	1.04	3624	169.6	2.83	189	9.1	0.67	121	6.2	0.57
Nebraska	125	3.4	0.30	2998	81.3	1.49	836	22.8	0.79	5786	156.7	2.06	298	8.2	0.47	251	6.8	0.43
Nevada	222	5.8	0.39	3632	98.1	1.69	736	20.6	0.79	5305	141.1	2.02	270	7.7	0.49	201	5.9	0.42
New Jersey	949	6.4	0.21	13327	88.9	0.77	3883	26.4	0.43	29184	192.7	1.13	1861	12.7	0.30	1003	7.5	0.24
New Mexico	126	5.3	0.48	1707	70.4	1.72	472	20.0	0.93	4013	161.8	2.57	167	7.2	0.56	134	7.0	0.61
New York	2123	6.8	0.15	27502	87.7	0.53	7275	23.6	0.28	48971	154.7	0.70	3868	12.6	0.20	1931	6.6	0.15
Ohio	1040	4.6	0.15	22522	99.1	0.67	5258	23.0	0.32	31184	136.5	0.78	1847	8.3	0.20	1388	5.9	0.16
Oklahoma	356	5.3	0.28	7862	115.5	1.31	1478	22.2	0.58	9788	143.1	1.46	486	7.3	0.33	330	5.2	0.29
Oregon	310	4.1	0.23	6222	84.9	1.08	1670	22.7	0.56	12044	162.5	1.49	610	8.4	0.34	544	7.6	0.33
Pennsylvania	1528	5.5	0.14	25474	90.8	0.57	6646	24.3	0.30	46316	164.1	0.77	2923	10.7	0.20	1645	6.7	0.17
Rhode Island	139	6.1	0.52	2322	102.5	2.13	515	23.0	1.02	3978	175.6	2.79	287	12.8	0.76	139	6.5	0.55
South Carolina	373	5.6	0.30	7012	105.9	1.30	1401	21.6	0.60	10012	149.1	1.53	510	8.0	0.37	340	5.1	0.28
Utah	100	2.8	0.29	1407	42.7	1.15	819	23.8	0.85	6138	185.0	2.38	225	6.9	0.46	313	6.5	0.37
Washington	620	5.4	0.22	9533	87.2	0.90	2845	25.3	0.48	19369	174.8	1.27	986	9.0	0.29	876	7.2	0.24
WA: Seattle/Puget Sound	451	5.9	0.28	6361	88.1	1.12	2034	27.1	0.61	13414	183.0	1.59	701	9.7	0.37	642	7.5	0.30
West Virginia	199	4.5	0.33	5489	122.1	1.67	961	21.8	0.71	6793	151.0	1.86	366	8.6	0.46	198	4.8	0.34
Wisconsin	619	5.4	0.22	9494	83.1	0.86	2580	22.6	0.45	18647	161.9	1.19	1110	9.8	0.30	865	7.3	0.25

1. Rates are per 100,000 population and were age-adjusted by five-year age groups to the 2000 U.S. standard population based on single years of age. Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for the U.S. combined.
2. Selected cancer sites by gender among the U.S. total population and the U.S. Hispanic/Latino population.
3. See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.
4. Abbreviations: S.E. - Standard Error; NH - Non-Hodgkin.

N.B. In areas with small Latino populations, methods to indirectly identify Latinos (like NHIA) can overestimate the ethnicity-specific counts of cancer cases. Also, even small errors in Latino population estimates can affect the magnitude of the cancer rates.

Total Cancer Cases and Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates[1] for Selected Cancer Sites[2]
Selected Areas in the United States[3], 1998-2002, Non-Hispanic White, Females
{Page 1 of 4}

	All Sites			Breast			Cervix Uteri			Colon & Rectum			Corpus uteri & Uterus, NOS [4]			Gallbladder		
	Cases	*Rate*	*S.E.*[4]	*Cases*	*Rate*	*S.E.*	*Cases*	*Rate*	*S.E.*	*Cases*	*Rate*	*S.E.*	*Cases*	*Rate*	*S.E.*	*Cases*	*Rate*	*S.E.*
NAACCR U.S. Combined	**2103352**	**431.1**	**0.30**	**647973**	**136.6**	**0.17**	**34603**	**8.1**	**0.04**	**251862**	**47.8**	**0.10**	**122799**	**25.7**	**0.07**	**6869**	**1.3**	**0.02**
Alabama	36598	370.5	1.96	11526	119.3	1.12	709	8.3	0.31	4122	40.0	0.63	1603	16.3	0.41	86	0.8	0.09
Alaska	3471	430.2	7.83	1256	143.1	4.33	80	7.9	0.96	293	44.8	2.75	191	22.2	1.71	7	1.2	0.48
Arizona	43345	389.2	1.91	13949	128.5	1.11	620	6.8	0.28	4801	39.9	0.58	2143	19.3	0.42	119	1.0	0.09
California	234876	444.5	0.93	78394	152.0	0.55	3478	7.6	0.13	25005	43.6	0.28	12994	24.7	0.22	591	1.0	0.04
CA: Greater Bay Area	46358	452.4	2.13	16422	163.2	1.29	587	6.3	0.26	4846	44.0	0.64	2765	27.0	0.52	105	1.0	0.10
CA: Los Angeles	49417	469.3	2.16	16250	160.1	1.28	738	8.3	0.31	5590	47.7	0.65	2854	27.7	0.53	154	1.4	0.11
Colorado	34823	406.6	2.19	12098	140.2	1.28	555	6.4	0.27	3504	40.9	0.69	1740	20.4	0.49	83	1.0	0.11
Connecticut	41169	458.1	2.31	12779	146.8	1.32	510	6.4	0.29	5088	51.2	0.73	2589	29.5	0.59	123	1.2	0.11
Delaware	7911	434.5	4.93	2345	131.3	2.74	150	9.4	0.78	938	48.5	1.59	485	26.8	1.22	22	1.1	0.24
District of Columbia	1892	446.0	10.62	737	177.0	6.68	17	3.9	1.00	194	42.4	3.15	113	27.6	2.65	-	-	-
Florida	180020	435.5	1.08	51866	133.3	0.61	2976	10.0	0.19	21957	46.9	0.33	8647	21.3	0.24	489	1.0	0.05
GA: Metropolitan Atlanta	16674	423.6	3.30	5888	147.0	1.93	315	7.5	0.43	1634	42.2	1.05	853	21.5	0.74	31	0.8	0.15
Hawaii	3533	472.0	8.00	1216	160.1	4.61	71	9.6	1.15	316	41.6	2.35	200	26.5	1.88	-	-	-
Idaho	12236	400.7	3.64	3986	132.2	2.10	185	6.5	0.48	1262	39.7	1.12	694	22.9	0.87	38	1.2	0.20
Illinois	117051	435.3	1.29	35714	137.2	0.74	2119	8.9	0.20	14740	50.4	0.42	6933	26.6	0.32	474	1.6	0.08
Indiana	63414	416.3	1.67	19296	129.0	0.94	1195	8.7	0.25	7954	49.3	0.56	3897	26.0	0.42	203	1.3	0.09
Iowa	37035	425.0	2.27	11053	132.1	1.29	613	8.5	0.35	5350	54.6	0.77	2322	27.8	0.59	147	1.5	0.12
Kentucky	46213	439.5	2.05	13065	125.9	1.11	1089	11.2	0.34	5778	53.0	0.70	2342	22.5	0.47	125	1.1	0.10
Louisiana	33540	403.7	2.22	10173	124.5	1.24	672	9.1	0.35	4070	47.0	0.74	1446	17.8	0.47	92	1.1	0.11
Maine	17252	445.2	3.42	5034	132.3	1.88	266	7.6	0.47	2183	53.0	1.14	1126	29.8	0.89	61	1.5	0.19
Massachusetts	77802	456.7	1.67	23909	145.7	0.96	951	6.2	0.20	9506	50.8	0.53	4656	28.4	0.42	255	1.3	0.08
Michigan	101356	437.0	1.38	30712	134.7	0.77	1670	7.9	0.19	11392	46.4	0.44	6390	27.9	0.35	348	1.4	0.08
MI: Metropolitan Detroit	38831	462.7	2.37	11538	140.5	1.32	614	8.0	0.33	4349	48.5	0.74	2339	28.4	0.59	135	1.5	0.13
Minnesota	51187	410.8	1.84	17071	139.2	1.08	717	6.2	0.23	6037	45.2	0.59	3236	26.6	0.47	179	1.4	0.10
Missouri	59241	408.2	1.70	17944	127.5	0.96	1131	8.9	0.27	7466	47.7	0.56	3627	25.5	0.43	237	1.5	0.10

N.B. In areas with small Latino populations, methods to indirectly identify Latinos (like NHIA) can overestimate the ethnicity-specific counts of cancer cases. Also, even small errors in Latino population estimates can affect the magnitude of the cancer rates.

Total Cancer Cases and Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates[1] for Selected Cancer Sites[2]
Selected Areas in the United States[3], 1998-2002, Non-Hispanic White, Females
{Page 2 of 4}

	All Sites			Breast			Cervix Uteri			Colon & Rectum			Corpus uteri & Uterus, NOS [4]			Gallbladder		
	Cases	*Rate*	*S.E.*[4]	*Cases*	*Rate*	*S.E.*	*Cases*	*Rate*	*S.E.*	*Cases*	*Rate*	*S.E.*	*Cases*	*Rate*	*S.E.*	*Cases*	*Rate*	*S.E.*
Montana	9870	404.1	4.11	3086	128.2	2.33	180	8.3	0.63	1088	42.1	1.29	515	21.2	0.94	35	1.3	0.22
Nebraska	18937	409.9	3.05	6059	135.3	1.77	305	7.8	0.45	2493	49.3	1.02	1218	27.4	0.80	67	1.3	0.17
Nevada	16965	433.5	3.35	4982	126.6	1.80	353	9.6	0.51	1791	45.6	1.08	757	19.1	0.70	28	0.7	0.14
New Jersey	90655	473.0	1.61	26626	144.8	0.90	1438	8.7	0.23	11456	54.0	0.52	5873	31.6	0.42	312	1.4	0.08
New Mexico	11315	410.1	3.91	3837	139.2	2.27	203	8.5	0.61	1085	37.0	1.13	600	21.4	0.88	30	1.0	0.18
New York	182337	457.3	1.09	53452	140.0	0.62	2612	7.5	0.15	23628	53.9	0.36	11411	29.5	0.28	769	1.7	0.06
Ohio	120576	411.8	1.20	36660	128.1	0.68	2147	8.3	0.18	14855	47.7	0.40	7786	27.2	0.31	409	1.3	0.06
Oklahoma	34249	407.3	2.23	11042	134.8	1.30	718	10.1	0.38	4101	45.6	0.72	1693	20.2	0.50	93	1.0	0.11
Oregon	39007	438.5	2.24	12806	146.1	1.30	600	7.6	0.31	4123	43.5	0.69	2096	23.7	0.52	117	1.2	0.11
Pennsylvania	153492	432.8	1.14	44891	131.6	0.64	2396	8.2	0.17	20392	51.6	0.37	10571	31.2	0.31	622	1.5	0.06
Rhode Island	13540	454.0	4.02	3832	134.0	2.22	173	7.1	0.54	1768	52.8	1.30	771	27.0	0.99	46	1.5	0.23
South Carolina	31964	393.3	2.21	10296	127.9	1.27	665	9.2	0.36	3598	42.8	0.72	1521	18.7	0.48	91	1.1	0.11
Utah	14517	356.9	2.98	4907	123.2	1.76	255	6.2	0.39	1534	37.8	0.97	944	23.8	0.78	43	1.1	0.16
Washington	61039	453.1	1.85	20252	151.5	1.07	851	6.7	0.23	6076	43.2	0.56	3464	25.8	0.44	172	1.2	0.09
WA: Seattle/Puget Sound	42378	470.6	2.30	14270	159.3	1.34	572	6.5	0.27	4127	44.3	0.70	2404	26.8	0.55	117	1.3	0.12
West Virginia	24161	432.1	2.82	6474	118.7	1.49	569	11.7	0.50	3116	52.6	0.95	1526	27.9	0.72	62	1.0	0.13
Wisconsin	60089	423.2	1.75	18760	135.6	1.00	1049	8.5	0.26	7168	46.7	0.56	3826	27.8	0.45	257	1.7	0.11

1. Rates are per 100,000 population and were age-adjusted by five-year age groups to the 2000 U.S. standard population based on single years of age. Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for the U.S. combined.
2. Selected cancer sites by gender among the U.S. total population and the U.S. Hispanic/Latino population.
3. See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.
4. Abbreviations: S.E. - Standard Error; NOS - Not Otherwise Specified.

N.B. In areas with small Latino populations, methods to indirectly identify Latinos (like NHIA) can overestimate the ethnicity-specific counts of cancer cases. Also, even small errors in Latino population estimates can affect the magnitude of the cancer rates.

Total Cancer Cases and Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates[1] for Selected Cancer Sites[2]
Selected Areas in the United States[3], 1998-2002, Non-Hispanic White, Females
{Page 3 of 4}

	Liver			Lung & Bronchus			NH[4] Lymphoma			Ovary			Stomach			Thyroid		
	Cases	*Rate*	*S.E.*[4]	*Cases*	*Rate*	*S.E.*	*Cases*	*Rate*	*S.E.*	*Cases*	*Rate*	*S.E.*	*Cases*	*Rate*	*S.E.*	*Cases*	*Rate*	*S.E.*
NAACCR U.S. Combined	**9368**	**1.8**	**0.02**	**291851**	**58.0**	**0.11**	**82738**	**16.6**	**0.06**	**70439**	**14.7**	**0.06**	**21810**	**4.1**	**0.03**	**46949**	**11.4**	**0.05**
Alabama	213	2.1	0.14	5419	52.8	0.72	1451	14.3	0.38	1423	14.4	0.39	432	4.1	0.20	695	8.2	0.31
Alaska	21	2.9	0.67	449	63.5	3.14	153	19.3	1.66	108	12.7	1.32	22	3.3	0.73	120	10.9	1.02
Arizona	184	1.6	0.12	6401	53.8	0.68	1609	14.2	0.36	1399	12.8	0.35	354	3.0	0.16	1024	11.5	0.36
California	1103	2.0	0.06	31812	57.9	0.33	8729	16.3	0.18	7813	15.0	0.17	2372	4.1	0.09	4713	10.6	0.16
CA: Greater Bay Area	210	2.0	0.14	5662	54.1	0.73	1843	17.8	0.42	1548	15.3	0.39	478	4.3	0.20	829	9.3	0.33
CA: Los Angeles	265	2.4	0.15	6225	55.8	0.72	1854	17.2	0.41	1623	15.9	0.40	590	5.1	0.21	1167	13.4	0.40
Colorado	130	1.5	0.13	3785	45.2	0.74	1436	16.8	0.45	1279	14.9	0.42	277	3.2	0.19	1004	11.6	0.37
Connecticut	175	1.9	0.15	5399	58.5	0.81	1584	17.2	0.44	1277	14.5	0.41	546	5.3	0.23	915	12.4	0.42
Delaware	37	1.9	0.32	1142	60.2	1.79	305	16.4	0.95	248	13.7	0.88	69	3.5	0.42	210	13.4	0.93
District of Columbia	-	-	-	194	46.8	3.45	72	17.0	2.07	55	13.1	1.82	18	4.0	0.96	53	12.1	1.74
Florida	819	1.8	0.07	30351	67.8	0.40	6884	16.0	0.20	6104	15.2	0.21	1815	3.8	0.09	3022	10.2	0.19
GA: Metropolitan Atlanta	49	1.3	0.19	2079	55.0	1.21	604	15.5	0.64	585	14.9	0.62	134	3.5	0.30	490	11.9	0.54
Hawaii	12	1.6	0.46	417	55.5	2.73	123	16.7	1.52	106	13.9	1.36	43	5.7	0.87	79	11.0	1.25
Idaho	51	1.6	0.22	1373	44.6	1.21	529	17.3	0.75	422	13.8	0.67	121	3.9	0.35	343	12.1	0.65
Illinois	575	2.0	0.09	15845	57.6	0.46	4673	16.9	0.25	3790	14.5	0.24	1288	4.3	0.12	2664	11.6	0.23
Indiana	255	1.6	0.10	9074	58.7	0.62	2467	15.9	0.32	2048	13.6	0.30	523	3.2	0.14	1321	9.7	0.27
Iowa	138	1.4	0.13	4404	49.5	0.76	1540	16.9	0.44	1220	14.3	0.42	303	3.1	0.18	818	11.6	0.41
Kentucky	187	1.8	0.13	7704	72.0	0.82	1810	16.9	0.40	1409	13.5	0.36	401	3.7	0.18	977	10.2	0.33
Louisiana	200	2.3	0.17	5032	59.0	0.84	1449	17.1	0.45	1040	12.6	0.39	352	4.0	0.22	789	10.7	0.38
Maine	51	1.2	0.17	2433	61.7	1.26	668	17.1	0.67	532	13.8	0.60	178	4.3	0.32	330	9.9	0.55
Massachusetts	296	1.6	0.10	10772	61.9	0.61	2843	16.3	0.31	2456	14.8	0.30	946	4.9	0.16	1836	12.6	0.30
Michigan	519	2.2	0.10	13772	58.5	0.50	4133	17.5	0.27	3427	14.9	0.26	1016	4.1	0.13	2338	11.3	0.23
MI: Metropolitan Detroit	208	2.4	0.17	5589	65.3	0.88	1543	18.1	0.47	1279	15.4	0.44	474	5.2	0.24	927	12.5	0.41
Minnesota	158	1.2	0.10	5685	46.0	0.62	2287	18.0	0.38	1645	13.5	0.34	497	3.6	0.17	1142	10.0	0.30
Missouri	213	1.4	0.10	8609	58.3	0.63	2418	16.2	0.34	1934	13.6	0.31	475	3.0	0.14	1352	10.8	0.30

N.B. In areas with small Latino populations, methods to indirectly identify Latinos (like NHIA) can overestimate the ethnicity-specific counts of cancer cases. Also, even small errors in Latino population estimates can affect the magnitude of the cancer rates.

Total Cancer Cases and Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates[1] for Selected Cancer Sites[2]
Selected Areas in the United States[3], 1998-2002, Non-Hispanic White, Females
{Page 4 of 4}

	Liver			Lung & Bronchus			NH[4] Lymphoma			Ovary			Stomach			Thyroid		
	Cases	*Rate*	*S.E.*[4]	*Cases*	*Rate*	*S.E.*	*Cases*	*Rate*	*S.E.*	*Cases*	*Rate*	*S.E.*	*Cases*	*Rate*	*S.E.*	*Cases*	*Rate*	*S.E.*
Montana	37	1.4	0.24	1378	55.5	1.50	384	15.4	0.79	364	14.9	0.79	109	4.3	0.41	298	14.2	0.83
Nebraska	60	1.3	0.17	2149	45.7	1.00	806	17.0	0.61	624	13.9	0.57	171	3.3	0.26	434	11.1	0.54
Nevada	99	2.5	0.25	3117	77.1	1.39	556	14.3	0.61	591	15.0	0.62	152	3.9	0.32	490	13.8	0.63
New Jersey	376	1.8	0.09	12009	60.1	0.56	3724	18.9	0.32	3247	17.4	0.31	1171	5.4	0.16	2208	14.2	0.31
New Mexico	73	2.6	0.30	1348	46.2	1.26	449	15.9	0.76	385	14.1	0.73	79	2.7	0.31	372	16.1	0.85
New York	955	2.2	0.07	25012	60.5	0.39	7060	17.4	0.21	6455	16.7	0.21	2454	5.5	0.11	4376	13.4	0.21
Ohio	539	1.8	0.08	17218	57.4	0.44	4802	16.0	0.23	3830	13.3	0.22	1147	3.6	0.11	2279	8.9	0.19
Oklahoma	195	2.2	0.16	5449	62.6	0.86	1312	15.1	0.42	1147	13.9	0.42	282	3.1	0.19	506	7.2	0.33
Oregon	153	1.7	0.14	5580	61.6	0.83	1511	16.6	0.43	1241	14.0	0.40	285	3.0	0.18	716	9.2	0.35
Pennsylvania	623	1.6	0.07	19399	52.1	0.38	6245	17.0	0.22	5152	15.0	0.21	1805	4.5	0.11	4266	15.1	0.23
Rhode Island	62	1.9	0.25	1914	63.2	1.48	535	17.4	0.78	394	14.0	0.72	210	6.0	0.43	325	13.5	0.76
South Carolina	126	1.5	0.13	4482	53.2	0.80	1267	15.3	0.43	1055	13.0	0.40	301	3.5	0.21	654	9.1	0.36
Utah	60	1.5	0.19	854	21.5	0.74	616	15.2	0.61	560	13.8	0.59	156	3.8	0.31	633	14.5	0.59
Washington	261	1.9	0.12	8420	62.4	0.68	2380	17.4	0.36	2178	16.2	0.35	518	3.7	0.16	1383	11.0	0.30
WA: Seattle/Puget Sound	190	2.1	0.15	5767	64.5	0.85	1653	18.2	0.45	1518	16.9	0.44	355	3.8	0.20	1009	11.7	0.37
West Virginia	93	1.6	0.17	4061	69.6	1.10	960	16.9	0.55	805	14.6	0.52	218	3.6	0.25	523	11.0	0.49
Wisconsin	266	1.8	0.11	7310	50.9	0.60	2364	16.2	0.34	2091	15.1	0.33	570	3.7	0.16	1251	10.1	0.29

1. Rates are per 100,000 population and were age-adjusted by five-year age groups to the 2000 U.S. standard population based on single years of age. Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for the U.S. combined.
2. Selected cancer sites by gender among the U.S. total population and the U.S. Hispanic/Latino population.
3. See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.
4. Abbreviations: S.E. - Standard Error; NH - Non-Hodgkin.

N.B. In areas with small Latino populations, methods to indirectly identify Latinos (like NHIA) can overestimate the ethnicity-specific counts of cancer cases. Also, even small errors in Latino population estimates can affect the magnitude of the cancer rates.

Total Cancer Cases and Average Annual Age-adjusted Cancer Incidence Rates*
NAACCR (U.S.) Combined, 1998-2002, Non-Hispanic Black

	MALES (NON-HISPANIC BLACK)					FEMALES (NON-HISPANIC BLACK)				
SITES	*Inc.*	*Adj. Rate 2000 US*	*S.E.*	*Adj. Rate World*	*S.E.*	*Inc.*	*Adj. Rate 2000 US*	*S.E.*	*Adj. Rate World*	*S.E.*
All Sites	232873	652.8	1.42	439.4	0.93	203771	391.3	0.87	270.5	0.63
Buccal Cavity & Pharynx	7693	19.0	0.22	14.4	0.17	3154	5.9	0.11	4.4	0.08
Lip	50	0.1	0.02	0.1	0.01	36	0.1	0.01	0.0	0.01
Tongue	1664	4.1	0.10	3.2	0.08	639	1.2	0.05	0.9	0.04
Major salivary gland	455	1.1	0.06	0.8	0.04	421	0.8	0.04	0.6	0.03
Floor of mouth	755	1.9	0.07	1.4	0.05	234	0.4	0.03	0.3	0.02
Gum and other mouth	1018	2.6	0.08	1.9	0.06	658	1.3	0.05	0.9	0.04
Nasopharynx	494	1.1	0.05	0.9	0.04	259	0.5	0.03	0.4	0.02
Oropharynx	578	1.4	0.06	1.1	0.05	167	0.3	0.02	0.2	0.02
Hypopharynx	1067	2.7	0.09	2.1	0.06	231	0.4	0.03	0.3	0.02
Digestive System	47932	137.7	0.66	88.7	0.41	46429	92.5	0.43	57.8	0.28
Esophagus	4625	12.7	0.19	8.9	0.13	1870	3.7	0.09	2.5	0.06
Stomach	6104	18.3	0.25	11.1	0.15	4666	9.4	0.14	5.4	0.08
Small intestine	1179	3.3	0.10	2.1	0.06	1159	2.3	0.07	1.5	0.05
Colon excl. rectum	18332	54.7	0.42	33.9	0.26	21838	43.6	0.30	27.1	0.19
Rectum & rectosigmoid	6046	16.7	0.23	11.2	0.15	5607	11.0	0.15	7.3	0.10
Anus	645	1.5	0.06	1.1	0.04	688	1.3	0.05	0.9	0.04
Liver	4045	10.3	0.17	7.4	0.12	1594	3.1	0.08	2.1	0.05
Gallbladder	293	0.9	0.05	0.5	0.03	871	1.7	0.06	1.1	0.04
Pancreas	5574	16.1	0.23	10.4	0.14	6750	13.6	0.17	8.2	0.11
Respiratory System	44971	127.8	0.63	85.0	0.41	28032	55.3	0.33	37.9	0.24
Larynx	4542	12.1	0.19	8.7	0.13	1271	2.4	0.07	1.8	0.05
Lung & bronchus	39896	114.4	0.60	75.4	0.38	26445	52.3	0.32	35.6	0.23
Bones & Joints	479	0.9	0.04	0.8	0.04	422	0.7	0.03	0.6	0.03
Soft Tissue (incl. heart)	1497	3.3	0.09	2.6	0.07	1603	2.8	0.07	2.3	0.06
Skin (excl. basal and sq.)	882	2.1	0.08	1.5	0.05	1008	1.8	0.06	1.3	0.04
Melanomas of the skin	421	1.1	0.06	0.8	0.04	494	0.9	0.04	0.6	0.03
Breast (excl. *in situ*)	718	2.0	0.08	1.3	0.05	60689	114.0	0.47	82.7	0.35
Female Genital System	-	-	-	-	-	24829	46.8	0.30	33.9	0.22
Cervix uteri	-	-	-	-	-	7573	13.6	0.16	10.3	0.12
Corpus uteri	-	-	-	-	-	9502	18.5	0.19	13.3	0.14
Uterus, NOS	-	-	-	-	-	705	1.4	0.05	0.9	0.04
Ovary	-	-	-	-	-	5228	10.0	0.14	7.0	0.10
Male Genital System	86705	249.3	0.88	169.1	0.58	-	-	-	-	-
Prostate	85517	246.7	0.88	167.1	0.58	-	-	-	-	-
Testis	789	1.4	0.05	1.3	0.05	-	-	-	-	-
Penis	308	0.9	0.06	0.6	0.03	-	-	-	-	-
Urinary System	13169	37.7	0.35	24.3	0.22	8572	16.8	0.18	11.1	0.13
Bladder (incl. *in situ*)	5968	18.7	0.25	10.8	0.14	3482	7.1	0.12	4.1	0.07
Kidney & renal pelvis	6956	18.2	0.23	13.0	0.16	4816	9.2	0.13	6.7	0.10
Ureter	97	0.3	0.03	0.2	0.02	90	0.2	0.02	0.1	0.01
Eye	158	0.3	0.03	0.3	0.03	114	0.2	0.02	0.2	0.02
Brain & Other Nervous System	2136	4.6	0.11	3.9	0.09	2004	3.5	0.08	2.9	0.07
Brain	1966	4.2	0.10	3.6	0.08	1763	3.1	0.08	2.6	0.06
Endocrine System	1334	3.0	0.08	2.4	0.07	4169	7.3	0.11	5.9	0.09
Thyroid gland	927	2.1	0.07	1.7	0.06	3804	6.6	0.11	5.4	0.09
Lymphomas	8620	19.7	0.23	14.9	0.16	7196	13.1	0.16	9.8	0.12
Hodgkin's disease	1473	2.7	0.07	2.4	0.06	1197	1.9	0.06	1.8	0.05
Non-Hodgkin's lymphoma	7147	17.0	0.21	12.5	0.15	5999	11.2	0.15	8.1	0.11
Multiple Myeloma	4354	12.7	0.20	8.1	0.12	4696	9.4	0.14	6.0	0.09
Leukemias	4585	12.0	0.19	8.3	0.12	4108	7.7	0.12	5.5	0.09
Acute lymphocytic	582	1.0	0.04	1.1	0.04	501	0.8	0.03	0.9	0.04
Chronic lymphocytic	1226	3.7	0.11	2.3	0.07	948	1.9	0.06	1.2	0.04
Acute myeloid	1417	3.7	0.11	2.5	0.07	1402	2.6	0.07	1.9	0.05
Chronic myeloid	714	1.8	0.07	1.2	0.05	588	1.1	0.05	0.8	0.03
Other	646	1.8	0.08	1.1	0.05	669	1.3	0.05	0.8	0.03
Mesothelioma	355	1.1	0.06	0.6	0.03	119	0.2	0.02	0.2	0.02
Kaposi Sarcoma	1385	2.6	0.07	2.1	0.06	174	0.3	0.02	0.2	0.02
Ill-defined & Unknown	5900	17.0	0.23	10.9	0.14	6453	12.9	0.16	7.8	0.10
Breast *in situ* (not in All Sites)	94	0.2	0.03	0.2	0.02	12269	23.1	0.21	17.3	0.16

* Rates are per 100,000 population and were age-adjusted by five-year age groups to the 2000 U.S. standard population based on single years of age and to the World population standard.

- Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for All Sites combined.

N.B. In areas with small Latino populations, methods to indirectly identify Latinos (like NHIA) can overestimate the ethnicity-specific counts of cancer cases. Also, even small errors in Latino population estimates can affect the magnitude of the cancer rates.

Total Cancer Cases and Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates[1] for Selected Cancer Sites[2]
Selected Areas in the United States[3], 1998-2002, Non-Hispanic Black, Males
{Page 1 of 4}

	All Sites			Bladder			Brain & ONS[4]			Colon & Rectum			Gallbladder			Kidney & Renal Pelvis		
	Cases	*Rate*	*S.E.*[4]	*Cases*	*Rate*	*S.E.*	*Cases*	*Rate*	*S.E.*	*Cases*	*Rate*	*S.E.*	*Cases*	*Rate*	*S.E.*	*Cases*	*Rate*	*S.E.*
NAACCR U.S. Combined	**232873**	**652.8**	**1.42**	**5968**	**18.7**	**0.25**	**2136**	**4.6**	**0.11**	**24378**	**71.4**	**0.48**	**293**	**0.9**	**0.05**	**6956**	**18.2**	**0.23**
Alabama	10065	555.4	5.65	223	13.4	0.91	95	4.3	0.46	1088	60.8	1.88	9	0.5	0.17	301	15.8	0.93
Alaska	153	646.2	59.50	-	-	-	-	-	-	15	45.4	14.12	-	-	-	6	24.7	11.88
Arizona	1128	488.4	15.96	24	12.2	2.83	12	3.3	1.08	132	62.0	5.94	-	-	-	50	20.1	3.20
California	23118	633.6	4.40	665	20.4	0.84	204	4.3	0.32	2305	67.6	1.50	24	0.7	0.15	680	17.5	0.71
CA: Greater Bay Area	5229	615.5	8.83	147	19.3	1.66	54	5.6	0.80	507	63.0	2.92	-	-	-	135	15.5	1.38
CA: Los Angeles	10938	681.4	6.79	312	21.7	1.28	75	3.8	0.45	1104	73.5	2.32	11	0.7	0.21	328	19.7	1.14
Colorado	1280	538.4	16.99	33	16.9	3.43	11	3.2	1.00	134	57.9	5.73	-	-	-	42	16.7	2.82
Connecticut	2915	659.6	13.28	73	18.9	2.48	25	4.5	1.04	270	63.2	4.19	-	-	-	89	18.5	2.10
Delaware	1513	675.2	18.47	42	20.0	3.31	17	6.0	1.55	159	75.1	6.38	-	-	-	47	18.2	2.76
District of Columbia	5319	738.8	10.25	147	21.7	1.82	49	6.4	0.93	558	80.2	3.45	6	0.8	0.31	128	17.5	1.56
Florida	20399	612.3	4.56	470	16.5	0.81	222	5.0	0.37	2125	66.5	1.54	36	1.3	0.23	504	13.9	0.66
GA: Metropolitan Atlanta	7174	671.3	9.22	157	18.7	1.69	86	4.7	0.65	734	75.2	3.24	-	-	-	244	19.6	1.46
Hawaii	151	502.7	48.48	-	-	-	-	-	-	15	60.6	19.03	-	-	-	-	-	-
Idaho	21	403.7	102.07	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Illinois	18788	667.6	5.11	478	19.0	0.92	169	4.8	0.40	2119	79.0	1.81	23	0.8	0.18	591	19.7	0.85
Indiana	4957	634.6	9.36	138	18.7	1.66	47	4.3	0.67	586	79.1	3.40	-	-	-	180	22.1	1.72
Iowa	521	662.8	31.30	14	19.7	5.80	9	8.5	3.05	61	90.1	12.55	-	-	-	19	24.2	6.02
Kentucky	3217	708.2	12.87	107	25.1	2.50	25	4.3	0.90	336	77.0	4.34	6	1.3	0.54	113	22.9	2.23
Louisiana	14210	678.9	5.88	362	18.7	1.01	123	4.6	0.45	1609	78.0	2.01	16	0.9	0.23	478	21.5	1.02
Maine	46	594.0	100.76	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Massachusetts	3062	612.1	11.78	83	19.6	2.29	28	4.1	0.88	279	57.1	3.67	-	-	-	81	14.2	1.69
Michigan	16424	750.1	6.05	427	21.1	1.06	141	5.1	0.45	1650	78.4	2.00	18	0.8	0.19	515	22.1	1.01
MI: Metropolitan Detroit	12633	778.2	7.10	321	21.3	1.23	102	5.1	0.53	1270	81.6	2.36	12	0.7	0.21	380	22.4	1.18
Minnesota	1076	679.1	24.67	34	24.8	5.02	12	3.2	1.02	96	68.4	8.18	-	-	-	49	25.4	4.49
Missouri	6185	656.8	8.66	165	18.5	1.50	61	4.9	0.66	711	79.2	3.08	8	1.0	0.37	217	21.4	1.51

N.B. In areas with small Latino populations, methods to indirectly identify Latinos (like NHIA) can overestimate the ethnicity-specific counts of cancer cases. Also, even small errors in Latino population estimates can affect the magnitude of the cancer rates.

Total Cancer Cases and Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates[1] for Selected Cancer Sites[2]
Selected Areas in the United States[3], 1998-2002, Non-Hispanic Black, Males
{Page 2 of 4}

	All Sites			Bladder			Brain & ONS[4]			Colon & Rectum			Gallbladder			Kidney & Renal Pelvis		
	Cases	*Rate*	*S.E.*[4]	*Cases*	*Rate*	*S.E.*	*Cases*	*Rate*	*S.E.*	*Cases*	*Rate*	*S.E.*	*Cases*	*Rate*	*S.E.*	*Cases*	*Rate*	*S.E.*
Montana	26	841.9	219.42	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Nebraska	545	610.6	28.34	6	8.2	3.50	7	3.5	1.35	57	67.5	9.68	-	-	-	22	22.0	4.97
Nevada	1135	543.1	18.53	42	28.0	5.00	15	5.6	1.64	138	71.4	7.14	-	-	-	45	17.3	2.78
New Jersey	12652	715.0	6.76	312	20.0	1.22	111	5.0	0.50	1325	79.2	2.34	15	1.0	0.28	345	18.7	1.06
New Mexico	316	540.5	32.10	8	16.0	5.94	-	-	-	31	51.0	9.63	-	-	-	6	8.3	3.41
New York	27880	619.3	3.93	667	17.1	0.71	268	4.7	0.31	2719	64.1	1.31	39	1.0	0.16	662	13.7	0.56
Ohio	13221	619.9	5.58	386	20.1	1.07	129	4.9	0.45	1413	68.0	1.88	11	0.5	0.16	403	18.0	0.92
Oklahoma	2297	623.3	13.52	61	17.7	2.36	24	4.7	1.04	278	76.6	4.77	-	-	-	78	19.8	2.34
Oregon	563	668.2	31.08	11	19.4	6.35	-	-	-	55	73.0	10.75	-	-	-	29	30.1	5.93
Pennsylvania	14660	721.6	6.13	418	22.3	1.13	98	4.0	0.43	1522	78.1	2.07	23	1.3	0.27	493	23.0	1.06
Rhode Island	376	588.9	33.28	12	18.7	5.82	-	-	-	27	49.2	10.33	-	-	-	14	17.5	4.81
South Carolina	12436	678.2	6.30	266	16.4	1.05	105	4.5	0.46	1335	73.6	2.09	19	1.3	0.30	348	17.9	0.99
Utah	104	540.9	59.99	-	-	-	-	-	-	11	69.2	24.02	-	-	-	-	-	-
Washington	1763	652.1	17.23	56	20.7	3.15	11	3.0	1.08	166	61.3	5.36	-	-	-	55	18.2	2.68
WA: Seattle/Puget Sound	1607	667.7	18.59	51	21.4	3.46	11	3.4	1.23	147	60.7	5.72	-	-	-	55	20.0	2.95
West Virginia	677	615.3	23.92	22	20.6	4.42	-	-	-	73	65.3	7.75	-	-	-	22	18.6	4.03
Wisconsin	2500	731.7	16.40	47	16.0	2.67	18	2.8	0.71	235	72.7	5.34	-	-	-	85	22.1	2.77

1. Rates are per 100,000 population and were age-adjusted by five-year age groups to the 2000 U.S. standard population based on single years of age. Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for the U.S. combined.
2. Selected cancer sites by gender among the U.S. total population and the U.S. Hispanic/Latino population.
3. See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.
4. Abbreviations: S.E. - Standard Error; ONS - Other Nervous System

N.B. In areas with small Latino populations, methods to indirectly identify Latinos (like NHIA) can overestimate the ethnicity-specific counts of cancer cases. Also, even small errors in Latino population estimates can affect the magnitude of the cancer rates.

Total Cancer Cases and Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates[1] for Selected Cancer Sites[2]
Selected Areas in the United States[3], 1998-2002, Non-Hispanic Black, Males
{Page 3 of 4}

	Liver			Lung & Bronchus			NH[4] Lymphoma			Prostate			Stomach			Testis		
	Cases	*Rate*	*S.E.*[4]	*Cases*	*Rate*	*S.E.*	*Cases*	*Rate*	*S.E.*	*Cases*	*Rate*	*S.E.*	*Cases*	*Rate*	*S.E.*	*Cases*	*Rate*	*S.E.*
NAACCR U.S. Combined	**4045**	**10.3**	**0.17**	**39896**	**114.4**	**0.60**	**7147**	**17.0**	**0.21**	**85517**	**246.7**	**0.88**	**6104**	**18.3**	**0.25**	**789**	**1.4**	**0.05**
Alabama	112	5.8	0.56	1960	109.7	2.52	246	12.0	0.78	3448	195.7	3.38	320	17.9	1.02	34	1.4	0.24
Alaska	-	-	-	20	95.6	23.59	8	22.5	9.25	66	267.4	36.54	-	-	-	-	-	-
Arizona	18	6.5	1.58	183	83.3	6.67	33	11.2	2.21	368	162.9	9.04	30	16.3	3.27	-	-	-
California	481	12.1	0.58	3659	104.0	1.81	760	18.1	0.70	8888	245.3	2.72	577	17.0	0.75	98	1.7	0.18
CA: Greater Bay Area	129	13.4	1.21	890	106.7	3.69	192	20.1	1.51	1874	222.6	5.29	153	19.5	1.63	32	2.6	0.46
CA: Los Angeles	208	12.6	0.91	1683	108.0	2.74	325	18.6	1.08	4393	271.2	4.24	253	16.8	1.10	29	1.4	0.28
Colorado	27	10.3	2.19	183	87.7	7.16	49	17.2	2.96	480	200.1	10.00	32	14.9	2.96	8	1.5	0.53
Connecticut	51	11.2	1.68	440	104.1	5.37	87	16.9	2.02	1138	258.2	8.18	94	25.1	2.84	13	1.7	0.49
Delaware	15	5.9	1.60	233	107.8	7.49	37	13.7	2.39	577	261.0	11.41	47	23.0	3.66	-	-	-
District of Columbia	93	12.6	1.31	885	122.2	4.15	171	23.3	1.80	1923	267.3	6.16	117	16.8	1.58	7	0.9	0.33
Florida	310	8.0	0.49	3306	102.2	1.88	708	16.7	0.68	7693	238.9	2.87	599	19.1	0.84	69	1.4	0.18
GA: Metropolitan Atlanta	103	7.7	0.93	1104	112.4	3.88	262	16.0	1.22	2675	272.8	5.90	161	17.1	1.57	37	1.3	0.21
Hawaii	-	-	-	20	72.0	16.66	7	10.2	4.54	55	216.7	31.85	-	-	-	-	-	-
Idaho	-	-	-	-	-	-	-	-	-	9	167.7	62.35	-	-	-	-	-	-
Illinois	313	10.6	0.63	3557	127.2	2.22	523	15.8	0.73	6199	228.5	3.03	497	18.7	0.89	59	1.3	0.17
Indiana	93	11.0	1.18	950	123.5	4.13	145	16.3	1.42	1587	209.0	5.40	106	14.0	1.42	14	1.2	0.33
Iowa	10	11.5	4.05	102	140.1	14.74	17	15.4	4.28	154	204.8	17.31	15	16.1	4.42	-	-	-
Kentucky	36	7.0	1.19	659	149.3	5.95	107	20.6	2.09	1052	241.1	7.59	68	14.7	1.86	14	1.9	0.50
Louisiana	208	8.8	0.63	2831	135.3	2.61	363	15.3	0.84	4669	233.4	3.50	408	20.5	1.05	43	1.4	0.22
Maine	-	-	-	7	97.5	38.93	-	-	-	13	197.2	61.93	-	-	-	-	-	-
Massachusetts	53	8.7	1.26	389	80.4	4.34	113	18.3	1.86	1284	267.2	7.80	85	19.4	2.26	13	1.5	0.41
Michigan	275	11.2	0.70	2780	128.5	2.51	423	16.9	0.86	6452	302.9	3.87	377	18.4	0.99	41	1.2	0.20
MI: Metropolitan Detroit	215	12.0	0.84	2087	129.7	2.90	319	17.8	1.03	5106	321.3	4.59	284	18.3	1.13	29	1.3	0.24
Minnesota	39	22.4	3.95	181	126.0	10.72	47	24.8	4.54	312	233.2	14.77	21	13.8	3.55	13	2.2	0.64
Missouri	134	13.1	1.17	1233	133.1	3.91	164	15.2	1.25	1974	215.3	4.99	147	16.9	1.45	14	0.9	0.25

N.B. In areas with small Latino populations, methods to indirectly identify Latinos (like NHIA) can overestimate the ethnicity-specific counts of cancer cases. Also, even small errors in Latino population estimates can affect the magnitude of the cancer rates.

Total Cancer Cases and Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates[1] for Selected Cancer Sites[2]
Selected Areas in the United States[3], 1998-2002, Non-Hispanic Black, Males
{Page 4 of 4}

	Liver			Lung & Bronchus			NH[4] Lymphoma			Prostate			Stomach			Testis		
	Cases	*Rate*	*S.E.*[4]	*Cases*	*Rate*	*S.E.*	*Cases*	*Rate*	*S.E.*	*Cases*	*Rate*	*S.E.*	*Cases*	*Rate*	*S.E.*	*Cases*	*Rate*	*S.E.*
Montana	-	-	-	6	175.1	89.90	-	-	-	9	237.0	82.85	-	-	-	-	-	-
Nebraska	11	13.8	4.72	125	146.7	13.99	16	13.5	3.83	174	201.7	16.30	8	10.0	3.83	-	-	-
Nevada	21	10.5	2.71	215	104.0	7.85	26	9.1	1.93	376	181.3	10.68	30	13.4	2.65	-	-	-
New Jersey	199	10.3	0.78	1942	112.4	2.70	443	20.6	1.04	5016	288.6	4.30	332	20.5	1.20	37	1.3	0.22
New Mexico	6	10.8	4.51	55	102.8	14.61	8	13.8	5.22	129	219.0	20.06	-	-	-	-	-	-
New York	574	11.6	0.51	3999	90.4	1.51	1031	19.1	0.63	11319	258.3	2.56	785	19.3	0.74	95	1.5	0.15
Ohio	242	10.8	0.72	2627	125.8	2.53	383	16.4	0.88	4570	217.0	3.30	328	16.4	0.94	43	1.5	0.24
Oklahoma	32	8.4	1.56	453	127.5	6.18	69	16.6	2.12	775	218.5	8.07	54	14.9	2.10	10	1.8	0.63
Oregon	12	10.9	3.23	113	140.3	14.25	24	22.6	5.41	174	215.6	17.71	18	23.1	6.05	-	-	-
Pennsylvania	327	14.8	0.84	2624	132.0	2.64	437	19.5	0.97	5268	260.9	3.67	362	19.6	1.07	50	1.8	0.27
Rhode Island	-	-	-	66	106.8	14.47	16	19.8	5.18	143	235.0	21.25	8	12.5	4.51	-	-	-
South Carolina	135	6.8	0.61	2073	113.2	2.57	273	13.5	0.86	4742	269.8	4.04	364	20.4	1.11	27	1.0	0.19
Utah	-	-	-	-	-	-	-	-	-	44	238.8	37.98	-	-	-	-	-	-
Washington	45	15.2	2.57	320	128.5	7.88	60	16.6	2.41	680	260.6	10.82	35	15.0	2.88	9	1.6	0.55
WA: Seattle/Puget Sound	42	16.1	2.86	289	131.0	8.50	52	15.8	2.48	621	265.3	11.57	34	16.8	3.29	9	1.8	0.61
West Virginia	10	7.3	2.35	123	112.5	10.24	9	7.5	2.55	249	234.7	14.98	10	9.4	3.02	-	-	-
Wisconsin	50	11.4	1.71	466	146.5	7.56	77	18.7	2.36	833	250.7	9.39	56	15.5	2.30	15	2.2	0.59

1. Rates are per 100,000 population and were age-adjusted by five-year age groups to the 2000 U.S. standard population based on single years of age. Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for the U.S. combined.
2. Selected cancer sites by gender among the U.S. total population and the U.S. Hispanic/Latino population.
3. See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.
4. Abbreviations: S.E. - Standard Error; NH - Non-Hodgkin.

N.B. In areas with small Latino populations, methods to indirectly identify Latinos (like NHIA) can overestimate the ethnicity-specific counts of cancer cases. Also, even small errors in Latino population estimates can affect the magnitude of the cancer rates.

Total Cancer Cases and Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates[1] for Selected Cancer Sites[2]
Selected Areas in the United States[3], 1998-2002, Non-Hispanic Black, Females
{Page 1 of 4}

	All Sites			Breast			Cervix Uteri			Colon & Rectum			Corpus uteri & Uterus, NOS [4]			Gallbladder		
	Cases	*Rate*	*S.E.*[4]	*Cases*	*Rate*	*S.E.*	*Cases*	*Rate*	*S.E.*	*Cases*	*Rate*	*S.E.*	*Cases*	*Rate*	*S.E.*	*Cases*	*Rate*	*S.E.*
NAACCR U.S. Combined	**203771**	**391.3**	**0.87**	**60689**	**114.0**	**0.47**	**7573**	**13.6**	**0.16**	**27445**	**54.6**	**0.33**	**10207**	**19.8**	**0.20**	**871**	**1.7**	**0.06**
Alabama	9166	333.4	3.50	2826	103.2	1.95	388	13.8	0.71	1328	48.6	1.34	452	16.8	0.79	37	1.3	0.22
Alaska	109	402.9	45.56	44	147.9	26.55	6	14.6	6.34	11	64.0	20.81	7	27.0	12.52	-	-	-
Arizona	823	307.5	11.00	261	93.7	5.94	20	7.0	1.62	113	45.4	4.36	27	10.7	2.10	-	-	-
California	19024	389.8	2.85	6062	120.5	1.56	508	9.5	0.43	2498	53.6	1.08	895	18.4	0.62	54	1.2	0.16
CA: Greater Bay Area	4447	394.2	5.93	1354	118.0	3.22	112	9.3	0.88	519	46.9	2.07	226	20.2	1.35	14	1.3	0.34
CA: Los Angeles	9138	400.5	4.21	2922	125.9	2.34	245	10.2	0.65	1289	57.8	1.62	445	19.4	0.92	30	1.3	0.25
Colorado	987	343.4	11.31	318	103.0	5.96	28	7.8	1.53	120	45.0	4.24	42	14.6	2.32	-	-	-
Connecticut	2570	395.2	7.96	813	121.4	4.34	82	11.6	1.31	375	60.6	3.19	124	19.0	1.73	12	2.1	0.62
Delaware	1341	426.3	11.84	396	119.1	6.09	44	12.3	1.88	177	60.7	4.62	65	21.3	2.67	-	-	-
District of Columbia	4613	424.9	6.32	1320	125.9	3.50	148	14.8	1.23	676	59.3	2.30	233	21.2	1.40	17	1.5	0.38
Florida	17141	367.7	2.85	4953	102.9	1.48	815	16.0	0.57	2299	52.0	1.10	970	21.5	0.70	82	1.9	0.21
GA: Metropolitan Atlanta	6638	379.5	4.98	2241	115.8	2.62	246	12.2	0.85	838	55.2	2.00	283	16.9	1.06	14	1.1	0.30
Hawaii	67	290.7	41.20	21	95.0	22.95	6	11.8	5.52	9	39.1	14.49	-	-	-	-	-	-
Idaho	9	189.4	67.05	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Illinois	17815	427.4	3.23	5314	124.6	1.72	767	17.1	0.62	2410	60.0	1.23	832	20.0	0.70	63	1.5	0.19
Indiana	4509	410.2	6.14	1334	119.5	3.29	143	12.2	1.02	588	55.2	2.28	218	19.9	1.35	25	2.4	0.48
Iowa	409	422.0	21.23	126	128.5	11.59	12	10.5	3.08	51	56.8	8.01	9	8.8	2.96	-	-	-
Kentucky	2960	449.7	8.31	865	130.9	4.47	110	15.8	1.52	464	71.1	3.32	108	16.5	1.59	17	2.6	0.64
Louisiana	12293	395.6	3.59	3814	120.6	1.96	507	15.3	0.68	1705	56.5	1.37	546	18.1	0.78	58	1.9	0.25
Maine	28	360.1	74.28	8	90.6	33.01	-	-	-	7	110.8	44.74	-	-	-	-	-	-
Massachusetts	2497	350.4	7.14	766	102.6	3.77	95	11.9	1.26	295	44.9	2.65	132	19.0	1.67	18	2.6	0.62
Michigan	13521	432.8	3.74	3857	121.0	1.96	423	12.7	0.62	1822	60.3	1.42	653	21.1	0.83	55	1.8	0.25
MI: Metropolitan Detroit	10420	437.7	4.30	2999	123.9	2.27	333	13.2	0.73	1428	61.5	1.63	509	21.5	0.96	45	1.9	0.29
Minnesota	799	398.6	15.60	239	109.3	7.87	30	10.5	2.24	89	52.5	5.94	33	17.5	3.26	-	-	-
Missouri	5828	413.8	5.45	1708	119.2	2.90	214	14.3	0.98	837	60.7	2.11	281	20.2	1.21	28	2.0	0.39

N.B. In areas with small Latino populations, methods to indirectly identify Latinos (like NHIA) can overestimate the ethnicity-specific counts of cancer cases. Also, even small errors in Latino population estimates can affect the magnitude of the cancer rates.

Total Cancer Cases and Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates[1] for Selected Cancer Sites[2]
Selected Areas in the United States[3], 1998-2002, Non-Hispanic Black, Females
{Page 2 of 4}

	All Sites			Breast			Cervix Uteri			Colon & Rectum			Corpus uteri & Uterus, NOS [4]			Gallbladder		
	Cases	*Rate*	*S.E.*[4]	*Cases*	*Rate*	*S.E.*	*Cases*	*Rate*	*S.E.*	*Cases*	*Rate*	*S.E.*	*Cases*	*Rate*	*S.E.*	*Cases*	*Rate*	*S.E.*
Montana	6	199.9	88.55	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Nebraska	458	370.5	17.68	138	107.5	9.33	11	6.9	2.12	61	52.3	6.79	22	18.4	4.02	-	-	-
Nevada	911	363.3	12.53	276	103.2	6.42	24	8.1	1.66	107	45.4	4.57	33	12.8	2.30	-	-	-
New Jersey	10811	415.3	4.04	3154	117.7	2.12	434	15.7	0.76	1449	58.0	1.54	604	23.2	0.96	49	2.0	0.28
New Mexico	180	283.8	21.32	60	95.2	12.36	7	9.5	3.61	20	33.6	7.57	10	15.8	5.02	-	-	-
New York	26676	373.2	2.30	7450	102.1	1.19	1150	15.5	0.46	3560	51.4	0.87	1651	22.9	0.57	147	2.1	0.18
Ohio	11946	391.2	3.59	3609	117.1	1.95	300	9.5	0.55	1595	53.2	1.33	564	18.5	0.78	52	1.7	0.24
Oklahoma	1982	378.7	8.59	636	120.2	4.80	74	13.0	1.53	252	49.0	3.11	82	16.2	1.80	6	1.1	0.47
Oregon	391	397.7	20.45	128	123.3	11.07	12	10.9	3.22	49	54.1	7.79	12	12.3	3.55	-	-	-
Pennsylvania	13051	428.8	3.76	3609	119.0	1.98	411	13.4	0.66	1705	56.1	1.36	591	19.4	0.80	64	2.1	0.26
Rhode Island	321	369.1	21.16	89	100.1	10.82	-	-	-	34	43.7	7.69	15	17.9	4.67	-	-	-
South Carolina	9893	358.4	3.62	3030	108.5	1.98	400	14.0	0.70	1398	51.6	1.38	555	20.6	0.88	39	1.4	0.23
Utah	49	267.7	39.69	15	79.3	21.15	-	-	-	7	43.5	16.63	-	-	-	-	-	-
Washington	1213	394.9	11.80	372	111.0	5.98	31	8.8	1.67	133	49.0	4.34	52	19.0	2.69	-	-	-
WA: Seattle/Puget Sound	1120	403.8	12.58	346	112.7	6.32	28	9.0	1.78	130	53.3	4.78	47	19.5	2.90	-	-	-
West Virginia	639	394.1	15.97	186	117.9	8.84	18	12.2	2.93	109	64.5	6.34	22	14.3	3.08	-	-	-
Wisconsin	2097	421.7	9.68	646	124.9	5.15	98	15.1	1.60	249	55.9	3.68	79	17.0	1.98	-	-	-

1. Rates are per 100,000 population and were age-adjusted by five-year age groups to the 2000 U.S. standard population based on single years of age. Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for the U.S. combined.
2. Selected cancer sites by gender among the U.S. total population and the U.S. Hispanic/Latino population.
3. See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.
4. Abbreviations: S.E. - Standard Error; NOS - Not Otherwise Specified.

N.B. In areas with small Latino populations, methods to indirectly identify Latinos (like NHIA) can overestimate the ethnicity-specific counts of cancer cases. Also, even small errors in Latino population estimates can affect the magnitude of the cancer rates.

Total Cancer Cases and Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates[1] for Selected Cancer Sites[2]
Selected Areas in the United States[3], 1998-2002, Non-Hispanic Black, Females
{Page 3 of 4}

	Liver			Lung & Bronchus			NH[4] Lymphoma			Ovary			Stomach			Thyroid		
	Cases	*Rate*	*S.E.*[4]	*Cases*	*Rate*	*S.E.*	*Cases*	*Rate*	*S.E.*	*Cases*	*Rate*	*S.E.*	*Cases*	*Rate*	*S.E.*	*Cases*	*Rate*	*S.E.*
NAACCR U.S. Combined	**1594**	**3.1**	**0.08**	**26445**	**52.3**	**0.32**	**5999**	**11.2**	**0.15**	**5228**	**10.0**	**0.14**	**4666**	**9.4**	**0.14**	**3804**	**6.6**	**0.11**
Alabama	65	2.4	0.29	998	37.0	1.18	240	8.6	0.56	269	9.8	0.60	197	7.1	0.51	171	6.0	0.46
Alaska	-	-	-	9	36.6	13.21	-	-	-	-	-	-	-	-	-	7	15.0	6.32
Arizona	6	2.4	1.00	113	44.9	4.30	26	8.8	1.79	18	6.7	1.59	25	10.1	2.07	24	7.3	1.53
California	167	3.5	0.27	2552	54.1	1.08	590	11.9	0.50	457	9.2	0.43	430	9.4	0.45	294	5.4	0.32
CA: Greater Bay Area	42	3.8	0.58	679	61.5	2.37	143	12.6	1.06	120	10.5	0.96	105	9.6	0.94	55	4.6	0.62
CA: Los Angeles	76	3.4	0.40	1162	52.0	1.53	283	12.2	0.73	207	9.1	0.63	213	9.7	0.66	125	5.2	0.47
Colorado	8	2.9	1.07	130	51.0	4.56	34	12.4	2.22	13	4.8	1.38	15	5.3	1.42	35	9.9	1.73
Connecticut	16	2.3	0.59	326	52.5	2.96	87	12.7	1.40	46	6.8	1.03	46	7.7	1.15	48	6.3	0.92
Delaware	-	-	-	182	61.2	4.58	51	15.4	2.19	24	7.2	1.49	36	12.6	2.12	49	13.2	1.92
District of Columbia	44	4.0	0.61	612	55.2	2.25	106	9.9	0.97	120	11.2	1.03	117	10.0	0.93	105	10.6	1.04
Florida	120	2.6	0.24	1822	40.8	0.97	590	11.8	0.50	496	10.6	0.49	412	9.6	0.48	341	6.3	0.34
GA: Metropolitan Atlanta	48	3.0	0.46	718	46.3	1.81	216	10.9	0.80	199	11.5	0.86	132	8.7	0.79	154	6.4	0.56
Hawaii	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Idaho	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Illinois	166	4.1	0.32	2497	61.1	1.23	444	10.5	0.50	416	9.8	0.48	378	9.7	0.50	274	6.1	0.37
Indiana	31	2.9	0.52	712	66.6	2.50	113	10.1	0.95	102	9.1	0.91	81	7.7	0.86	72	5.9	0.70
Iowa	-	-	-	62	69.3	8.86	7	6.7	2.54	11	10.5	3.25	7	8.0	3.05	11	9.3	2.85
Kentucky	16	2.4	0.59	500	78.0	3.50	70	10.5	1.26	75	11.1	1.29	48	7.4	1.07	43	6.0	0.93
Louisiana	79	2.6	0.29	1479	49.1	1.28	332	10.5	0.58	311	10.0	0.57	328	11.0	0.61	207	6.2	0.43
Maine	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Massachusetts	20	2.9	0.67	308	46.0	2.65	90	12.3	1.33	42	5.9	0.93	53	8.4	1.17	64	7.4	0.94
Michigan	103	3.3	0.33	1989	65.3	1.47	392	12.2	0.62	337	10.8	0.59	315	10.5	0.59	238	7.1	0.46
MI: Metropolitan Detroit	81	3.4	0.38	1538	66.1	1.69	285	11.8	0.70	257	10.9	0.68	251	10.8	0.69	178	7.0	0.53
Minnesota	7	4.1	1.66	111	63.8	6.39	24	8.8	2.01	16	5.6	1.57	11	7.7	2.50	20	4.9	1.21
Missouri	48	3.5	0.51	858	62.5	2.14	170	12.0	0.92	131	9.4	0.82	102	7.4	0.73	93	6.1	0.63

N.B. In areas with small Latino populations, methods to indirectly identify Latinos (like NHIA) can overestimate the ethnicity-specific counts of cancer cases. Also, even small errors in Latino population estimates can affect the magnitude of the cancer rates.

Total Cancer Cases and Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates[1] for Selected Cancer Sites[2]
Selected Areas in the United States[3], 1998-2002, Non-Hispanic Black, Females
{Page 4 of 4}

	Liver			Lung & Bronchus			NH[4] Lymphoma			Ovary			Stomach			Thyroid		
	Cases	*Rate*	*S.E.*[4]	*Cases*	*Rate*	*S.E.*	*Cases*	*Rate*	*S.E.*	*Cases*	*Rate*	*S.E.*	*Cases*	*Rate*	*S.E.*	*Cases*	*Rate*	*S.E.*
Montana	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Nebraska	-	-	-	71	60.8	7.32	14	11.8	3.24	8	6.1	2.21	11	9.5	2.90	-	-	-
Nevada	13	5.0	1.47	144	61.7	5.33	22	9.2	2.07	31	12.4	2.30	14	6.6	1.83	22	6.8	1.47
New Jersey	67	2.7	0.33	1347	53.0	1.46	365	13.4	0.71	295	11.3	0.66	244	9.8	0.64	201	7.0	0.50
New Mexico	-	-	-	16	25.8	6.47	-	-	-	-	-	-	6	9.3	3.80	6	9.0	3.69
New York	257	3.7	0.23	3030	43.0	0.79	870	11.8	0.40	812	11.3	0.40	744	11.0	0.40	512	6.7	0.30
Ohio	85	2.8	0.31	1846	61.2	1.43	334	10.8	0.60	254	8.3	0.52	254	8.6	0.54	168	5.2	0.40
Oklahoma	14	2.6	0.71	298	59.9	3.49	53	10.0	1.40	61	12.0	1.55	35	6.8	1.15	24	3.9	0.82
Oregon	-	-	-	49	52.8	7.61	9	9.6	3.23	9	10.2	3.41	10	10.5	3.36	7	6.2	2.40
Pennsylvania	111	3.7	0.35	2071	68.4	1.50	345	11.3	0.61	296	9.7	0.56	294	9.6	0.56	361	11.8	0.62
Rhode Island	-	-	-	57	67.5	9.08	6	6.1	2.62	8	8.4	3.10	13	16.0	4.52	13	11.4	3.24
South Carolina	53	1.9	0.26	969	36.0	1.16	257	9.1	0.57	271	9.8	0.60	230	8.5	0.56	160	5.4	0.43
Utah	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Washington	12	4.4	1.30	151	56.6	4.66	48	13.6	2.08	28	9.0	1.76	29	10.5	2.00	32	8.0	1.49
WA: Seattle/Puget Sound	11	4.4	1.37	141	58.8	5.02	40	12.5	2.11	23	8.0	1.74	28	11.1	2.17	30	8.4	1.61
West Virginia	-	-	-	86	52.7	5.80	15	10.0	2.64	21	13.0	2.88	9	4.8	1.64	13	9.3	2.60
Wisconsin	15	3.2	0.84	317	66.9	3.88	70	13.1	1.66	41	7.9	1.31	39	8.8	1.49	26	4.4	0.90

1. Rates are per 100,000 population and were age-adjusted by five-year age groups to the 2000 U.S. standard population based on single years of age. Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for the U.S. combined.
2. Selected cancer sites by gender among the U.S. total population and the U.S. Hispanic/Latino population.
3. See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.
4. Abbreviations: S.E. - Standard Error; NH - Non-Hodgkin.

N.B. In areas with small Latino populations, methods to indirectly identify Latinos (like NHIA) can overestimate the ethnicity-specific counts of cancer cases. Also, even small errors in Latino population estimates can affect the magnitude of the cancer rates.

Average Annual Age-adjusted (2000 U.S. Standard) NAACCR (U.S.) Combined[2] Cancer Incidence Rates [1] and Rate Ratios for Selected Cancer Sites[3] by Race/Ethnicity and Gender, 1998-2002

Males

	Rates			Rate Ratios	
Cancer Site	**Hispanic**	**Non-Hispanic White**	**Non-Hispanic Black**	**Hispanic:Non-Hispanic White**	**Hispanic:Non-Hispanic Black**
All Sites	445.6	562.5	652.8	0.8	0.7
Bladder	22.5	42.2	18.7	0.5	1.2
Brain & ONS[4]	6.2	8.6	4.6	0.7	1.3
Colon & Rectum	54.0	66.3	71.4	0.8	0.8
Gallbladder	1.4	0.8	0.9	1.9	1.6
Kidney & Renal Pelvis	15.5	17.3	18.2	0.9	0.9
Liver	12.6	5.4	10.3	2.3	1.2
Lung & Bronchus	53.4	91.7	114.4	0.6	0.5
NH[4] Lymphoma	20.0	23.2	17.0	0.9	1.2
Prostate	142.6	156.4	246.7	0.9	0.6
Stomach	16.1	9.4	18.3	1.7	0.9
Testis	3.4	6.5	1.4	0.5	2.4

Females

	Rates			Rate Ratios	
Cancer Site	**Hispanic**	**Non-Hispanic White**	**Non-Hispanic Black**	**Hispanic:Non-Hispanic White**	**Hispanic:Non-Hispanic Black**
All Sites	328.4	431.1	391.3	0.8	0.8
Breast	94.5	136.6	114.0	0.7	0.8
Cervix Uteri	14.6	8.1	13.6	1.8	1.1
Colon & Rectum	38.6	47.8	54.6	0.8	0.7
Corpus uteri & Uterus NOS[4]	18.7	25.7	19.8	0.7	0.9
Gallbladder	3.1	1.3	1.7	2.4	1.8
Liver	4.5	1.8	3.1	2.5	1.4
Lung & Bronchus	27.0	58.0	52.3	0.5	0.5
NH[4] Lymphoma	14.3	16.6	11.2	0.9	1.3
Ovary	11.4	14.7	10.0	0.8	1.1
Stomach	9.1	4.1	9.4	2.2	1.0
Thyroid	10.8	11.4	6.6	1.0	1.6

1. Rates are per 100,000 population and were age-adjusted by five-year age groups to the 2000 U.S. standard population based on single years of age. Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for the U.S. combined.
2. See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.
3. Selected cancer sites by gender among the U.S. total population and the U.S. Hispanic/Latino population.
4. Abbreviations: ONS - Other Nervous System; NOS - Not Otherwise Specified; NH - Non-Hodgkin

N.B. In areas with small Latino populations, methods to indirectly identify Latinos (like NHIA) can overestimate the ethnicity-specific counts of cancer cases. Also, even small errors in Latino population estimates can affect the magnitude of the cancer rates.