





Section V

Comparative Graphs of Average-annual Age-adjusted Cancer Incidence Rates by Registry Among Hispanic Latino Populations, Selected Sites



**Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates¹ and 95% Confidence Intervals²
Selected Areas in the United States³, 2000-2004, Hispanic/Latino, All Races, Males**




¹ Rates are per 100,000 and are adjusted to the 2000 U.S. population standard. Rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the rates for the U.S. combined.

² See <http://smm.sagepub.com/cgi/content/abstract/15/6/547> for the methodology used in the calculation of the age-adjusted 95% confidence intervals.

³ See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.

N.B. In areas with small Latino populations, methods to indirectly identify Latinos (like NHIA) can overestimate the ethnicity-specific counts of cancer cases. Also, even small errors in Latino population estimates can affect the magnitude of the cancer rates.

**Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates¹ and 95% Confidence Intervals²
Selected Areas in the United States³, 2000-2004, Hispanic/Latina, All Races, Females**




¹ Rates are per 100,000 and are adjusted to the 2000 U.S. population standard. Rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the rates for the U.S. combined.

² See <http://smm.sagepub.com/cgi/content/abstract/15/6/547> for the methodology used in the calculation of the age-adjusted 95% confidence intervals.

³ See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.

N.B. In areas with small Latino populations, methods to indirectly identify Latinos (like NHIA) can overestimate the ethnicity-specific counts of cancer cases. Also, even small errors in Latino population estimates can affect the magnitude of the cancer rates.

**Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates¹ and 95% Confidence Intervals²
Selected Areas in the United States³, 2000-2004, Hispanic/Latino, All Races, Males**



¹ Rates are per 100,000 and are adjusted to the 2000 U.S. population standard. Rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the rates for the U.S. combined.


² See <http://smm.sagepub.com/cgi/content/abstract/15/6/547> for the methodology used in the calculation of the age-adjusted 95% confidence intervals.

³ See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics

N.B. In areas with small Latino populations, methods to indirectly identify Latinos (like NHIA) can overestimate the ethnicity-specific counts of cancer cases. Also, even small errors in Latino population estimates can affect the magnitude of the cancer rates.

**Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates¹ and 95% Confidence Intervals²
Selected Areas in the United States³, 2000-2004, Hispanic/Latina, All Races, Females**

Bladder



¹ Rates are per 100,000 and are adjusted to the 2000 U.S. population standard. Rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the rates for the U.S. combined.


² See <http://smm.sagepub.com/cgi/content/abstract/15/6/547> for the methodology used in the calculation of the age-adjusted 95% confidence intervals.

³ See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics

N.B. In areas with small Latino populations, methods to indirectly identify Latinos (like NHIA) can overestimate the ethnicity-specific counts of cancer cases. Also, even small errors in Latino population estimates can affect the magnitude of the cancer rates.

Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates¹ and 95% Confidence Intervals²
 Selected Areas in the United States³, 2000-2004, Hispanic/Latino, All Races, Males

Brain and Other Nervous System



¹ Rates are per 100,000 and are adjusted to the 2000 U.S. population standard. Rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the rates for the U.S. combined.


² See <http://smm.sagepub.com/cgi/content/abstract/15/6/547> for the methodology used in the calculation of the age-adjusted 95% confidence intervals.

³ See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics

N.B. In areas with small Latino populations, methods to indirectly identify Latinos (like NHIA) can overestimate the ethnicity-specific counts of cancer cases. Also, even small errors in Latino population estimates can affect the magnitude of the cancer rates.

**Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates¹ and 95% Confidence Intervals²
Selected Areas in the United States³, 2000-2004, Hispanic/Latina, All Races, Females**

Brain and Other Nervous System




¹ Rates are per 100,000 and are adjusted to the 2000 U.S. population standard. Rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the rates for the U.S. combined.

² See <http://smm.sagepub.com/cgi/content/abstract/15/6/547> for the methodology used in the calculation of the age-adjusted 95% confidence intervals.

³ See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics

N.B. In areas with small Latino populations, methods to indirectly identify Latinos (like NHIA) can overestimate the ethnicity-specific counts of cancer cases. Also, even small errors in Latino population estimates can affect the magnitude of the cancer rates.

Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates¹ and 95% Confidence Intervals²
 Selected Areas in the United States³, 2000-2004, Hispanic/Latina, All Races, Females




¹ Rates are per 100,000 and are adjusted to the 2000 U.S. population standard. Rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the rates for the U.S. combined.

² See <http://smm.sagepub.com/cgi/content/abstract/15/6/547> for the methodology used in the calculation of the age-adjusted 95% confidence intervals.

³ See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics

N.B. In areas with small Latino populations, methods to indirectly identify Latinos (like NHIA) can overestimate the ethnicity-specific counts of cancer cases. Also, even small errors in Latino population estimates can affect the magnitude of the cancer rates.

**Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates¹ and 95% Confidence Intervals²
Selected Areas in the United States³, 2000-2004, Hispanic/Latina, All Races, Females**



¹ Rates are per 100,000 and are adjusted to the 2000 U.S. population standard. Rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the rates for the U.S. combined.


² See <http://smm.sagepub.com/cgi/content/abstract/15/6/547> for the methodology used in the calculation of the age-adjusted 95% confidence intervals.

³ See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.

N.B. In areas with small Latino populations, methods to indirectly identify Latinos (like NHIA) can overestimate the ethnicity-specific counts of cancer cases. Also, even small errors in Latino population estimates can affect the magnitude of the cancer rates.

Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates¹ and 95% Confidence Intervals²
 Selected Areas in the United States³, 2000-2004, Hispanic/Latino, All Races, Males

Colon and Rectum




¹ Rates are per 100,000 and are adjusted to the 2000 U.S. population standard. Rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the rates for the U.S. combined.

² See <http://smm.sagepub.com/cgi/content/abstract/15/6/547> for the methodology used in the calculation of the age-adjusted 95% confidence intervals.

³ See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.

N.B. In areas with small Latino populations, methods to indirectly identify Latinos (like NHIA) can overestimate the ethnicity-specific counts of cancer cases. Also, even small errors in Latino population estimates can affect the magnitude of the cancer rates.

**Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates¹ and 95% Confidence Intervals²
Selected Areas in the United States³, 2000-2004, Hispanic/Latina, All Races, Females**



¹ Rates are per 100,000 and are adjusted to the 2000 U.S. population standard. Rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the rates for the U.S. combined.


² See <http://smm.sagepub.com/cgi/content/abstract/15/6/547> for the methodology used in the calculation of the age-adjusted 95% confidence intervals.

³ See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.

N.B. In areas with small Latino populations, methods to indirectly identify Latinos (like NHIA) can overestimate the ethnicity-specific counts of cancer cases. Also, even small errors in Latino population estimates can affect the magnitude of the cancer rates.

This Page Intentionally Left Blank

**Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates¹ and 95% Confidence Intervals²
Selected Areas in the United States³, 2000-2004, Hispanic/Latina, All Races, Females**




¹ Rates are per 100,000 and are adjusted to the 2000 U.S. population standard. Rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the rates for the U.S. combined.
² See <http://smm.sagepub.com/cgi/content/abstract/15/6/547> for the methodology used in the calculation of the age-adjusted 95% confidence intervals.
³ See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.

N.B. In areas with small Latino populations, methods to indirectly identify Latinos (like NHIA) can overestimate the ethnicity-specific counts of cancer cases. Also, even small errors in Latino population estimates can affect the magnitude of the cancer rates.

Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates¹ and 95% Confidence Intervals²
 Selected Areas in the United States³, 2000-2004, Hispanic/Latino, All Races, Males

Gallbladder




¹ Rates are per 100,000 and are adjusted to the 2000 U.S. population standard. Rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the rates for the U.S. combined.

² See <http://smm.sagepub.com/cgi/content/abstract/15/6/547> for the methodology used in the calculation of the age-adjusted 95% confidence intervals.

³ See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.

N.B. In areas with small Latino populations, methods to indirectly identify Latinos (like NHIA) can overestimate the ethnicity-specific counts of cancer cases. Also, even small errors in Latino population estimates can affect the magnitude of the cancer rates.

**Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates¹ and 95% Confidence Intervals²
Selected Areas in the United States³, 2000-2004, Hispanic/Latina, All Races, Females**



¹ Rates are per 100,000 and are adjusted to the 2000 U.S. population standard. Rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the rates for the U.S. combined.


² See <http://smm.sagepub.com/cgi/content/abstract/15/6/547> for the methodology used in the calculation of the age-adjusted 95% confidence intervals.

³ See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.

N.B. In areas with small Latino populations, methods to indirectly identify Latinos (like NHIA) can overestimate the ethnicity-specific counts of cancer cases. Also, even small errors in Latino population estimates can affect the magnitude of the cancer rates.

Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates¹ and 95% Confidence Intervals²
 Selected Areas in the United States³, 2000-2004, Hispanic/Latino, All Races, Males

Kidney and Renal Pelvis



¹ Rates are per 100,000 and are adjusted to the 2000 U.S. population standard. Rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the rates for the U.S. combined.


² See <http://smm.sagepub.com/cgi/content/abstract/15/6/547> for the methodology used in the calculation of the age-adjusted 95% confidence intervals.

³ See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.

N.B. In areas with small Latino populations, methods to indirectly identify Latinos (like NHIA) can overestimate the ethnicity-specific counts of cancer cases. Also, even small errors in Latino population estimates can affect the magnitude of the cancer rates.

**Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates¹ and 95% Confidence Intervals²
Selected Areas in the United States³, 2000-2004, Hispanic/Latina, All Races, Females**

Kidney and Renal Pelvis




¹ Rates are per 100,000 and are adjusted to the 2000 U.S. population standard. Rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the rates for the U.S. combined.

² See <http://smm.sagepub.com/cgi/content/abstract/15/6/547> for the methodology used in the calculation of the age-adjusted 95% confidence intervals.

³ See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.

N.B. In areas with small Latino populations, methods to indirectly identify Latinos (like NHIA) can overestimate the ethnicity-specific counts of cancer cases. Also, even small errors in Latino population estimates can affect the magnitude of the cancer rates.

**Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates¹ and 95% Confidence Intervals²
Selected Areas in the United States³, 2000-2004, Hispanic/Latino, All Races, Males**




¹ Rates are per 100,000 and are adjusted to the 2000 U.S. population standard. Rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the rates for the U.S. combined.

² See <http://smm.sagepub.com/cgi/content/abstract/15/6/547> for the methodology used in the calculation of the age-adjusted 95% confidence intervals.

³ See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.

N.B. In areas with small Latino populations, methods to indirectly identify Latinos (like NHIA) can overestimate the ethnicity-specific counts of cancer cases. Also, even small errors in Latino population estimates can affect the magnitude of the cancer rates.

**Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates¹ and 95% Confidence Intervals²
Selected Areas in the United States³, 2000-2004, Hispanic/Latina, All Races, Females**




¹ Rates are per 100,000 and are adjusted to the 2000 U.S. population standard. Rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the rates for the U.S. combined.

² See <http://smm.sagepub.com/cgi/content/abstract/15/6/547> for the methodology used in the calculation of the age-adjusted 95% confidence intervals.

³ See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.

N.B. In areas with small Latino populations, methods to indirectly identify Latinos (like NHIA) can overestimate the ethnicity-specific counts of cancer cases. Also, even small errors in Latino population estimates can affect the magnitude of the cancer rates.

Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates¹ and 95% Confidence Intervals²
 Selected Areas in the United States³, 2000-2004, Hispanic/Latino, All Races, Males




¹ Rates are per 100,000 and are adjusted to the 2000 U.S. population standard. Rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the rates for the U.S. combined.

² See <http://smm.sagepub.com/cgi/content/abstract/15/6/547> for the methodology used in the calculation of the age-adjusted 95% confidence intervals.

³ See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.

N.B. In areas with small Latino populations, methods to indirectly identify Latinos (like NHIA) can overestimate the ethnicity-specific counts of cancer cases. Also, even small errors in Latino population estimates can affect the magnitude of the cancer rates.

**Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates¹ and 95% Confidence Intervals²
Selected Areas in the United States³, 2000-2004, Hispanic/Latina, All Races, Females**




¹ Rates are per 100,000 and are adjusted to the 2000 U.S. population standard. Rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the rates for the U.S. combined.

² See <http://smm.sagepub.com/cgi/content/abstract/15/6/547> for the methodology used in the calculation of the age-adjusted 95% confidence intervals.

³ See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.

N.B. In areas with small Latino populations, methods to indirectly identify Latinos (like NHIA) can overestimate the ethnicity-specific counts of cancer cases. Also, even small errors in Latino population estimates can affect the magnitude of the cancer rates.

Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates¹ and 95% Confidence Intervals²
 Selected Areas in the United States³, 2000-2004, Hispanic/Latino, All Races, Males



¹ Rates are per 100,000 and are adjusted to the 2000 U.S. population standard. Rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the rates for the U.S. combined.


² See <http://smm.sagepub.com/cgi/content/abstract/15/6/547> for the methodology used in the calculation of the age-adjusted 95% confidence intervals.

³ See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.

N.B. In areas with small Latino populations, methods to indirectly identify Latinos (like NHIA) can overestimate the ethnicity-specific counts of cancer cases. Also, even small errors in Latino population estimates can affect the magnitude of the cancer rates.

**Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates¹ and 95% Confidence Intervals²
Selected Areas in the United States³, 2000-2004, Hispanic/Latina, All Races, Females**

Lung and Bronchus



¹ Rates are per 100,000 and are adjusted to the 2000 U.S. population standard. Rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the rates for the U.S. combined.


² See <http://smm.sagepub.com/cgi/content/abstract/15/6/547> for the methodology used in the calculation of the age-adjusted 95% confidence intervals.

³ See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.

N.B. In areas with small Latino populations, methods to indirectly identify Latinos (like NHIA) can overestimate the ethnicity-specific counts of cancer cases. Also, even small errors in Latino population estimates can affect the magnitude of the cancer rates.

Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates¹ and 95% Confidence Intervals²
 Selected Areas in the United States³, 2000-2004, Hispanic/Latino, All Races, Males

Lymphomas



¹ Rates are per 100,000 and are adjusted to the 2000 U.S. population standard. Rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the rates for the U.S. combined.


² See <http://smm.sagepub.com/cgi/content/abstract/15/6/547> for the methodology used in the calculation of the age-adjusted 95% confidence intervals.

³ See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.

N.B. In areas with small Latino populations, methods to indirectly identify Latinos (like NHIA) can overestimate the ethnicity-specific counts of cancer cases. Also, even small errors in Latino population estimates can affect the magnitude of the cancer rates.

**Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates¹ and 95% Confidence Intervals²
Selected Areas in the United States³, 2000-2004, Hispanic/Latina, All Races, Females**

Lymphomas



¹ Rates are per 100,000 and are adjusted to the 2000 U.S. population standard. Rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the rates for the U.S. combined.


² See <http://smm.sagepub.com/cgi/content/abstract/15/6/547> for the methodology used in the calculation of the age-adjusted 95% confidence intervals.

³ See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.

N.B. In areas with small Latino populations, methods to indirectly identify Latinos (like NHIA) can overestimate the ethnicity-specific counts of cancer cases. Also, even small errors in Latino population estimates can affect the magnitude of the cancer rates.

Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates¹ and 95% Confidence Intervals²
 Selected Areas in the United States³, 2000-2004, Hispanic/Latina, All Races, Males

Oral Cavity and Pharynx



¹ Rates are per 100,000 and are adjusted to the 2000 U.S. population standard. Rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the rates for the U.S. combined.


² See <http://smm.sagepub.com/cgi/content/abstract/15/6/547> for the methodology used in the calculation of the age-adjusted 95% confidence intervals.

³ See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.

N.B. In areas with small Latino populations, methods to indirectly identify Latinos (like NHIA) can overestimate the ethnicity-specific counts of cancer cases. Also, even small errors in Latino population estimates can affect the magnitude of the cancer rates.

**Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates¹ and 95% Confidence Intervals²
Selected Areas in the United States³, 2000-2004, Hispanic/Latina, All Races, Females**

Oral Cavity and Pharynx



¹ Rates are per 100,000 and are adjusted to the 2000 U.S. population standard. Rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the rates for the U.S. combined.


² See <http://smm.sagepub.com/cgi/content/abstract/15/6/547> for the methodology used in the calculation of the age-adjusted 95% confidence intervals.

³ See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.

N.B. In areas with small Latino populations, methods to indirectly identify Latinos (like NHIA) can overestimate the ethnicity-specific counts of cancer cases. Also, even small errors in Latino population estimates can affect the magnitude of the cancer rates.

This Page Intentionally Left Blank

**Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates¹ and 95% Confidence Intervals²
Selected Areas in the United States³, 2000-2004, Hispanic/Latina, All Races, Females**




¹ Rates are per 100,000 and are adjusted to the 2000 U.S. population standard. Rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the rates for the U.S. combined.

² See <http://smm.sagepub.com/cgi/content/abstract/15/6/547> for the methodology used in the calculation of the age-adjusted 95% confidence intervals.

³ See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.

N.B. In areas with small Latino populations, methods to indirectly identify Latinos (like NHIA) can overestimate the ethnicity-specific counts of cancer cases. Also, even small errors in Latino population estimates can affect the magnitude of the cancer rates.

Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates¹ and 95% Confidence Intervals²
 Selected Areas in the United States³, 2000-2004, Hispanic/Latino, All Races, Males




¹ Rates are per 100,000 and are adjusted to the 2000 U.S. population standard. Rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the rates for the U.S. combined.

² See <http://smm.sagepub.com/cgi/content/abstract/15/6/547> for the methodology used in the calculation of the age-adjusted 95% confidence intervals.

³ See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.

N.B. In areas with small Latino populations, methods to indirectly identify Latinos (like NHIA) can overestimate the ethnicity-specific counts of cancer cases. Also, even small errors in Latino population estimates can affect the magnitude of the cancer rates.

**Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates¹ and 95% Confidence Intervals²
Selected Areas in the United States³, 2000-2004, Hispanic/Latina, All Races, Females**




¹ Rates are per 100,000 and are adjusted to the 2000 U.S. population standard. Rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the rates for the U.S. combined.

² See <http://smm.sagepub.com/cgi/content/abstract/15/6/547> for the methodology used in the calculation of the age-adjusted 95% confidence intervals.

³ See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.

N.B. In areas with small Latino populations, methods to indirectly identify Latinos (like NHIA) can overestimate the ethnicity-specific counts of cancer cases. Also, even small errors in Latino population estimates can affect the magnitude of the cancer rates.

**Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates¹ and 95% Confidence Intervals²
Selected Areas in the United States³, 2000-2004, Hispanic/Latino, All Races, Males**



¹ Rates are per 100,000 and are adjusted to the 2000 U.S. population standard. Rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the rates for the U.S. combined.


² See <http://smm.sagepub.com/cgi/content/abstract/15/6/547> for the methodology used in the calculation of the age-adjusted 95% confidence intervals.

³ See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.

N.B. In areas with small Latino populations, methods to indirectly identify Latinos (like NHIA) can overestimate the ethnicity-specific counts of cancer cases. Also, even small errors in Latino population estimates can affect the magnitude of the cancer rates.

This Page Intentionally Left Blank

Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates¹ and 95% Confidence Intervals²
 Selected Areas in the United States³, 2000-2004, Hispanic/Latino, All Races, Males




¹ Rates are per 100,000 and are adjusted to the 2000 U.S. population standard. Rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the rates for the U.S. combined.

² See <http://smm.sagepub.com/cgi/content/abstract/15/6/547> for the methodology used in the calculation of the age-adjusted 95% confidence intervals.

³ See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.

N.B. In areas with small Latino populations, methods to indirectly identify Latinos (like NHIA) can overestimate the ethnicity-specific counts of cancer cases. Also, even small errors in Latino population estimates can affect the magnitude of the cancer rates.

**Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates¹ and 95% Confidence Intervals²
Selected Areas in the United States³, 2000-2004, Hispanic/Latina, All Races, Females**




¹ Rates are per 100,000 and are adjusted to the 2000 U.S. population standard. Rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the rates for the U.S. combined.

² See <http://smm.sagepub.com/cgi/content/abstract/15/6/547> for the methodology used in the calculation of the age-adjusted 95% confidence intervals.

³ See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.

N.B. In areas with small Latino populations, methods to indirectly identify Latinos (like NHIA) can overestimate the ethnicity-specific counts of cancer cases. Also, even small errors in Latino population estimates can affect the magnitude of the cancer rates.

Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates¹ and 95% Confidence Intervals²
 Selected Areas in the United States³, 2000-2004, Hispanic/Latino, All Races, Males



¹ Rates are per 100,000 and are adjusted to the 2000 U.S. population standard. Rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the rates for the U.S. combined.

² See <http://smm.sagepub.com/cgi/content/abstract/15/6/547> for the methodology used in the calculation of the age-adjusted 95% confidence intervals.


³ See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.

N.B. In areas with small Latino populations, methods to indirectly identify Latinos (like NHIA) can overestimate the ethnicity-specific counts of cancer cases. Also, even small errors in Latino population estimates can affect the magnitude of the cancer rates.

This Page Intentionally Left Blank

Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates¹ and 95% Confidence Intervals²
 Selected Areas in the United States³, 2000-2004, Hispanic/Latino, All Races, Males

Thyroid




¹ Rates are per 100,000 and are adjusted to the 2000 U.S. population standard. Rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the rates for the U.S. combined.

² See <http://smm.sagepub.com/cgi/content/abstract/15/6/547> for the methodology used in the calculation of the age-adjusted 95% confidence intervals.

³ See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.

N.B. In areas with small Latino populations, methods to indirectly identify Latinos (like NHIA) can overestimate the ethnicity-specific counts of cancer cases. Also, even small errors in Latino population estimates can affect the magnitude of the cancer rates.

**Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates¹ and 95% Confidence Intervals²
Selected Areas in the United States³, 2000-2004, Hispanic/Latina, All Races, Females**




¹ Rates are per 100,000 and are adjusted to the 2000 U.S. population standard. Rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the rates for the U.S. combined.

² See <http://smm.sagepub.com/cgi/content/abstract/15/6/547> for the methodology used in the calculation of the age-adjusted 95% confidence intervals.

³ See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.

N.B. In areas with small Latino populations, methods to indirectly identify Latinos (like NHIA) can overestimate the ethnicity-specific counts of cancer cases. Also, even small errors in Latino population estimates can affect the magnitude of the cancer rates.


Average Annual Age-adjusted (2000 U.S. Standard) U.S. Cancer Incidence Rates¹ (NAACCR Combined²) for Selected Cancer Sites by Race/Ethnicity, Males, 2000-2004



¹ Rates are per 100,000 population and are age-adjusted by five-year age groups to the 2000 U.S. standard population based on single years of age. Rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the rates for the U.S. combined.
² See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.

N.B. In areas with small Latino populations, methods to indirectly identify Latinos (like NHIA) can overestimate the ethnicity-specific counts of cancer cases. Also, even small errors in Latino population estimates can affect the magnitude of the cancer rates.

Average Annual Age-adjusted (2000 U.S. Standard) U.S. Cancer Incidence Rates¹ (NAACCR Combined²) for Selected Cancer Sites by Race/Ethnicity, Females, 2000-2004



¹ Rates are per 100,000 population and are age-adjusted by five-year age groups to the 2000 U.S. standard population based on single years of age. Rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the rates for the U.S. combined.

² See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.

N.B. In areas with small Latino populations, methods to indirectly identify Latinos (like NHIA) can overestimate the ethnicity-specific counts of cancer cases. Also, even small errors in Latino population estimates can affect the magnitude of the cancer rates.