Section IV

Average-annual Age-adjusted Pediatric Cancer Incidence for the NAACCR (U.S.) Combined by Sex and Hispanic/Latino Ethnicity



Registries Included in the NAACCR U.S. Combined Pediatric Cancer Incidence 1998-2002



Total Cancer Cases and Average Annual Age-specific Incidence Rates[1] for Pediatric Cancers
United States (NAACCR Combined[2]), 1998-2002, Hispanic/Latino, All Races, Males

	Ages 0-4			Ages 5-9			Ages 10-14			Ages 15-19		
Sites	*Cases*	*Rate*	*S.E.*[3]	*Cases*	*Rate*	*S.E.*	*Cases*	*Rate*	*S.E.*	*Cases*	*Rate*	*S.E.*
All Sites	**1542**	**213.3**	**5.43**	**842**	**119.7**	**4.12**	**766**	**123.1**	**4.45**	**1219**	**196.8**	**5.64**
Leukemias	630	87.1	3.47	331	47.0	2.59	222	35.7	2.39	270	43.6	2.65
Lymphoid leukemia	529	73.2	3.18	277	39.4	2.37	161	25.9	2.04	183	29.5	2.18
Acute non-lymphocytic leukemia	67	9.3	1.13	42	6.0	0.92	42	6.7	1.04	60	9.7	1.25
Lymphomas & Other Reticuloendothelial Neoplasms	88	12.2	1.30	152	21.6	1.75	153	24.6	1.99	265	42.8	2.63
Hodgkin disease	19	2.6	0.60	60	8.5	1.10	66	10.6	1.31	146	23.6	1.95
Non-Hodgkin lymphoma	38	5.3	0.85	54	7.7	1.04	66	10.6	1.31	90	14.5	1.53
CNS[3] & Misc. Intracranial & Intraspinal Neoplasms	241	33.3	2.15	185	26.3	1.93	156	25.1	2.01	112	18.1	1.71
Astrocytoma	92	12.7	1.33	89	12.6	1.34	88	14.1	1.51	47	7.6	1.11
Primitive neuroectodermal tumors	73	10.1	1.18	44	6.3	0.94	31	5.0	0.89	36	5.8	0.97
Sympathetic Nervous System Tumors	147	20.3	1.68	12	1.7	0.49	-	-	-	7	1.1	0.43
Neuroblastoma & ganglioneuroblastoma	144	19.9	1.66	10	1.4	0.45	-	-	-	-	-	-
Retinoblastoma	108	14.9	1.44	-	-	-	-	-	-	-	-	-
Renal Tumors	106	14.7	1.42	29	4.1	0.77	-	-	-	-	-	-
Wilms' tumor	105	14.5	1.42	28	4.0	0.75	-	-	-	-	-	-
Hepatic Tumors	38	5.3	0.85	-	-	-	-	-	-	7	1.1	0.43
Malignant Bone Tumors	9	1.2	0.42	41	5.8	0.91	74	11.9	1.38	103	16.6	1.64
Osteosarcoma	-	-	-	28	4.0	0.75	47	7.6	1.10	63	10.2	1.28
Ewing’s sarcoma	6	0.8	0.34	9	1.3	0.43	22	3.5	0.75	27	4.4	0.84
Soft Tissue Sarcomas	86	11.9	1.28	51	7.2	1.01	66	10.6	1.31	94	15.2	1.57
Rhabdomyosarcoma & embryonal sarcoma	49	6.8	0.97	26	3.7	0.72	16	2.6	0.64	14	2.3	0.60
Germ Cell, Trophoblastic & Other Gonadal Neoplasms	76	10.5	1.21	10	1.4	0.45	50	8.0	1.14	259	41.8	2.60
Carcinomas and Other Malignant Epithelial Neoplasms	8	1.1	0.39	20	2.8	0.64	34	5.5	0.94	94	15.2	1.57
Thyroid carcinoma	-	-	-	7	1.0	0.38	10	1.6	0.51	22	3.6	0.76
Other and Unspecified Epithelial Neoplasms	-	-	-	-	-	-	10	1.6	0.51	43	6.9	1.06
Other and Unspecified Malignant Tumors	-	-	-	-	-	-	-	-	-	-	-	-

[1] Rates are **per million** population and were age-adjusted by five-year age groups to the 2000 U.S. standard population based on single years of age and to the World population standard. Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for All Sites combined.
[2] See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.
[3] Abbreviations: S.E. - Standard Error; CNS - Central Nervous System.

N.B. In areas with small Latino populations, methods to indirectly identify Latinos (like NHIA) can overestimate the ethnicity-specific counts of cancer cases. Also, even small errors in Latino population estimates can affect the magnitude of the cancer rates.

Total Cancer Cases and Average Annual Age-specific Incidence Rates[1] for Pediatric Cancers
United States (NAACCR Combined[2]), 1998-2002, Hispanic/Latina, All Races, Females

	Ages 0-4			Ages 5-9			Ages 10-14			Ages 15-19		
Sites	*Cases*	*Rate*	*S.E.*[3]	*Cases*	*Rate*	*S.E.*	*Cases*	*Rate*	*S.E.*	*Cases*	*Rate*	*S.E.*
All Sites	**1287**	**186.2**	**5.19**	**719**	**106.8**	**3.98**	**725**	**121.8**	**4.52**	**913**	**164.5**	**5.44**
Leukemias	526	76.1	3.32	311	46.2	2.62	195	32.8	2.35	145	26.1	2.17
Lymphoid leukemia	412	59.6	2.94	265	39.4	2.42	137	23.0	1.97	76	13.7	1.57
Acute non-lymphocytic leukemia	85	12.3	1.33	34	5.1	0.87	48	8.1	1.16	44	7.9	1.20
Lymphomas & Other Reticuloendothelial Neoplasms	43	6.2	0.95	39	5.8	0.93	100	16.8	1.68	200	36.0	2.55
Hodgkin disease	9	1.3	0.43	11	1.6	0.49	58	9.7	1.28	134	24.1	2.09
Non-Hodgkin lymphoma	17	2.5	0.60	20	3.0	0.66	32	5.4	0.95	49	8.8	1.26
CNS[3] & Misc. Intracranial & Intraspinal Neoplasms	221	32.0	2.15	179	26.6	1.99	100	16.8	1.68	83	15.0	1.64
Astrocytoma	78	11.3	1.28	78	11.6	1.31	53	8.9	1.22	47	8.5	1.24
Primitive neuroectodermal tumors	78	11.3	1.28	40	5.9	0.94	20	3.4	0.75	14	2.5	0.67
Sympathetic Nervous System Tumors	134	19.4	1.67	15	2.2	0.58	-	-	-	-	-	-
Neuroblastoma & ganglioneuroblastoma	132	19.1	1.66	15	2.2	0.58	-	-	-	-	-	-
Retinoblastoma	96	13.9	1.42	-	-	-	-	-	-	-	-	-
Renal Tumors	110	15.9	1.52	26	3.9	0.76	-	-	-	-	-	-
Wilms' tumor	110	15.9	1.52	26	3.9	0.76	-	-	-	-	-	-
Hepatic Tumors	31	4.5	0.81	-	-	-	-	-	-	-	-	-
Malignant Bone Tumors	7	1.0	0.38	36	5.3	0.89	95	16.0	1.64	67	12.1	1.48
Osteosarcoma	-	-	-	25	3.7	0.74	62	10.4	1.32	37	6.7	1.10
Ewing's sarcoma	6	0.9	0.35	7	1.0	0.39	25	4.2	0.84	20	3.6	0.81
Soft Tissue Sarcomas	66	9.5	1.18	52	7.7	1.07	70	11.8	1.41	72	13.0	1.53
Rhabdomyosarcoma & embryonal sarcoma	37	5.4	0.88	21	3.1	0.68	24	4.0	0.82	13	2.3	0.65
Germ Cell, Trophoblastic & Other Gonadal Neoplasms	39	5.6	0.90	32	4.8	0.84	67	11.3	1.38	116	20.9	1.94
Carcinomas and Other Malignant Epithelial Neoplasms	9	1.3	0.43	19	2.8	0.65	82	13.8	1.52	211	38.0	2.62
Thyroid carcinoma	-	-	-	7	1.0	0.39	42	7.1	1.09	127	22.9	2.03
Other and Unspecified Epithelial Neoplasms	-	-	-	6	0.9	0.36	25	4.2	0.84	62	11.2	1.42
Other and Unspecified Malignant Tumors	-	-	-	-	-	-	6	1.0	0.41	6	1.1	0.44

[1] Rates are **per million** population and were age-adjusted by five-year age groups to the 2000 U.S. standard population based on single years of age and to the World population standard. Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for All Sites combined.
[2] See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.
[3] Abbreviations: S.E. - Standard Error; CNS - Central Nervous System.

N.B. In areas with small Latino populations, methods to indirectly identify Latinos (like NHIA) can overestimate the ethnicity-specific counts of cancer cases. Also, even small errors in Latino population estimates can affect the magnitude of the cancer rates.

Total Cancer Cases and Average Annual Age-specific Incidence Rates[1] for Pediatric Cancers
United States (NAACCR Combined[2]), 1998-2002, Non-Hispanic White, Males

	Ages 0-4			Ages 5-9			Ages 10-14			Ages 15-19		
Sites	*Cases*	*Rate*	*S.E.*[3]	*Cases*	*Rate*	*S.E.*	*Cases*	*Rate*	*S.E.*	*Cases*	*Rate*	*S.E.*
All Sites	**5487**	**238.4**	**3.22**	**3166**	**126.5**	**2.25**	**3429**	**130.6**	**2.23**	**5968**	**230.0**	**2.98**
Leukemias	2019	87.7	1.95	1030	41.1	1.28	700	26.7	1.01	813	31.3	1.10
Lymphoid leukemia	1609	69.9	1.74	882	35.2	1.19	500	19.0	0.85	507	19.5	0.87
Acute non-lymphocytic leukemia	308	13.4	0.76	103	4.1	0.41	157	6.0	0.48	228	8.8	0.58
Lymphomas & Other Reticuloendothelial Neoplasms	211	9.2	0.63	483	19.3	0.88	774	29.5	1.06	1487	57.3	1.49
Hodgkin disease	26	1.1	0.22	116	4.6	0.43	349	13.3	0.71	913	35.2	1.16
Non-Hodgkin lymphoma	110	4.8	0.46	210	8.4	0.58	273	10.4	0.63	443	17.1	0.81
CNS[3] & Misc. Intracranial & Intraspinal Neoplasms	980	42.6	1.36	923	36.9	1.21	765	29.1	1.05	606	23.4	0.95
Astrocytoma	446	19.4	0.92	396	15.8	0.80	419	16.0	0.78	340	13.1	0.71
Primitive neuroectodermal tumors	267	11.6	0.71	272	10.9	0.66	161	6.1	0.48	90	3.5	0.37
Sympathetic Nervous System Tumors	787	34.2	1.22	95	3.8	0.39	34	1.3	0.22	20	0.8	0.17
Neuroblastoma & ganglioneuroblastoma	782	34.0	1.21	90	3.6	0.38	27	1.0	0.20	10	0.4	0.12
Retinoblastoma	261	11.3	0.70	8	0.3	0.11	-	-	-	-	-	-
Renal Tumors	460	20.0	0.93	105	4.2	0.41	28	1.1	0.20	28	1.1	0.20
Wilms' tumor	453	19.7	0.92	99	4.0	0.40	15	0.6	0.15	7	0.3	0.10
Hepatic Tumors	141	6.1	0.52	14	0.6	0.15	17	0.6	0.16	32	1.2	0.22
Malignant Bone Tumors	34	1.5	0.25	152	6.1	0.49	383	14.6	0.75	544	21.0	0.90
Osteosarcoma	8	0.3	0.12	62	2.5	0.31	199	7.6	0.54	288	11.1	0.65
Ewing's sarcoma	20	0.9	0.19	83	3.3	0.36	159	6.1	0.48	197	7.6	0.54
Soft Tissue Sarcomas	355	15.4	0.82	238	9.5	0.62	300	11.4	0.66	435	16.8	0.80
Rhabdomyosarcoma & embryonal sarcoma	224	9.7	0.65	140	5.6	0.47	96	3.7	0.37	134	5.2	0.45
Germ Cell, Trophoblastic & Other Gonadal Neoplasms	146	6.3	0.52	32	1.3	0.23	130	5.0	0.43	1087	41.9	1.27
Carcinomas and Other Malignant Epithelial Neoplasms	70	3.0	0.36	78	3.1	0.35	286	10.9	0.64	891	34.3	1.15
Thyroid carcinoma	-	-	-	26	1.0	0.20	69	2.6	0.32	171	6.6	0.50
Other and Unspecified Epithelial Neoplasms	29	1.3	0.23	16	0.6	0.16	84	3.2	0.35	253	9.8	0.61
Other and Unspecified Malignant Tumors	23	1.0	0.21	8	0.3	0.11	12	0.5	0.13	25	1.0	0.19

[1] Rates are **per million** population and were age-adjusted by five-year age groups to the 2000 U.S. standard population based on single years of age and to the World population standard. Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for All Sites combined.
[2] See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.
[3] Abbreviations: S.E. - Standard Error; CNS - Central Nervous System.

N.B. In areas with small Latino populations, methods to indirectly identify Latinos (like NHIA) can overestimate the ethnicity-specific counts of cancer cases. Also, even small errors in Latino population estimates can affect the magnitude of the cancer rates.

Total Cancer Cases and Average Annual Age-specific Incidence Rates[1] for Pediatric Cancers
United States (NAACCR Combined[2]), 1998-2002, Non-Hispanic White, Females

	Ages 0-4			Ages 5-9			Ages 10-14			Ages 15-19		
Sites	*Cases*	*Rate*	*S.E.*[3]	*Cases*	*Rate*	*S.E.*	*Cases*	*Rate*	*S.E.*	*Cases*	*Rate*	*S.E.*
All Sites	**4508**	**206.1**	**3.07**	**2473**	**104.3**	**2.10**	**3031**	**122.0**	**2.22**	**5202**	**210.8**	**2.92**
Leukemias	1623	74.2	1.84	784	33.1	1.18	591	23.8	0.98	534	21.6	0.94
Lymphoid leukemia	1319	60.3	1.66	644	27.2	1.07	399	16.1	0.80	256	10.4	0.65
Acute non-lymphocytic leukemia	227	10.4	0.69	103	4.3	0.43	152	6.1	0.50	225	9.1	0.61
Lymphomas & Other Reticuloendothelial Neoplasms	128	5.9	0.52	190	8.0	0.58	531	21.4	0.93	1294	52.4	1.46
Hodgkin disease	-	-	-	46	1.9	0.29	331	13.3	0.73	993	40.2	1.28
Non-Hodgkin lymphoma	79	3.6	0.41	96	4.0	0.41	144	5.8	0.48	260	10.5	0.65
CNS[3] & Misc. Intracranial & Intraspinal Neoplasms	816	37.3	1.31	757	31.9	1.16	637	25.6	1.02	476	19.3	0.88
Astrocytoma	387	17.7	0.90	399	16.8	0.84	359	14.5	0.76	295	12.0	0.70
Primitive neuroectodermal tumors	183	8.4	0.62	147	6.2	0.51	103	4.1	0.41	48	1.9	0.28
Sympathetic Nervous System Tumors	687	31.4	1.20	77	3.2	0.37	32	1.3	0.23	16	0.6	0.16
Neuroblastoma & ganglioneuroblastoma	681	31.1	1.19	71	3.0	0.36	24	1.0	0.20	-	-	-
Retinoblastoma	222	10.2	0.68	10	0.4	0.13	-	-	-	-	-	-
Renal Tumors	438	20.0	0.96	157	6.6	0.53	32	1.3	0.23	26	1.1	0.21
Wilms' tumor	433	19.8	0.95	151	6.4	0.52	21	0.8	0.18	-	-	-
Hepatic Tumors	104	4.8	0.47	-	-	-	18	0.7	0.17	23	0.9	0.19
Malignant Bone Tumors	29	1.3	0.25	135	5.7	0.49	331	13.3	0.73	302	12.2	0.70
Osteosarcoma	-	-	-	53	2.2	0.31	181	7.3	0.54	154	6.2	0.50
Ewing's sarcoma	23	1.1	0.22	72	3.0	0.36	133	5.4	0.46	107	4.3	0.42
Soft Tissue Sarcomas	248	11.3	0.72	173	7.3	0.55	271	10.9	0.66	371	15.0	0.78
Rhabdomyosarcoma & embryonal sarcoma	148	6.8	0.56	86	3.6	0.39	74	3.0	0.35	65	2.6	0.33
Germ Cell, Trophoblastic & Other Gonadal Neoplasms	132	6.0	0.53	72	3.0	0.36	156	6.3	0.50	331	13.4	0.74
Carcinomas and Other Malignant Epithelial Neoplasms	54	2.5	0.34	108	4.6	0.44	414	16.7	0.82	1788	72.5	1.71
Thyroid carcinoma	-	-	-	38	1.6	0.26	181	7.3	0.54	806	32.7	1.15
Other and Unspecified Epithelial Neoplasms	13	0.6	0.16	25	1.1	0.21	96	3.9	0.39	315	12.8	0.72
Other and Unspecified Malignant Tumors	27	1.2	0.24	7	0.3	0.11	18	0.7	0.17	41	1.7	0.26

[1] Rates are **per million** population and were age-adjusted by five-year age groups to the 2000 U.S. standard population based on single years of age and to the World population standard. Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for All Sites combined.
[2] See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.
[3] Abbreviations: S.E. - Standard Error; CNS - Central Nervous System.

N.B. In areas with small Latino populations, methods to indirectly identify Latinos (like NHIA) can overestimate the ethnicity-specific counts of cancer cases. Also, even small errors in Latino population estimates can affect the magnitude of the cancer rates.

Total Cancer Cases and Average Annual Age-specific Incidence Rates[1] for Pediatric Cancers
United States (NAACCR Combined[2]), 1998-2002, Non-Hispanic Black, Males

	Ages 0-4			Ages 5-9			Ages 10-14			Ages 15-19		
Sites	*Cases*	*Rate*	*S.E.*[3]	*Cases*	*Rate*	*S.E.*	*Cases*	*Rate*	*S.E.*	*Cases*	*Rate*	*S.E.*
All Sites	**761**	**146.7**	**5.32**	**513**	**89.4**	**3.95**	**569**	**101.8**	**4.27**	**777**	**150.8**	**5.41**
Leukemias	183	35.3	2.61	120	20.9	1.91	136	24.3	2.09	129	25.0	2.20
Lymphoid leukemia	133	25.6	2.22	96	16.7	1.71	86	15.4	1.66	64	12.4	1.55
Acute non-lymphocytic leukemia	42	8.1	1.25	17	3.0	0.72	39	7.0	1.12	46	8.9	1.32
Lymphomas & Other Reticuloendothelial Neoplasms	16	3.1	0.77	86	15.0	1.62	148	26.5	2.18	202	39.2	2.76
Hodgkin disease	-	-	-	23	4.0	0.84	66	11.8	1.45	103	20.0	1.97
Non-Hodgkin lymphoma	8	1.5	0.55	42	7.3	1.13	68	12.2	1.48	87	16.9	1.81
CNS[3] & Misc. Intracranial & Intraspinal Neoplasms	158	30.5	2.42	167	29.1	2.25	85	15.2	1.65	93	18.0	1.87
Astrocytoma	55	10.6	1.43	81	14.1	1.57	42	7.5	1.16	56	10.9	1.45
Primitive neuroectodermal tumors	44	8.5	1.28	35	6.1	1.03	22	3.9	0.84	16	3.1	0.78
Sympathetic Nervous System Tumors	115	22.2	2.07	23	4.0	0.84	6	1.1	0.44	-	-	-
Neuroblastoma & ganglioneuroblastoma	114	22.0	2.06	22	3.8	0.82	6	1.1	0.44	-	-	-
Retinoblastoma	64	12.3	1.54	-	-	-	-	-	-	-	-	-
Renal Tumors	98	18.9	1.91	27	4.7	0.91	11	2.0	0.59	15	2.9	0.75
Wilms' tumor	97	18.7	1.90	25	4.4	0.87	-	-	-	-	-	-
Hepatic Tumors	21	4.0	0.88	-	-	-	6	1.1	0.44	7	1.4	0.51
Malignant Bone Tumors	6	1.2	0.47	21	3.7	0.80	51	9.1	1.28	94	18.2	1.88
Osteosarcoma	-	-	-	18	3.1	0.74	38	6.8	1.10	73	14.2	1.66
Ewing's sarcoma	-	-	-	-	-	-	10	1.8	0.57	10	1.9	0.61
Soft Tissue Sarcomas	72	13.9	1.64	50	8.7	1.23	65	11.6	1.44	98	19.0	1.92
Rhabdomyosarcoma & embryonal sarcoma	43	8.3	1.26	32	5.6	0.99	22	3.9	0.84	34	6.6	1.13
Germ Cell, Trophoblastic & Other Gonadal Neoplasms	20	3.9	0.86	-	-	-	8	1.4	0.51	49	9.5	1.36
Carcinomas and Other Malignant Epithelial Neoplasms	6	1.2	0.47	10	1.7	0.55	48	8.6	1.24	81	15.7	1.75
Thyroid carcinoma	-	-	-	-	-	-	-	-	-	8	1.6	0.55
Other and Unspecified Epithelial Neoplasms	-	-	-	-	-	-	17	3.0	0.74	43	8.3	1.27
Other and Unspecified Malignant Tumors	-	-	-	-	-	-	-	-	-	-	-	-

[1] Rates are **per million** population and were age-adjusted by five-year age groups to the 2000 U.S. standard population based on single years of age and to the World population standard. Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for All Sites combined.
[2] See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.
[3] Abbreviations: S.E. - Standard Error; CNS - Central Nervous System.

N.B. In areas with small Latino populations, methods to indirectly identify Latinos (like NHIA) can overestimate the ethnicity-specific counts of cancer cases. Also, even small errors in Latino population estimates can affect the magnitude of the cancer rates.

Total Cancer Cases and Average Annual Age-specific Incidence Rates[1] for Pediatric Cancers
United States (NAACCR Combined[2]), 1998-2002, Non-Hispanic Black, Females

	Ages 0-4			Ages 5-9			Ages 10-14			Ages 15-19		
Sites	*Cases*	*Rate*	*S.E.*[3]	*Cases*	*Rate*	*S.E.*	*Cases*	*Rate*	*S.E.*	*Cases*	*Rate*	*S.E.*
All Sites	**700**	**139.5**	**5.27**	**422**	**76.1**	**3.70**	**539**	**99.5**	**4.28**	**662**	**131.9**	**5.12**
Leukemias	181	36.1	2.68	120	21.6	1.97	102	18.8	1.86	99	19.7	1.98
Lymphoid leukemia	133	26.5	2.30	91	16.4	1.72	57	10.5	1.39	40	8.0	1.26
Acute non-lymphocytic leukemia	39	7.8	1.24	23	4.1	0.86	35	6.5	1.09	39	7.8	1.24
Lymphomas & Other Reticuloendothelial Neoplasms	17	3.4	0.82	36	6.5	1.08	86	15.9	1.71	149	29.7	2.43
Hodgkin disease	-	-	-	10	1.8	0.57	42	7.7	1.20	89	17.7	1.88
Non-Hodgkin lymphoma	12	2.4	0.69	25	4.5	0.90	38	7.0	1.14	55	11.0	1.48
CNS[3] & Misc. Intracranial & Intraspinal Neoplasms	126	25.1	2.24	107	19.3	1.86	101	18.6	1.85	60	12.0	1.54
Astrocytoma	52	10.4	1.44	48	8.7	1.25	58	10.7	1.41	32	6.4	1.13
Primitive neuroectodermal tumors	38	7.6	1.23	22	4.0	0.85	15	2.8	0.71	12	2.4	0.69
Sympathetic Nervous System Tumors	88	17.5	1.87	16	2.9	0.72	7	1.3	0.49	-	-	-
Neuroblastoma & ganglioneuroblastoma	87	17.3	1.86	14	2.5	0.67	6	1.1	0.45	-	-	-
Retinoblastoma	57	11.4	1.50	-	-	-	-	-	-	-	-	-
Renal Tumors	119	23.7	2.17	50	9.0	1.27	13	2.4	0.67	11	2.2	0.66
Wilms' tumor	117	23.3	2.16	46	8.3	1.22	7	1.3	0.49	-	-	-
Hepatic Tumors	12	2.4	0.69	-	-	-	-	-	-	-	-	-
Malignant Bone Tumors	-	-	-	22	4.0	0.85	63	11.6	1.46	35	7.0	1.18
Osteosarcoma	-	-	-	18	3.2	0.76	57	10.5	1.39	20	4.0	0.89
Ewing's sarcoma	-	-	-	-	-	-	-	-	-	-	-	-
Soft Tissue Sarcomas	54	10.8	1.46	40	7.2	1.14	67	12.4	1.51	73	14.5	1.70
Rhabdomyosarcoma & embryonal sarcoma	29	5.8	1.07	16	2.9	0.72	16	3.0	0.74	16	3.2	0.80
Germ Cell, Trophoblastic & Other Gonadal Neoplasms	35	7.0	1.18	21	3.8	0.83	40	7.4	1.17	84	16.7	1.83
Carcinomas and Other Malignant Epithelial Neoplasms	7	1.4	0.53	6	1.1	0.44	51	9.4	1.32	128	25.5	2.25
Thyroid carcinoma	-	-	-	-	-	-	20	3.7	0.83	45	9.0	1.34
Other and Unspecified Epithelial Neoplasms	-	-	-	-	-	-	15	2.8	0.71	61	12.1	1.56
Other and Unspecified Malignant Tumors	-	-	-	-	-	-	7	1.3	0.49	14	2.8	0.75

[1] Rates are **per million** population and were age-adjusted by five-year age groups to the 2000 U.S. standard population based on single years of age and to the World population standard. Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for All Sites combined.
[2] See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.
[3] Abbreviations: S.E. - Standard Error; CNS - Central Nervous System.

N.B. In areas with small Latino populations, methods to indirectly identify Latinos (like NHIA) can overestimate the ethnicity-specific counts of cancer cases. Also, even small errors in Latino population estimates can affect the magnitude of the cancer rates.

Total Cancer Cases and Average Annual Age-adjusted Incidence Rates[1] for Pediatric Cancers
United States (NAACCR Combined[2]), 1998-2002, Hispanic/Latino, All Races, Males

	Ages 0-14					Ages 0-19				
		2000 U.S.		World			2000 U.S.		World	
Sites	*Cases*	*Rate*	*S.E.*[3]	*Rate*	*S.E.*	*Cases*	*Rate*	*S.E.*	*Rate*	*S.E.*
All Sites	**3150**	**151.0**	**2.70**	**156.9**	**2.80**	**4369**	**162.5**	**2.47**	**165.9**	**2.52**
Leukemias	1183	56.3	1.64	59.5	1.73	1453	53.1	1.40	55.9	1.47
Lymphoid leukemia	967	45.9	1.48	48.8	1.57	1150	41.8	1.24	44.4	1.31
Acute non-lymphocytic leukemia	151	7.3	0.60	7.5	0.61	211	7.9	0.55	8.0	0.55
Lymphomas & Other Reticuloendothelial Neoplasms	393	19.6	0.99	18.9	0.95	658	25.4	0.99	24.2	0.95
Hodgkin disease	145	7.3	0.61	6.9	0.57	291	11.4	0.67	10.6	0.62
Non-Hodgkin lymphoma	158	7.9	0.63	7.6	0.61	248	9.6	0.61	9.2	0.58
CNS[3] & Misc. Intracranial & Intraspinal Neoplasms	582	28.2	1.17	28.7	1.19	694	25.6	0.98	26.3	1.00
Astrocytoma	269	13.2	0.81	13.1	0.80	316	11.8	0.67	11.9	0.67
Primitive neuroectodermal tumors	148	7.1	0.58	7.4	0.61	184	6.7	0.50	7.0	0.52
Sympathetic Nervous System Tumors	163	7.2	0.57	8.5	0.67	170	5.7	0.44	6.8	0.52
Neuroblastoma & ganglioneuroblastoma	155	6.9	0.55	8.1	0.65	157	5.2	0.42	6.3	0.51
Retinoblastoma	112	4.9	0.47	5.9	0.56	112	3.7	0.35	4.6	0.43
Renal Tumors	139	6.3	0.54	7.2	0.61	142	4.9	0.41	5.7	0.48
Wilms' tumor	135	6.1	0.53	7.0	0.60	136	4.6	0.40	5.5	0.47
Hepatic Tumors	44	2.0	0.30	2.3	0.35	51	1.8	0.25	2.0	0.29
Malignant Bone Tumors	124	6.4	0.58	5.8	0.52	227	9.0	0.60	8.3	0.55
Osteosarcoma	76	4.0	0.46	3.5	0.41	139	5.5	0.47	5.0	0.43
Ewing's sarcoma	37	1.9	0.31	1.8	0.29	64	2.5	0.32	2.3	0.29
Soft Tissue Sarcomas	203	9.9	0.70	10.0	0.70	297	11.2	0.65	11.2	0.65
Rhabdomyosarcoma & embryonal sarcoma	91	4.3	0.45	4.6	0.48	105	3.8	0.37	4.0	0.40
Germ Cell, Trophoblastic & Other Gonadal Neoplasms	136	6.5	0.56	6.8	0.58	395	15.4	0.78	14.7	0.74
Carcinomas and Other Malignant Epithelial Neoplasms	62	3.2	0.40	2.9	0.37	156	6.2	0.50	5.7	0.46
Thyroid carcinoma	17	0.9	0.21	0.8	0.19	39	1.6	0.25	1.4	0.23
Other and Unspecified Epithelial Neoplasms	15	0.8	0.20	0.7	0.18	58	2.3	0.31	2.1	0.28
Other and Unspecified Malignant Tumors	9	0.4	0.14	0.4	0.15	14	0.5	0.14	0.5	0.14

[1] Rates are **per million** population and were age-adjusted by five-year age groups to the 2000 U.S. standard population based on single years of age and to the World population standard. Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for All Sites combined.
[2] See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.
[3] Abbreviations: S.E. - Standard Error; CNS - Central Nervous System.

N.B. In areas with small Latino populations, methods to indirectly identify Latinos (like NHIA) can overestimate the ethnicity-specific counts of cancer cases. Also, even small errors in Latino population estimates can affect the magnitude of the cancer rates.

Total Cancer Cases and Average Annual Age-adjusted Incidence Rates[1] for Pediatric Cancers
United States (NAACCR Combined[2]), 1998-2002, Hispanic/Latina, All Races, Females

	Ages 0-14					Ages 0-19				
		2000 U.S.		World			2000 U.S.		World	
Sites	*Cases*	*Rate*	*S.E.*[3]	*Rate*	*S.E.*	*Cases*	*Rate*	*S.E.*	*Rate*	*S.E.*
All Sites	**2731**	**137.4**	**2.64**	**141.8**	**2.72**	**3644**	**144.2**	**2.40**	**146.9**	**2.44**
Leukemias	1032	51.4	1.60	54.0	1.69	1177	45.0	1.32	47.7	1.39
Lymphoid leukemia	814	40.5	1.42	42.6	1.50	890	33.7	1.14	36.1	1.21
Acute non-lymphocytic leukemia	167	8.4	0.65	8.7	0.67	211	8.3	0.57	8.5	0.59
Lymphomas & Other Reticuloendothelial Neoplasms	182	9.7	0.72	9.2	0.68	382	16.3	0.84	15.2	0.78
Hodgkin disease	78	4.3	0.49	3.9	0.44	212	9.3	0.64	8.4	0.58
Non-Hodgkin lymphoma	69	3.6	0.44	3.5	0.42	118	4.9	0.46	4.7	0.43
CNS[3] & Misc. Intracranial & Intraspinal Neoplasms	500	25.0	1.12	25.8	1.16	583	22.5	0.94	23.4	0.97
Astrocytoma	209	10.6	0.73	10.7	0.74	256	10.0	0.63	10.2	0.64
Primitive neuroectodermal tumors	138	6.8	0.58	7.3	0.62	152	5.7	0.47	6.2	0.50
Sympathetic Nervous System Tumors	152	7.1	0.58	8.3	0.67	154	5.4	0.44	6.5	0.53
Neuroblastoma & ganglioneuroblastoma	149	7.0	0.57	8.1	0.67	150	5.3	0.43	6.3	0.52
Retinoblastoma	100	4.6	0.46	5.5	0.55	101	3.5	0.35	4.3	0.43
Renal Tumors	140	6.7	0.56	7.6	0.64	145	5.2	0.43	6.1	0.51
Wilms' tumor	139	6.6	0.56	7.6	0.64	139	4.9	0.42	5.9	0.50
Hepatic Tumors	37	1.7	0.29	2.0	0.33	42	1.5	0.24	1.7	0.27
Malignant Bone Tumors	138	7.6	0.65	6.8	0.58	205	8.7	0.61	8.0	0.56
Osteosarcoma	88	4.8	0.52	4.3	0.46	125	5.3	0.48	4.8	0.43
Ewing's sarcoma	38	2.1	0.34	1.9	0.31	58	2.4	0.32	2.3	0.30
Soft Tissue Sarcomas	188	9.7	0.71	9.6	0.70	260	10.5	0.66	10.3	0.64
Rhabdomyosarcoma & embryonal sarcoma	82	4.2	0.46	4.3	0.47	95	3.7	0.38	3.8	0.39
Germ Cell, Trophoblastic & Other Gonadal Neoplasms	138	7.2	0.62	6.9	0.59	254	10.7	0.67	10.1	0.63
Carcinomas and Other Malignant Epithelial Neoplasms	110	6.1	0.58	5.4	0.52	321	14.1	0.79	12.8	0.71
Thyroid carcinoma	50	2.8	0.40	2.4	0.35	177	7.9	0.59	7.0	0.53
Other and Unspecified Epithelial Neoplasms	33	1.8	0.32	1.6	0.28	95	4.2	0.43	3.8	0.39
Other and Unspecified Malignant Tumors	14	0.7	0.19	0.7	0.19	20	0.8	0.18	0.8	0.18

[1] Rates are **per million** population and were age-adjusted by five-year age groups to the 2000 U.S. standard population based on single years of age and to the World population standard. Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for All Sites combined.
[2] See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.
[3] Abbreviations: S.E. - Standard Error; CNS - Central Nervous System.

N.B. In areas with small Latino populations, methods to indirectly identify Latinos (like NHIA) can overestimate the ethnicity-specific counts of cancer cases. Also, even small errors in Latino population estimates can affect the magnitude of the cancer rates.

Total Cancer Cases and Average Annual Age-adjusted Incidence Rates[1] for Pediatric Cancers
United States (NAACCR Combined[2]), 1998-2002, Non-Hispanic White, Males

	Ages 0-14					Ages 0-19				
		2000 U.S.		World			2000 U.S.		World	
Sites	*Cases*	*Rate*	*S.E.*[3]	*Rate*	*S.E.*	*Cases*	*Rate*	*S.E.*	*Rate*	*S.E.*
All Sites	**12082**	**163.9**	**1.49**	**171.0**	**1.58**	**18050**	**180.6**	**1.34**	**184.3**	**1.40**
Leukemias	3749	51.2	0.84	54.9	0.91	4562	46.2	0.68	49.6	0.75
Lymphoid leukemia	2991	40.8	0.75	43.9	0.81	3498	35.5	0.60	38.4	0.66
Acute non-lymphocytic leukemia	568	7.7	0.32	8.3	0.35	796	8.0	0.28	8.4	0.30
Lymphomas & Other Reticuloendothelial Neoplasms	1468	19.5	0.51	18.3	0.48	2955	29.0	0.53	27.1	0.50
Hodgkin disease	491	6.5	0.29	5.8	0.26	1404	13.7	0.37	12.4	0.33
Non-Hodgkin lymphoma	593	7.9	0.32	7.6	0.32	1036	10.2	0.32	9.7	0.30
CNS[3] & Misc. Intracranial & Intraspinal Neoplasms	2668	36.1	0.70	36.8	0.72	3274	32.9	0.57	33.8	0.60
Astrocytoma	1261	17.0	0.48	17.2	0.49	1601	16.0	0.40	16.3	0.41
Primitive neuroectodermal tumors	700	9.5	0.36	9.8	0.37	790	8.0	0.28	8.4	0.30
Sympathetic Nervous System Tumors	916	12.8	0.42	14.9	0.49	936	9.7	0.32	11.7	0.38
Neuroblastoma & ganglioneuroblastoma	899	12.5	0.42	14.7	0.49	909	9.5	0.31	11.4	0.38
Retinoblastoma	269	3.8	0.23	4.5	0.28	269	2.8	0.17	3.5	0.21
Renal Tumors	593	8.2	0.34	9.4	0.39	621	6.4	0.26	7.5	0.30
Wilms' tumor	567	7.9	0.33	9.1	0.38	574	6.0	0.25	7.1	0.30
Hepatic Tumors	172	2.4	0.18	2.7	0.21	204	2.1	0.15	2.4	0.17
Malignant Bone Tumors	569	7.5	0.31	6.8	0.29	1113	10.9	0.33	10.0	0.30
Osteosarcoma	269	3.5	0.22	3.1	0.19	557	5.4	0.23	4.9	0.21
Ewing's sarcoma	262	3.5	0.21	3.2	0.20	459	4.5	0.21	4.2	0.20
Soft Tissue Sarcomas	893	12.1	0.40	12.4	0.42	1328	13.3	0.36	13.4	0.37
Rhabdomyosarcoma & embryonal sarcoma	460	6.3	0.29	6.6	0.31	594	6.0	0.25	6.3	0.26
Germ Cell, Trophoblastic & Other Gonadal Neoplasms	308	4.2	0.24	4.3	0.25	1395	13.7	0.37	12.8	0.35
Carcinomas and Other Malignant Epithelial Neoplasms	434	5.7	0.28	5.4	0.26	1325	12.9	0.36	11.9	0.33
Thyroid carcinoma	100	1.3	0.13	1.2	0.12	271	2.6	0.16	2.4	0.15
Other and Unspecified Epithelial Neoplasms	129	1.7	0.15	1.6	0.15	382	3.7	0.19	3.5	0.18
Other and Unspecified Malignant Tumors	43	0.6	0.09	0.6	0.10	68	0.7	0.08	0.7	0.09

[1] Rates are **per million** population and were age-adjusted by five-year age groups to the 2000 U.S. standard population based on single years of age and to the World population standard. Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for All Sites combined.
[2] See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.
[3] Abbreviations: S.E. - Standard Error; CNS - Central Nervous System.

N.B. In areas with small Latino populations, methods to indirectly identify Latinos (like NHIA) can overestimate the ethnicity-specific counts of cancer cases. Also, even small errors in Latino population estimates can affect the magnitude of the cancer rates.

Total Cancer Cases and Average Annual Age-adjusted Incidence Rates[1] for Pediatric Cancers
United States (NAACCR Combined[2]), 1998-2002, Non-Hispanic White, Females

	Ages 0-14					Ages 0-19				
		2000 U.S.		World			2000 U.S.		World	
Sites	*Cases*	*Rate*	*S.E.*[3]	*Rate*	*S.E.*	*Cases*	*Rate*	*S.E.*	*Rate*	*S.E.*
All Sites	**10012**	**143.2**	**1.43**	**148.9**	**1.51**	**15214**	**160.2**	**1.30**	**162.8**	**1.34**
Leukemias	2998	43.1	0.79	46.3	0.86	3532	37.7	0.64	40.7	0.70
Lymphoid leukemia	2362	34.0	0.70	36.7	0.77	2618	28.1	0.55	30.8	0.61
Acute non-lymphocytic leukemia	482	6.9	0.31	7.2	0.33	707	7.5	0.28	7.6	0.29
Lymphomas & Other Reticuloendothelial Neoplasms	849	11.9	0.41	11.1	0.38	2143	22.1	0.48	20.4	0.44
Hodgkin disease	380	5.2	0.27	4.5	0.23	1373	14.0	0.38	12.6	0.34
Non-Hodgkin lymphoma	319	4.5	0.25	4.4	0.25	579	6.0	0.25	5.8	0.24
CNS[3] & Misc. Intracranial & Intraspinal Neoplasms	2210	31.5	0.67	32.2	0.69	2686	28.4	0.55	29.3	0.57
Astrocytoma	1145	16.3	0.48	16.5	0.49	1440	15.2	0.40	15.5	0.41
Primitive neuroectodermal tumors	433	6.2	0.30	6.4	0.31	481	5.1	0.23	5.4	0.25
Sympathetic Nervous System Tumors	796	11.7	0.41	13.6	0.49	812	8.9	0.31	10.7	0.38
Neuroblastoma & ganglioneuroblastoma	776	11.4	0.41	13.3	0.48	781	8.6	0.31	10.4	0.37
Retinoblastoma	232	3.4	0.22	4.1	0.27	232	2.6	0.17	3.2	0.21
Renal Tumors	627	9.1	0.36	10.3	0.41	653	7.1	0.28	8.2	0.32
Wilms' tumor	605	8.8	0.36	10.0	0.41	610	6.6	0.27	7.8	0.32
Hepatic Tumors	125	1.8	0.16	2.1	0.19	148	1.6	0.13	1.8	0.15
Malignant Bone Tumors	495	6.9	0.31	6.2	0.28	797	8.2	0.29	7.6	0.27
Osteosarcoma	236	3.3	0.21	2.9	0.19	390	4.0	0.20	3.6	0.18
Ewing's sarcoma	228	3.2	0.21	2.9	0.20	335	3.5	0.19	3.3	0.18
Soft Tissue Sarcomas	692	9.8	0.37	9.9	0.38	1063	11.1	0.34	11.1	0.35
Rhabdomyosarcoma & embryonal sarcoma	308	4.4	0.25	4.7	0.27	373	4.0	0.21	4.2	0.22
Germ Cell, Trophoblastic & Other Gonadal Neoplasms	360	5.1	0.27	5.2	0.28	691	7.2	0.27	7.0	0.27
Carcinomas and Other Malignant Epithelial Neoplasms	576	8.0	0.33	7.3	0.31	2364	24.2	0.50	21.9	0.45
Thyroid carcinoma	223	3.1	0.21	2.7	0.18	1029	10.5	0.33	9.4	0.29
Other and Unspecified Epithelial Neoplasms	134	1.9	0.16	1.7	0.15	449	4.6	0.22	4.2	0.20
Other and Unspecified Malignant Tumors	52	0.7	0.10	0.8	0.11	93	1.0	0.10	1.0	0.10

[1] Rates are **per million** population and were age-adjusted by five-year age groups to the 2000 U.S. standard population based on single years of age and to the World population standard. Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for All Sites combined.
[2] See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.
[3] Abbreviations: S.E. - Standard Error; CNS - Central Nervous System.

N.B. In areas with small Latino populations, methods to indirectly identify Latinos (like NHIA) can overestimate the ethnicity-specific counts of cancer cases. Also, even small errors in Latino population estimates can affect the magnitude of the cancer rates.

Total Cancer Cases and Average Annual Age-adjusted Incidence Rates[1] for Pediatric Cancers
United States (NAACCR Combined[2]), 1998-2002, Non-Hispanic Black, Males

	Ages 0-14					Ages 0-19				
		2000 U.S.		World			2000 U.S.		World	
Sites	*Cases*	*Rate*	*S.E.*[3]	*Rate*	*S.E.*	*Cases*	*Rate*	*S.E.*	*Rate*	*S.E.*
All Sites	**1843**	**112.1**	**2.61**	**115.2**	**2.72**	**2620**	**121.8**	**2.38**	**123.2**	**2.43**
Leukemias	439	26.7	1.28	27.5	1.33	568	26.3	1.10	26.9	1.14
Lymphoid leukemia	315	19.2	1.08	19.8	1.13	379	17.5	0.90	18.1	0.94
Acute non-lymphocytic leukemia	98	6.0	0.60	6.1	0.63	144	6.7	0.56	6.7	0.57
Lymphomas & Other Reticuloendothelial Neoplasms	250	15.1	0.95	13.7	0.87	452	21.1	1.00	19.5	0.92
Hodgkin disease	92	5.6	0.58	4.9	0.52	195	9.2	0.66	8.3	0.60
Non-Hodgkin lymphoma	118	7.1	0.65	6.5	0.60	205	9.6	0.67	8.8	0.62
CNS[3] & Misc. Intracranial & Intraspinal Neoplasms	410	24.8	1.23	25.6	1.28	503	23.1	1.03	23.9	1.08
Astrocytoma	178	10.7	0.81	10.8	0.82	234	10.8	0.70	10.8	0.72
Primitive neuroectodermal tumors	101	6.1	0.61	6.4	0.64	117	5.4	0.50	5.7	0.53
Sympathetic Nervous System Tumors	144	8.9	0.74	10.2	0.85	148	6.8	0.56	8.1	0.67
Neuroblastoma & ganglioneuroblastoma	142	8.7	0.73	10.1	0.85	142	6.5	0.55	7.8	0.66
Retinoblastoma	67	4.1	0.51	4.9	0.60	68	3.2	0.38	3.9	0.47
Renal Tumors	136	8.3	0.72	9.4	0.81	151	7.0	0.57	7.9	0.65
Wilms' tumor	125	7.7	0.69	8.8	0.79	126	5.8	0.52	6.9	0.62
Hepatic Tumors	31	1.9	0.34	2.1	0.38	38	1.8	0.29	1.9	0.32
Malignant Bone Tumors	78	4.7	0.53	4.3	0.49	172	8.1	0.62	7.4	0.57
Osteosarcoma	59	3.6	0.46	3.2	0.42	132	6.2	0.54	5.7	0.50
Ewing's sarcoma	12	0.7	0.21	0.7	0.19	22	1.0	0.22	0.9	0.20
Soft Tissue Sarcomas	187	11.4	0.83	11.6	0.86	285	13.3	0.79	13.2	0.79
Rhabdomyosarcoma & embryonal sarcoma	97	5.9	0.60	6.2	0.63	131	6.1	0.53	6.3	0.55
Germ Cell, Trophoblastic & Other Gonadal Neoplasms	29	1.8	0.33	2.0	0.37	78	3.7	0.42	3.7	0.42
Carcinomas and Other Malignant Epithelial Neoplasms	64	3.9	0.49	3.5	0.44	145	6.9	0.57	6.3	0.52
Thyroid carcinoma	-	-	-	-	-	13	0.6	0.17	0.6	0.15
Other and Unspecified Epithelial Neoplasms	23	1.4	0.29	1.3	0.27	66	3.1	0.39	2.9	0.36
Other and Unspecified Malignant Tumors	8	0.5	0.17	0.5	0.17	12	0.6	0.16	0.5	0.16

[1] Rates are **per million** population and were age-adjusted by five-year age groups to the 2000 U.S. standard population based on single years of age and to the World population standard. Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for All Sites combined.
[2] See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.
[3] Abbreviations: S.E. - Standard Error; CNS - Central Nervous System.

N.B. In areas with small Latino populations, methods to indirectly identify Latinos (like NHIA) can overestimate the ethnicity-specific counts of cancer cases. Also, even small errors in Latino population estimates can affect the magnitude of the cancer rates.

Total Cancer Cases and Average Annual Age-adjusted Incidence Rates[1] for Pediatric Cancers
United States (NAACCR Combined[2]), 1998-2002, Non-Hispanic Black, Females

	Ages 0-14					Ages 0-19				
		2000 U.S.		World			2000 U.S.		World	
Sites	*Cases*	*Rate*	*S.E.*[3]	*Rate*	*S.E.*	*Cases*	*Rate*	*S.E.*	*Rate*	*S.E.*
All Sites	**1661**	**104.4**	**2.56**	**107.4**	**2.67**	**2323**	**111.3**	**2.31**	**112.9**	**2.37**
Leukemias	403	25.3	1.26	26.4	1.33	502	23.9	1.07	24.9	1.13
Lymphoid leukemia	281	17.7	1.05	18.6	1.12	321	15.2	0.85	16.2	0.92
Acute non-lymphocytic leukemia	97	6.1	0.62	6.2	0.64	136	6.5	0.56	6.6	0.57
Lymphomas & Other Reticuloendothelial Neoplasms	139	8.7	0.74	8.0	0.69	288	14.0	0.82	12.9	0.76
Hodgkin disease	52	3.2	0.45	2.8	0.39	141	6.9	0.58	6.2	0.52
Non-Hodgkin lymphoma	75	4.7	0.54	4.4	0.51	130	6.3	0.55	5.9	0.52
CNS[3] & Misc. Intracranial & Intraspinal Neoplasms	334	20.9	1.15	21.4	1.18	394	18.7	0.94	19.2	0.98
Astrocytoma	158	9.9	0.79	9.9	0.80	190	9.0	0.65	9.1	0.67
Primitive neuroectodermal tumors	75	4.7	0.55	5.0	0.59	87	4.1	0.44	4.4	0.48
Sympathetic Nervous System Tumors	111	7.1	0.67	8.1	0.77	116	5.5	0.51	6.5	0.61
Neuroblastoma & ganglioneuroblastoma	107	6.8	0.66	7.8	0.76	109	5.2	0.50	6.2	0.59
Retinoblastoma	57	3.7	0.48	4.4	0.58	57	2.7	0.36	3.4	0.45
Renal Tumors	182	11.5	0.85	12.8	0.96	193	9.2	0.66	10.4	0.76
Wilms' tumor	170	10.7	0.82	12.1	0.93	173	8.2	0.62	9.5	0.73
Hepatic Tumors	15	1.0	0.25	1.1	0.28	19	0.9	0.21	1.0	0.24
Malignant Bone Tumors	88	5.5	0.58	4.9	0.52	123	5.9	0.53	5.4	0.48
Osteosarcoma	78	4.9	0.55	4.3	0.49	98	4.6	0.47	4.3	0.43
Ewing's sarcoma	7	0.4	0.16	0.4	0.15	9	0.4	0.14	0.4	0.13
Soft Tissue Sarcomas	161	10.1	0.80	10.1	0.81	234	11.2	0.73	11.1	0.73
Rhabdomyosarcoma & embryonal sarcoma	61	3.8	0.49	4.0	0.52	77	3.7	0.42	3.8	0.44
Germ Cell, Trophoblastic & Other Gonadal Neoplasms	96	6.0	0.62	6.1	0.63	180	8.7	0.65	8.5	0.64
Carcinomas and Other Malignant Epithelial Neoplasms	64	4.0	0.50	3.6	0.46	192	9.4	0.68	8.5	0.62
Thyroid carcinoma	23	1.4	0.30	1.3	0.26	68	3.3	0.40	3.0	0.36
Other and Unspecified Epithelial Neoplasms	20	1.3	0.28	1.1	0.26	81	4.0	0.44	3.6	0.40
Other and Unspecified Malignant Tumors	11	0.7	0.21	0.6	0.19	25	1.2	0.24	1.1	0.22

[1] Rates are **per million** population and were age-adjusted by five-year age groups to the 2000 U.S. standard population based on single years of age and to the World population standard. Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for All Sites combined.
[2] See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.
[3] Abbreviations: S.E. - Standard Error; CNS - Central Nervous System.

N.B. In areas with small Latino populations, methods to indirectly identify Latinos (like NHIA) can overestimate the ethnicity-specific counts of cancer cases. Also, even small errors in Latino population estimates can affect the magnitude of the cancer rates.