

Top Five Most Commonly Diagnosed Cancers<sup>1</sup> in the U.S.  
by Race/Ethnic Group, 1999-2003  
Males

<b>All Races (total cases = 2,853,795)</b>		<b>White (total cases = 2,462,866)</b>	
Prostate	29.4%	Prostate	28.3%
Lung	15.8%	Lung	15.9%
Colon & Rectum	11.2%	Colon & Rectum	11.3%
Bladder	6.5%	Bladder	7.1%
Non-Hodgkin Lymphoma	4.1%	Non-Hodgkin Lymphoma	4.2%
<b>African American (total cases = 275,080)</b>		<b>Filipino (total cases = 10,974)</b>	
Prostate	36.9%	Prostate	30.1%
Lung	16.8%	Lung	18.3%
Colon & Rectum	10.5%	Colon & Rectum	12.8%
Kidney	3.1%	Non-Hodgkin Lymphoma	5.1%
Non-Hodgkin Lymphoma	3.1%	Liver	4.3%
<b>Japanese (total cases = 8,428)</b>		<b>Chinese (total cases = 14,305)</b>	
Prostate	28.4%	Prostate	21.6%
Colon & Rectum	17.5%	Lung	15.7%
Lung	12.0%	Colon & Rectum	15.3%
Stomach	6.7%	Liver	7.5%
Bladder	5.6%	Stomach	5.4%
<b>Vietnamese (total cases =4,727 )</b>		<b>Korean (total cases = 4,397)</b>	
Lung	18.0%	Colon & Rectum	14.4%
Liver	15.0%	Stomach	14.2%
Prostate	15.0%	Prostate	13.7%
Colon & Rectum	11.8%	Lung	13.6%
Stomach	5.8%	Liver	10.4%
<b>American Indian/Alaskan Native (total cases = 6,171)</b>		<b>Hispanic Ethnicity (of any race) (total cases = 162,173)</b>	
Prostate	21.5%	Prostate	28.5%
Lung	17.3%	Colon & Rectum	11.1%
Colon & Rectum	13.4%	Lung	10.4%
Kidney	5.5%	Non-Hodgkin Lymphoma	5.3%
Non-Hodgkin Lymphoma	4.0%	Bladder	4.2%

<sup>1</sup> Compiled by the Data Evaluation & Publication Committee for the North American Association of Central Cancer Registries, based on cases included in the NAACCR U.S. combined incidence.