Average Annual Age-adjusted (2000 U.S. Standard) NAACCR (U.S.) Combined² Cancer Incidence Rates ¹ and Rate Ratios for Selected Cancer Sites³ by Race/Ethnicity and Gender, 2005-2009

Males								
		Rates		Rate Ratios				
Cancer Site	Hispanic	Non-Hispanic White	Non-Hispanic Black	Hispanic:Non-Hispanic White	Hispanic:Non-Hispanic Black			
All Sites	418.7	556.6	631.2	0.75	0.66			
Prostate	124.9	143.2	232.6	0.87	0.54			
Colon and Rectum	46.9	53.4	66.3	0.88	0.71			
Lung and Bronchus	45.4	85.7	101.6	0.53	0.45			
NH ⁴ Lymphoma	19.3	24.4	17.3	0.79	1.12			
Kidney and Renal Pelvis	19.8	21.4	23.8	0.92	0.83			
Liver and IBD ⁴	17.5	8.3	15.2	2.11	1.15			
Urinary Bladder	20.1	41.2	19.7	0.49	1.02			
Leukemia	12.0	16.7	12.5	0.72	0.96			
Stomach	13.5	7.9	16.6	1.72	0.82			
Oral Cavity and Pharynx	10.4	17.3	16.0	0.60	0.65			
Pancreas	11.5	13.6	17.2	0.85	0.67			

Females

	Rates			Rate Ratios	
Cancer Site	Hispanic	Non-Hispanic White	Non-Hispanic Black	Hispanic:Non-Hispanic White	Hispanic:Non-Hispanic Black
All Sites	333.2	433.9	403.0	0.77	0.83
Breast	93.0	126.6	120.0	0.73	0.78
Colon and Rectum	33.3	39.8	48.7	0.84	0.68
Lung and Bronchus	26.6	60.5	52.3	0.44	0.51
Thyroid	17.2	18.9	11.0	0.91	1.57
Corpus and Uterus, NOS ⁴	20.1	25.3	22.7	0.79	0.89
NH ⁴ Lymphoma	15.3	16.8	11.7	0.91	1.30
Cervix	11.8	7.2	10.6	1.63	1.11
Ovary	11.1	13.1	9.7	0.85	1.15
Kidney and Renal Pelvis	11.4	11.1	12.3	1.02	0.92
Leukemia	8.5	10.0	7.8	0.85	1.09
Pancreas	9.9	10.2	14.1	0.97	0.70

^{1.} Rates are per 100,000 population and are age-adjusted by five-year age groups to the 2000 U.S. standard population based on single years of age. Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for the U.S. combined. Statistics exclude cases diagnosed for AL, LA, MS and TX from July 2005 through December 2005. See Introduction for more information.

² See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.

^{3.} Selected cancer sites by gender among the U.S. total population and the U.S. Hispanic/Latino population.

^{4.} Abbreviations: NH - Non-Hodgkin; IBD - Intrahepatic Bile Duct NOS - Not Otherwise Specified

N.B. In areas with small Latino populations, methods to indirectly identify Latinos (like NHIAv2) can overestimate the ethnicity-specific counts of cancer cases. Also, even small errors in Latino population estimates can affect the magnitude of the cancer rates.