Bladder (C67.0-C67.9)

А.

- Summary Stage Rule Changes1. Involvement of the common iliac lymph nodes Regional (SS1977) to Distant (SS2000)
- 2. Extension in males only to: pubic bone rectum Regional (SS1977) to Distant (SS2000)

B. **Stage Shift**

Stage	<u>SS 1977</u>	<u>SS 2000</u>		
Localized	84.8%	84.8%		
Regional-Ext	5.7%	5.7%		
Regional-Node	0.8%	0.8%		
Reg-Ext + Node	1.7%	1.6%		
Distant	3.7%	3.7%		
Unknown	3.4%	3.4%		
Total	100.0%	100.0%		

C. **Data Interpretation**

The distribution of summary stage 1977 and summary stage 2000 codes is very similar. However, individual cases may be classified differently according to the two systems, and thus for some uses, this impact should be reviewed for comparability.

Indirect Coding to Summary Stage 1977 and Summary Stage 2000 based on Extent of Disease Codes SEER Program Data, 1998-2001, Bladder

ss1977	ss2000							
	0: In Situ	1: Localized	2: Regional	3: Reg LN	4: Reg + LN	7: Distant	9: Unknown	Total
1: Localized		23831	0	0	0	0	0	23831
2: Regional		0	1589	0	0	0	0	1589
3: Reg LN		0	0	225	0	3	0	228
4: Reg + LN		0	0	0	461	11	0	472
7: Distant		0	0	0	0	1032	0	1032
9: Unknown		0	0	0	0	0	957	957
Total		23831	1589	225	461	1046	957	28109
Proportion	0: In Situ	1: Localized	2: Regional	3: Reg LN	4: Reg + LN	7: Distant	9: Unknown	Total
1: Localized		0.847807	0	0	0	0	0	0.847807
2: Regional		0	0.05653	0	0	0	0	0.05653
3: Reg LN		0	0	0.008005	0	0.000107	0	0.008111
4: Reg + LN		0	0	0	0.0164	0.000391	0	0.016792
7: Distant		0	0	0	0	0.036714	0	0.036714
9: Unknown		0	0	0	0	0	0.034046	0.034046
Total		0.847807	0.05653	0.008005	0.0164	0.037212	0.034046	1

p0	0.999502
ре	0.724838
K	0.99819

