

Cervix Uteri (C 53.0-C53.1, C53.8-C53.9)

A. Summary Stage Rule Changes

1. Extension to:
 ureter, intra- and extramural
 vulva
 fallopian tubes
 ovary (ies),
 urethra
 frozen pelvis
Distant (SS1977) to Regional by direct extension only (SS2000)

B. Stage Shift

<u>Stage</u>	<u>SS 1977</u>	<u>SS 2000</u>
Localized	52.9%	52.9%
Regional-Ext	23.4%	23.6%
Regional-Node	3.6%	3.6%
Reg-Ext + Node	4.0%	4.1%
Distant	10.5%	10.3%
Unknown	5.6%	5.6%
Total	100.0%	100.0%

	} 31.0%		
	} 31.3%		

Data Interpretation

The distribution of Summary Stage 1977 and Summary Stage 2000 is very similar. However, individual cases may be classified differently according to the two systems, and thus for some uses, this impact should be reviewed for comparability.

Indirect Coding to Summary Stage 1977 and Summary Stage 2000 based on Extent of Disease Codes
SEER Program Data, 1998-2001, Cervix Uteri

ss1977

ss2000

	0: In Situ	1: Localized	2: Regional	3: Reg LN	4: Reg + LN	7: Distant	9: Unknown	Total
1: Localized		3800	0	0	0	0	0	3800
2: Regional		0	1684	0	0	0	0	1684
3: Reg LN		0	0	255	0	0	0	255
4: Reg + LN		0	0	0	286	0	0	286
7: Distant		0	11	0	4	742	0	757
9: Unknown		0	0	0	0	0	402	402
Total		3800	1695	255	290	742	402	7184

Proportion:	0: In Situ	1: Localized	2: Regional	3: Reg LN	4: Reg + LN	7: Distant	9: Unknown	Total
1: Localized		0.528953	0	0	0	0	0	0.528953
2: Regional		0	0.23441	0	0	0	0	0.23441
3: Reg LN		0	0	0.035496	0	0	0	0.035496
4: Reg + LN		0	0	0	0.039811	0	0	0.039811
7: Distant		0	0.001531	0	0.000557	0.103285	0	0.105373
9: Unknown		0	0	0	0	0	0.055958	0.055958
Total		0.528953	0.235941	0.035496	0.040367	0.103285	0.055958	1

Simple Kappa 0.9968

p0 0.997912
pe 0.35198
K 0.996778

