

Stomach (C 16)

A. Summary Stage Rule Changes

1. Any subsite of stomach (C16.0-C16.6, C16.8-C16.9) with intraluminal spread (only) to esophagus or duodenum.
Regional Extension (SS1977) to Localized (SS2000)
2. Any subsite of stomach (C16.0-C16.6, C16.8-C16.9) with linitis plastica (diffuse involvement of the entire stomach wall).
Regional Extension (SS1977) to Localized (SS2000)
3. Any subsite of stomach (C16.0-C16.6, C16.8-C16.9) involving celiac or hepatic lymph nodes.
Distant (SS1977) to Regional Lymph Node (SS2000)

B. Stage Shift

<u>Stage</u>	<u>SS 1977</u>	<u>SS 2000</u>
Localized	23.3%	23.3%
Regional-Ext	7.0%	7.0%
Regional-Node	9.2%	9.8%
Reg-Ext + Node	13.1%	14.1%
Distant	33.7%	31.9%
Unknown	13.7%	13.8%
Total	100.0%	100.0%

}	29.3%	}	30.9%
---	-------	---	-------

Data Interpretation

The distribution of Summary Stage 1977 and Summary Stage 2000 codes is very similar. However, individual cases may be classified differently according to the two systems, and thus for some uses, this impact should be reviewed for comparability.

Indirect Coding to Summary Stage 1977 and Summary Stage 2000 based on Extent of Disease Codes
SEER Program Data, 1998-2001, Stomach

ss1977

ss2000

	1: Localized	2: Regional	3: Reg LN	4: Reg + LN	7: Distant	9: Unknown	Total
1: Localized	2858	0	0	0	0	0	2858
2: Regional	0	864	0	0	0	0	864
3: Reg LN	0	0	1135	0	0	0	1135
4: Reg + LN	0	0	0	1609	0	0	1609
7: Distant	0	0	72	129	3923	20	4144
9: Unknown	0	0	0	0	0	1678	1678
Total	2858	864	1207	1738	3923	1698	12288

Proportions	1: Localized	2: Regional	3: Reg LN	4: Reg + LN	7: Distant	9: Unknown	Total
1: Localized	0.232585	0	0	0	0	0	0.232585
2: Regional	0	0.070313	0	0	0	0	0.070313
3: Reg LN	0	0	0.092367	0	0	0	0.092367
4: Reg + LN	0	0	0	0.130941	0	0	0.130941
7: Distant	0	0	0.005859	0.010498	0.319255	0.001628	0.33724
9: Unknown	0	0	0	0	0	0.136556	0.136556
Total	0.232585	0.070313	0.098226	0.141439	0.319255	0.138184	1

p0 0.982
pe 0.213
K 0.977

