

Corpus Uteri; Uterus, NOS
(C 54.0-C54.3, C54.8-C54.9, C55.9)

A. Summary Stage Rule Changes

1. Extension or metastasis to:
 - ureter (extension)
 - vulva (extension)
 - bladder, NOS excluding mucosa (metastasis)
 - bladder wall (metastasis)
 - rectal wall, NOS (metastasis)
 - rectum, NOS excluding mucosa (metastasis)
 - pelvis wall (metastasis)
 - frozen pelvis (metastasis)
 - vagina (extension or metastasis)*Distant (SS1977) to Regional by direct extension only (SS2000)*

2. Involvement of superficial inguinal (femoral) lymph nodes
Regional (SS1977) to Distant (SS2000)

B. Stage Shift

<u>Stage</u>	<u>SS 1977</u>	<u>SS 2000</u>
Localized	69.8%	69.8%
Regional-Ext	12.5%	13.3%
Regional-Node	1.8%	1.8%
Reg-Ext + Node	2.9%	3.2%
Distant	8.6%	7.4%
Unknown	4.4%	4.4%
Total	100.0%	100.0%

C. Data Interpretation

The distribution of summary stage 1977 and summary stage 2000 codes is very similar. However, individual cases may be classified differently according to the two systems, and thus for some uses, this impact should be reviewed for comparability.

Indirect Coding to Summary Stage 1977 and Summary Stage 2000 based on Extent of Disease Codes
SEER Program Data, 1998-2001, Corpus Uteri and Uterus NOS

ss1977

ss2000

	0: In Situ	1: Localized	2: Regional	3: Reg LN	4: Reg + LN	7: Distant	9: Unknown	Total
1: Localized		13080	0	0	0	0	0	13080
2: Regional		0	2334	0	0	0	0	2334
3: Reg LN		0	0	337	0	4	0	341
4: Reg + LN		0	0	0	528	7	0	535
7: Distant		0	170	0	65	1374	0	1609
9: Unknown		0	0	0	0	2	828	830
Total		13080	2504	337	593	1387	828	18729

Proportion:	0: In Situ	1: Localized	2: Regional	3: Reg LN	4: Reg + LN	7: Distant	9: Unknown	Total
1: Localized		0.698382	0	0	0	0	0	0.698382
2: Regional		0	0.12462	0	0	0	0	0.12462
3: Reg LN		0	0	0.017993	0	0.000214	0	0.018207
4: Reg + LN		0	0	0	0.028192	0.000374	0	0.028565
7: Distant		0	0.009077	0	0.003471	0.073362	0	0.08591
9: Unknown		0	0	0	0	0.000107	0.04421	0.044316
Total		0.698382	0.133696	0.017993	0.031662	0.074056	0.04421	1

p0 0.986759
pe 0.513952
K 0.972757

