Cancer in North America: 2010-2014
Volume One: Combined Cancer Incidence for the United States, Canada, and North America

North American Association of Central Cancer Registries, Inc.
NAACCR
North American Association of Central Cancer Registries

EXECUTIVE SUMMARY

Members of the North American Association of Central Cancer Registries, Inc. (NAACCR) participate voluntarily in an annual call for data to develop a multi-registry, aggregated data resource for cancer surveillance and research. In June of 2017, NAACCR released the four volume monograph, *Cancer in North America (CiNA): 2010-2014.* This release of cancer incidence statistics for the United States is the 27th publication in the series and the 25th release that includes statistics for both Canada and the United States.

This year's publication includes cancer incidence data for 50 states, the District of Columbia, Puerto Rico, California (excluding the Greater Bay area and Los Angeles), 4 SEER urban registries, and 12 Canadian registries. CiNA includes registry-specific data on stage at diagnosis, survival and delay adjusted estimates of counts and age-adjusted rates for selected cancers. These delay adjustment figures provide a projection of the likely volume of tardy reports of cancer for the time period and is recommended for use in assessing current cancer incidence trends.

Volume One presents the aggregated cancer incidence data, including pediatric cancer and cancer by stage at diagnosis, representing North America including data from the high quality registries in the United States and Canada. Age-adjusted cancer incidence rates for the Canadian and United States populations are provided standardized to the 2000 US, 1996 Canadian, and the world population standards. Combined United States data are presented by race and for Hispanic/Latino ethnicity. New for this year are registry specific delay adjusted rates in Volume One. Volume Two contains registry-specific cancer incidence rates by cancer site, sex, race, ethnicity and stage for all NAACCR members submitting data for inclusion in the monograph. Volume Three presents the registry-specific cancer death rates by cancer site, sex, race, and ethnicity. Volume Four provides cancer survival data for the US and Canada from 50 registries on more than 8 million cases diagnosed among North Americans between 2007 and 2013. All four volumes are available free of charge from the NAACCR website, along with population data, supporting appendices, and cancer rates age-adjusted to the US, Canadian and World population standards.

Note that this publication utilizes population estimates that are based upon the United States 2010 and Canada 2011 censuses.

In addition to publication of cancer statistics for the most recent five-year period, the *CiNA* data resource is used to create the *CiNA+ Online* database and NAACCR FastStats (graphic and tables by cancer sites, race/sex, race/ethnicity, age at diagnosis, sex, registry, and data type) for public queries regarding cancer incidence for the most recent five years. The CiNA data are also used to develop a discretionary research data set (*CiNA Deluxe)*, including data from 1995 to 2014, to facilitate analytic studies conducted by epidemiologists and other qualified researchers. The data within this report are also available as an electronic reproduction of all cancer statistics reported in the CiNA monograph for use in SAS analytic software and a discretionary data set of the data in *CiNA+ Online* (named *CiNA Plus in SEER*Stat)* in SEER*Stat software for approved uses by NAACCR members. These preeminent cancer data resources are unparalleled in geographic scope, timeliness of statistics, and assurance of standardization of information and data quality across all contributors, with standard data quality metrics reported for all contributors.

We hope that *Cancer in North America: 2010-2014*, with the companion *CiNA* products and resources, facilitate studies of cancer burden, so that we are better able to identify and understand appropriate and important measures to control the myriad diseases within the cancer rubric. The cancer surveillance infrastructure in the U.S. and Canada has been orchestrated to meet these objectives.

This publication is made possible by the continuing efforts of the NAACCR member registries. High quality standardized cancer incidence data aggregated across the states, provinces, territories, and regions in North America is made possible by the dedication of our members to cancer surveillance.

The Editors would like to acknowledge the fine work and support of the National Cancer Institute, the National Center for Health Statistics and Statistics Canada for their assistance in developing this publication.

The Editors
June 2017

CANCER IN NORTH AMERICA: 2010-2014

VOLUME ONE: COMBINED CANCER INCIDENCE FOR THE UNITED STATES, CANADA AND NORTH AMERICA

A Publication of the North American Association of Central Cancer Registries, Inc. (NAACCR)

Editors:
Glenn Copeland, Michigan Cancer Surveillance Program
Andrew Lake, Information Management Services, Inc.
Rick Firth, Information Management Services, Inc.
Brad Wohler, Florida Cancer Data System
Xiao-Cheng Wu, Louisiana Tumor Registry
Maria J. Schymura, New York State Cancer Registry
Prithwish De, Ontario Cancer Registry at Cancer Care Ontario
Jim Hofferkamp, NAACCR, Inc.
Recinda Sherman, NAACCR Inc.
Betsy Kohler, NAACCR, Inc.

June 2017

The Editorial Subcommittee of the NAACCR Standing Committee, Data Use and Research Committee publishes this monograph. We are grateful to the National Cancer Institute, National Institutes of Health (NCI/NIH) for providing support for the production under Contract No. HHSN261200444001C/ADB Contract No. PC-44401. We acknowledge partial support of NAACCR staff by cooperative agreement U75/CCU52346-05 from the Centers for Disease Control and Prevention (CDC) and by the NCI/NIH Contract No. HHSN261200444001C/ADB Contract No. PC-44401. Its contents are solely the responsibility of the editors and do not necessarily represent the official view of CDC.

CINA PARTICIPANTS
2010-2014

SUGGESTED CITATION:
Copeland G, Lake A, Firth R, Wohler B, Wu XC, Schymura M, De P, Hofferkamp J, Sherman R, Kohler B (eds). *Cancer in North America: 2010-2014. Volume One: Combined Cancer Incidence for the United States, Canada and North America*. Springfield, IL: North American Association of Central Cancer Registries, Inc. June 2017.

TABLE OF CONTENTS

SECTION I: Introduction and Technical Notes I-1

SECTION II: Top Five Most Commonly Diagnosed Cancers in North America
Top Five Most Commonly Diagnosed Cancers in North America II-1
Top Five Most Commonly Diagnosed Cancers in the U.S. by Race/Ethnicity
Males II-2
Females II-3

SECTION III: Average-annual Age-adjusted NAACCR Combined Cancer Incidence Rates for the United States, Canada and North America
Data Quality Indicators for the United States III-1
Map: Registries included in U.S. Combined Incidence for All Races III-3
Combined Registries, All Races (2000 U.S. and 1996 Canadian populations) III-4
Five Most Commonly Diagnosed Cancers, Incidence Rates by Registry,
All Races, Males (2000 U.S. population) III-5
All Races, Males (1996 Canadian population) III-7
All Races, Females (2000 U.S. population) III-9
All Races, Females (1996 Canadian population) III-11
Map: Registries included in U.S. Combined Incidence for White III-13
Combined Registries, White (2000 U.S. and 1996 Canadian populations) III-14
Five Most Commonly Diagnosed Cancers, Incidence Rates by Registry,
White, Males (2000 U.S. population) III-15
White, Males (1996 Canadian population) III-17
White, Females (2000 U.S. population) III-19
White, Females (1996 Canadian population) III-21
Map: Registries included in U.S. Combined Incidence for Black III-23
Combined Registries, Black (2000 U.S. and 1996 Canadian populations) III-24
Five Most Commonly Diagnosed Cancers, Incidence Rates by Registry,
Black, Males (2000 U.S. population) III-25
Black, Males (1996 Canadian population) III-27
Black, Females (2000 U.S. population) III-29
Black, Females (1996 Canadian population) III-31
Map: Registries included in U.S. Combined Incidence for Asian/Pacific Islander III-33
Combined Registries, Asian/Pacific Islander (2000 U.S. and 1996 Canadian populations) III-34
Five Most Commonly Diagnosed Cancers, Incidence Rates by Registry,
Asian/Pacific Islander, Males (2000 U.S. population) III-35
Asian/Pacific Islander, Males (1996 Canadian population) III-37
Asian/Pacific Islander, Females (2000 U.S. population) III-39
Asian/Pacific Islander, Females (1996 Canadian population) III-41
Map: Selected CHSDA Counties included in U.S. Combined Incidence for American Indian/Alaskan Native III-43
Combined Registries, American Indian/Alaskan Native (2000 U.S. and 1996 Canadian populations) III-44
Five Most Commonly Diagnosed Cancers, Incidence Rates by Registry,
American Indian/Alaskan Native, Males (2000 U.S. population) III-45
American Indian/Alaskan Native, Males (1996 Canadian population) III-47
American Indian/Alaskan Native, Females (2000 U.S. population) III-49
American Indian/Alaskan Native, Females (1996 Canadian population) III-51
Most Frequent Cancer Sites by IHS Region,
American Indian/Alaskan Native, Males (2000 U.S. population) III-53
American Indian/Alaskan Native, Females (2000 U.S. population) III-55
Map: Registries included in U.S. Combined Incidence for Hispanic/Latino, All Races III-57
Combined Registries, Hispanic/Latino, All Races (2000 U.S. and 1996 Canadian populations) III-58
Commonly Diagnosed Cancers, Incidence Rates by Registry,
Hispanic/Latino, All Races, Males (2000 U.S. population) III-59
Hispanic/Latina, All Races, Females (2000 U.S. population) III-63

Contents

Map: Registries included in U.S. Combined Incidence for Non-Hispanic White III-67
Combined Registries, Non-Hispanic White (2000 U.S. and 1996 Canadian populations) III-68
Commonly Diagnosed Cancers, Incidence Rates by Registry,
Non-Hispanic White, Males (2000 U.S. population) III-69
Non-Hispanic White, Females (2000 U.S. population) III-73
Map: Registries included in U.S. Combined Incidence for Non-Hispanic Black III-77
Combined Registries, Non-Hispanic Black (2000 U.S. and 1996 Canadian populations) III-78
Commonly Diagnosed Cancers, Incidence Rates by Registry,
Non-Hispanic Black, Males (2000 U.S. population) III-79
Non-Hispanic Black, Females (2000 U.S. population) III-83
Data Quality Indicators for Canada III-88
Map: Registries included in Canadian Combined Incidence for All Races III-89
Combined Registries, All Races (2000 U.S. and 1996 Canadian populations) III-90
Five Most Commonly Diagnosed Cancers, Incidence Rates by Registry,
All Races, Males (2000 U.S. and 1996 Canadian populations) III-91
All Races, Females (2000 U.S. and 1996 Canadian populations) III-92
Data Quality Indicators for North America III-94
Map: Registries included in North America Combined Incidence for All Races III-95
Combined Registries, All Races (2000 U.S. and 1996 Canadian populations) III-96
Five Most Commonly Diagnosed Cancers, Incidence Rates by Registry,
All Races, Males (2000 U.S. population) III-97
All Races, Males (1996 Canadian population) III-99
All Races, Females (2000 U.S. population) III-101
All Races, Females (1996 Canadian population) III-103
Map: Registries included in U.S. Pediatric Cancer Incidence III-106
Age-specific Rates for the United States
All Races, Males III-107
All Races, Females III-108
White, Males III-109
White, Females III-110
Black, Males III-111
Black, Females III-112
Asian/Pacific Islander, Males III-113
Asian/Pacific Islander, Females III-114
American Indian/Alaskan Native, Males III-115
American Indian/Alaskan Native, Females III-116
Hispanic/Latino, All Races, Males III-117
Hispanic/Latina, All Races, Females III-118
Non-Hispanic White, Males III-119
Non-Hispanic White, Females III-120
Non-Hispanic Black, Males III-121
Non-Hispanic Black, Females III-122
Age-adjusted Rates (2000 U.S. and 1996 Canadian population) (Ages 0-14, Ages 0-19)
All Races, Males III-123
All Races, Females III-124
White, Males III-125
White, Females III-126
Black, Males III-127
Black, Females III-128
Asian/Pacific Islander, Males III-129
Asian/Pacific Islander, Females III-130
American Indian/Alaskan Native, Males III-131
American Indian/Alaskan Native, Females III-132
Hispanic/Latino, All Races, Males III-133
Hispanic/Latina, All Races, Females III-134
Non-Hispanic White, Males III-135
Non-Hispanic White, Females III-136
Non-Hispanic Black, Males III-137
Non-Hispanic Black, Females III-138
Map: Registries included in Canadian Pediatric Cancer Incidence III-140
Age-specific Rates for Canada
All Races, Males III-141
All Races, Females III-142

Contents

Age-adjusted Rates (2000 U.S. and 1996 Canadian population) (Ages 0-14, Ages 0-19)
All Races, Males III-143
All Races, Females III-144
Map: Registries included in North American Pediatric Cancer Incidence III-146
Age-specific Rates for North America
All Races, Males III-147
All Races, Females III-148
Age-adjusted Rates (2000 U.S. and 1996 Canadian population) (Ages 0-14, Ages 0-19)
All Races, Males III-149
All Races, Females III-150

SECTION IV: Stage at Diagnosis by Age for Select Cancer Sites
United States Combined Registries IV-1
Canada Combined Registries IV-3
North America Combined Registries IV-4

SECTION V: Average-annual Age-adjusted Cancer Incidence Observed and Delay Adjusted Counts and Rates by Race, Sex and Selected Sites, for the United States, Canada and North America
Map: Registries included in U.S. Combined Delay Adjusted Incidence for All Races V-1
Combined Registries, All Races (2000 U.S. population) V-2
All Races, Males (2000 U.S. population) V-3
All Races, Females (2000 U.S. population) V-9
Map: Registries included in U.S. Combined Delay Adjusted Incidence for White V-15
Combined Registries, White (2000 U.S. population) V-16
White, Males (2000 U.S. population) V-17
White, Females (2000 U.S. population) V-23
Map: Registries included in U.S. Combined Delay Adjusted Incidence for Black V-29
Combined Registries, Black (2000 U.S. population) V-30
Black, Males (2000 U.S. population) V-31
Black, Females (2000 U.S. population) V-37
Map: Registries included in U.S. Combined Delay Adjusted Incidence for Asian/Pacific Islander V-43
Combined Registries, Asian/Pacific Islander (2000 U.S. population) V-44
Asian/Pacific Islander, Males (2000 U.S. population) V-45
Asian/Pacific Islander, Females (2000 U.S. population) V-51
Map: Selected CHSDA Counties included in U.S. Combined Incidence for American Indian/Alaskan Native V-57
Combined Registries, American Indian/Alaskan Native (2000 U.S. population) V-58
American Indian/Alaskan Native, Males (2000 U.S. population) by Region V-59
American Indian/Alaskan Native, Females (2000 U.S. population) by Region V-60
Map: Registries included in U.S. Combined Delay Adjusted Incidence for Hispanic V-61
Combined Registries, Hispanic (2000 U.S. population) V-62
Hispanic, Males (2000 U.S. population) V-63
Hispanic, Females (2000 U.S. population) V-69
Map: Registries included in U.S. Combined Delay Adjusted Incidence for Non-Hispanic White V-75
Combined Registries, Non-Hispanic White (2000 U.S. population) V-76
Non-Hispanic White, Males (2000 U.S. population) V-77
Non-Hispanic White, Females (2000 U.S. population) V-83
Map: Registries included in U.S. Combined Delay Adjusted Incidence for Non-Hispanic Black V-89
Combined Registries, Non-Hispanic Black (2000 U.S. population) V-90
Non-Hispanic Black, Males (2000 U.S. population) V-91
Non-Hispanic Black, Females (2000 U.S. population) V-97
Map: Registries included in Canadian Combined Delay Adjusted Incidence V-103
Combined Registries, All Races (2000 U.S. population) V-104
All Races, Males (2000 U.S. population) V-105
All Races, Females (2000 U.S. population) V-106
Map: Registries included in North American Combined Delay Adjusted Incidence V-107
Combined Registries, All Races (2000 U.S. population) V-108

All Appendices are available on the NAACCR website:
https://www.naaccr.org/cancer-in-north-america-cina-volumes/#Appendices

Section I

Introduction and Technical Notes

Cancer in North America: 2010-2014

INTRODUCTION

The North American Association of Central Cancer Registries, Inc. (NAACCR) is a professional organization that develops and promotes uniform data standards for cancer registration; provides education and training; certifies population based registries; aggregates and publishes data from central cancer registries; and promotes the use of cancer surveillance data and systems for cancer control and epidemiologic research, public health programs, and patient care to reduce the burden of cancer in North America. NAACCR annually produces this statistical monograph on cancer in North America to provide cancer incidence and mortality statistics for the United States, Canada, and North America.

This year marks the 27th annual release of the series of Cancer in North America (CiNA) monographs. The 2017 monograph includes data from 69 central population-based registries: 57 from the United States (50 states, the District of Columbia, Puerto Rico, California (excluding the Greater Bay area and Los Angeles) and 4 metropolitan areas) and 12 from Canada (9 provinces and 3 territories).

The submitted data reflect 97.7% population coverage of Canada and the United States. Combined, five-year average annual incidence statistics for the United States, for Canada and for North America are created from registries that met or exceeded the NAACCR standards of "fitness for use" in aggregated incidence data for all years, 2010-2014.

Volume One presents the combined U.S. and Canadian statistics for cancer incidence rates across North America. The combined statistics presented in Volume One reflect 96.7% of the U.S. population (47 states and the District of Columbia) and includes statistics from the 9 Canadian registries that met the CiNA inclusion requirements, reflecting 38.0% of the Canadian population.

NAACCR issues an annual call for data to member registries to create the CiNA monograph and other useful CiNA products for data analyses and cancer incidence information (e.g., CiNA+ Online, NAACCR FastStats, CiNA in SEER*Stat, CiNA Plus in SEER*Stat, CiNA Monograph data in SAS Data set and CiNA Deluxe). For more information on these products and the process for gaining access to them, please visit www.NAACCR.org and select *Cancer in North America CiNA Data Products* under the *Research & Analytic Tools* tab.

This monograph is a testament to the continuing commitment by population-based cancer registries throughout the United States and Canada The efforts of the NAACCR membership to collect timely, complete, and accurate data make this publication possible. An Editorial Board of member volunteers, with support from NAACCR staff and staff from Information Management Services, takes on the responsibility to oversee the call for data, evaluate all submissions, and make all editorial decisions in production of the monograph. The objective of this work is to provide timely and useful information on cancer incidence and mortality for all geographic areas and race/ethnic groups in the United States and Canada.

The publication includes U.S. cancer incidence and mortality for white, black, Hispanic/Latino and Asian/Pacific Islander as combined and registry specific rates and American Indian and Alaskan Native populations presented for U.S. counties located in Indian Health Service Contract Health Service Delivery Areas (CHSDA) of the United States (excepting only for Kansas).

The 2010-2014 Cancer Incidence in North America monograph, registry specific delay adjusted rates are provided to adjust for reporting delay. Each year, registries submit cases approximately two years after the close of a calendar year (e.g. cases for 2014 were first submitted in the December 2016 NAACCR Call For Data submission). These case counts are then revised in each subsequent submission. Delay adjusted rates represent the observed rate adjusted to estimate what will be the "eventual" case count, providing more accurate trends and levels of incidence rates.

Starting in 2015, a joint effort by NCI, CDC, and the North American Association of Central Cancer Registries (NAACCR) was mounted to develop a unified approach for estimating and reporting delay-adjusted rates across all of the U.S. and Canada. The delay-adjusted rates for NAACCR registries, including SEER registries reported in the Cancer Statistics Review, is based on the data submitted to the North American Association of Central Cancer Registries (NAACCR). Statistical models to estimate delay are estimated for each registry by age (<50, 50-69, 70+), gender (male and female) and race/ethnicity (all races combined, white, black, API, AIAN CHSDA region, Hispanic, non-Hispanic white and non-Hispanic black), and delay time (for the 2016 submission a delay of 2 years for diagnosis year 2014, 3 years for diagnosis year 2013, etc.) While the statistical models estimate reporting delay registry by age, race, gender, and delay time, delay adjusted rates for any grouping is done in SEER*Stat by weighting each group appropriately.

In this year's release, a different and improved statistical model has been used with the goal of producing more stable estimates. The details of the new modeling are available in the Development of the Delay Model of the website http://www.surveillance.cancer.gov/delay/.

For some registries, there were insufficient historical data available to calculate registry specific delay factors. In order to calculate delay adjusted national statistics, delay factors were applied to registries with missing data using an average of the U.S. or Canadian delay factors.

CONTENTS OF THE CiNA MONOGRAPH

This year the CiNA monograph is comprised of four volumes as described below. Rates are presented as average-annual, age-adjusted rates using three different standard million populations: the 2000 U.S., the 1996 Canadian, and the World standard populations. We believe that presenting cancer rates adjusted to all three standards enhances the meaning and utility of the rates for users in the United States, Canada, and international settings. Recent volumes of the CiNA publication, including sections containing incidence rates standardized to the World Standard, and the population counts used in the calculation of cancer rates are available for download on the NAACCR website, https://www.naaccr.org/cancer-in-north-america-cina-volumes/.

Volume One, Combined Cancer Incidence for the United States, Canada, and North America presents cancer incidence data that have been combined to create five-year, average-annual data for cancers occurring in all persons and for children, aged 0 to 14 and 0 to 19. Registries submitting the data must meet the NAACCR criteria of high quality incidence data indicating that the data are fit for use at the time of data submission. The criteria for the standard are described in detail below in the Data Quality Indicator section. In the United States, cancer counts and incidence rates are presented for all races combined, black, white, Asian/Pacific Islander, American Indian/Alaskan Native, and for Hispanic/Latino, non-Hispanic white, non-Hispanic black populations.

It should be noted that cancer incidence data for two of the largest provinces in Canada, namely Ontario and Quebec, are not included in Volume One as they did not submit data that met NAACCR standards for the December 2016 data call. These two provinces represent over 60% of new cancer cases in Canada. As a result, the statistics for Canada should be interpreted with this in mind.

Volume One is comprised of four data sections:

Section two includes the five most common types of cancer by sex in North America, the United States and Canada based on the combined United States and the combined Canadian data. The section also includes the top five most common types of cancer by sex, race and ethnicity in the United States based on the combined United States data.

Section three provides cancer incidence rates for the United States, Canada and North America. The rates for the United States are presented for all races combined, white, black, Asian or Pacific Islander, American Indian/Alaska Native, Hispanic, non-Hispanic white and non-Hispanic black populations for all ages combined. The rates for Canada and North America are presented for all races combined and for all ages combined. Also included in section three are tables of cancer rates for children, aged 0-14 and 0-19 for the United States, Canada

and North America. As with the tables for all ages, the pediatric tables are presented by race and ethnicity for the United States and for all races combined for Canada and North America.

Section four contains the frequency counts, percent distribution and age-adjusted rates of cancer by derived summary stage at diagnosis (derived from and following the Collaborative Stage staging system) for female breast, and for prostate, lung, uterine cervix and colorectal cancers. Combined United States data are provided for white, black, Asian or Pacific Islander, and American Indian/Alaska Native populations with American Indian/Alaska Native data limited to Contract Health Service Delivery Areas counties. In addition, combined United States data, combined Canadian data and combined North America data are provided for all races combined. U.S. data are also provided by Hispanic ethnicity.

Section five provides cancer counts and rates for the time period 2010-2014 that are "delay-adjusted." This adjustment is done to estimate the number of incident cancer cases that were diagnosed in the time period but whose reporting to a central registry is delayed. These estimates are derived from historic patterns of reporting lag adjusted by various factors including cancer type, sex, age and race/ethnicity. Registries otherwise included in Volume One data are excluded from Section five, if they did not supply the data needed to calculate the delay adjustment for that registry. There were three registries affected by this with one from the U.S. and two from Canada. A U.S. average and a Canadian average delay factor was calculated and applied to these excluded registries to assure the combined delay-adjusted data are consistent with the incidence rates otherwise presented in Volume One. For more information on the importance of this methodology see the articles by Clegg, Midthune and Zou listed in the references at the end of this section or visit http://surveillance.cancer.gov/delay/.

Volume Two, Registry-specific Cancer Incidence in the United States and Canada presents cancer incidence data for NAACCR-member, population-based central cancer registries in Canada and the United States that have agreed to participate in the CiNA monograph. Five-year averages of data are presented for the years 2010 to 2014.

Each set of data tables includes demographic and data quality information and registry descriptions to help interpret the statistics reported. The 2017 publication includes registry specific data on stage at diagnosis for female breast, lung, colon and rectum, prostate and uterine cervix cases for all registries submitting stage data for 2010 through 2014. In the United States, cancer incidence rates are calculated for the following race/ethnic populations: all races combined, white, black, Asian or Pacific Islander, American Indian/Alaskan Native for CHSDA counties, Hispanic/Latino (all races), non-Hispanic white, and non-Hispanic black. Race and ethnicity information is not collected in Canada, and thus is not presented.

Volume Three, Registry-specific Cancer Mortality in the United States and Canada presents cancer mortality data for all geographic areas in the United States and Canada.

Volume Four, Cancer Survival Rates in the United States and Canada includes data from 50 registries on more than 8 million cases diagnosed among North Americans between 2007 and 2013.

Volume Four is comprised of two data sections:

Section two includes counts, relative survival ratios and confidence intervals for the United States, Canada and North America combined. These statistics are presented for all races by sex and select cancer sites. The tables for each cancer site present statistics by a total, stage and specific age groups for each region. In the United States combined, survival statistics by white and black race are also presented.

Section three includes counts, relative survival ratios and confidence intervals by registry, sex and select cancer sites. Survival statistics are available for the United States and Canadian registries by all races and for the United States registries by white and black race.

SOURCES OF DATA

Cancer Incidence. A cancer registry must be population-based and a NAACCR member in good standing to be included in NAACCR data publications. All cancer registries in the United States and Canada are NAACCR members, including the three territories in Canada and three U.S. territories. An annual request is sent to all members to submit voluntarily a data file for use in CiNA, CiNA research and data information products, the U.S. *Annual Report to the Nation*, and the American Cancer Society's *Cancer Facts and Figures* annual publication. All NAACCR member registries receive support from the state, province or territory. In the U.S., they also participate in the National Cancer Institute's Surveillance, Epidemiology, and End Results (SEER) Program or the Centers for Disease Control and Prevention's National Program of Cancer Registries (NPCR) or both. In Canada, all registries submit data to the Canadian Cancer Registry maintained by Statistics Canada.

Mortality Data. Mortality data for 2010 to 2014 for United States and Puerto Rico were obtained from the National Center for Health Statistics (NCHS) of the Centers for Disease Control and Prevention (CDC), as provided to the National Cancer Institute (NCI). The Canadian mortality data for 2010 to 2014 were not yet available from Statistics Canada at initial publication, so data for the years 2009-2013 were used for completeness estimations. These data are available from Statistics Canada's Vital Statistics – Death Database (http://www23.statcan.gc.ca/imdb/p2SV.pl?Function=getSurvey&SDDS=3233).

Population Estimates. To assist the reader, population estimates by age, sex, and race/ethnicity (where relevant) used in the calculation of rates in the monograph are available for each central registry, the United States and Canada. These data are available online on the NAACCR website at https://www.naaccr.org/cancer-in-north-america-cina-volumes/.

Population estimates for 2010 through 2014 for the United States, individual states, and SEER metropolitan areas were obtained from the SEER program. These estimates were based on United States Census Bureau population estimates for these years and represent a modification of the annual time series population estimates produced by the Population Estimates Program of the Census Bureau with support from the NCI. The population estimates incorporate bridged, single race estimates that are derived from the original multiple race categories in the 2010 Census. These bridged estimates are consistent with the four race groups enumerated in the 1990 Census and were produced under a collaborative arrangement between the National Center for Health Statistics (NCHS) and the Census Bureau. The methodology implemented by the Census Bureau to develop these county estimates is comparable to that used to produce national and state 1990-2000 intercensal estimates and is described on the Census Bureau's website (National Center for Health Statistics 2003). In 2007, the U.S. Census revised their methodology for population estimation. This revision impacts all population denominators for 2010 to 2014. The main effect of this approach was on specific age groups, particularly, two of the oldest age groups, ages 65-74 and 75 and older. In many states, the new method results in a larger increase in the younger age category and fewer in the older age category than estimated in previous years. Subsequently, this revision contributed to a larger decrease in age-adjusted cancer mortality rates in some states than what was expected based on statistics from last year.

The NCI modifies the Census data for the population estimates for the state of Hawaii. The Epidemiology Program of the Hawaii Cancer Research Center has developed its own set of population estimates, based on sample survey data collected by the Hawaii Department of Health. This effort grew out of a concern that the native Hawaiian population had been vastly undercounted in previous censuses. The "Hawaii adjustment" to the Census Bureau estimates has the net result of reducing the estimated white population and increasing the Asian and Pacific Islander population for the state. The Census Bureau estimates for the total population, black population, and American Indian and Alaska Native populations in Hawaii are unaffected. Refer to the SEER *Cancer Statistics Review, 1975-2006* and its methodologies for specific documentation regarding modifications made by the NCI to the Census Bureau estimates (Horner 2009).

For **Canada**, Statistics Canada provided the estimates of the Canadian population for all Canadian provinces and territories, adjusted for census net under coverage (http://www5.statcan.gc.ca/cansim/a26?lang=eng&retrLang=eng&id=0510001&pattern=population+estimates&tabMode=dataTable&srchLan=-1&p1=1&p2=-1).

CANCER SITE CODING, 2010-2014

Incidence Data. All cancer registries use the International Classification of Diseases for Oncology, third edition (ICD-O-3) to code the anatomic site and morphology. Cancer incidence statistics include invasive cancers only, with the exception of *in situ* cancer of the bladder. Although tables include incidence statistics for breast cancer *in situ* and benign tumors of the brain and central nervous system, these cases are not included in any counts or rates of total cancer incidence.

The SEER program site recode groups were used for classifying types of cancer, using anatomic site and morphology. Using this standard ontology, only squamous and basal cell carcinomas of the lip and genital organs are included in the data reported. All categories used to present pediatric cancer rates in Volume One are based on the International Classification of Childhood Cancer, third edition (ICCC-3) (Steliarova-Foucher 2005).

Summary tables of all codes and site groups for incidence can be found in the electronic copy of the CiNA Monograph Appendices on the NAACCR website, https://www.naaccr.org/cancer-in-north-america-cina-volumes/.

Mortality Data. Underlying cause of death was coded using the International Classification of Diseases version 10 (ICD-10). Cancer deaths were defined as those coded C00 through C97 in ICD-10.

The SEER mortality recode scheme was used to classify cancer deaths into the groupings used in the volume III (see Appendix B on the NAACCR website, https://www.naaccr.org/cancer-in-north-america-cina-volumes/).

Hispanic/Latino Ethnicity Identification. The ethnicity available in medical records and reported to cancer registries is enhanced by the use of the NAACCR Hispanic Identification Algorithm, version 2 (NHIAv2). NHIAv2 uses a combination of NAACCR variables including direct identification of ethnicity from the Spanish/Hispanic Origin variable (NAACCR data element 190 values 1-6), and information indirectly derived based on an evaluation of the strength of the birthplace, race, and surname (including maiden name when available) associations with Hispanic ethnicity status. After applying NHIAv2, cases not ultimately classified as Hispanic are classified as non-Hispanic, leaving no cases with "unknown" Hispanic status. The NHIAv2 algorithm allows the user to run the program using one of three possible options:

1) All Records: Surname portion is run on 0 (non-Hispanic), 7 (surname only), and 9 (unknown) for all counties. After the algorithm, item 190 codes (0, 7, 9) are either NHIA - Hispanic or non-Hispanic.

2) Option 1: (The default) For counties with 5% or more Hispanics - as for all records option. For counties with < than 5% Hispanic, surname portion is run on 7 (surname only) and 9 (unknown). After the algorithm, item 190 codes (7, 9) are either NHIA - Hispanic or non-Hispanic.

3) Option 2: For counties with 5% or more Hispanics - as for all records option. For counties with < than 5% Hispanic, surname portion is run on 7 (surname only). These become either NHIA - Hispanic or non-Hispanic. All cases coded to 9 on item 190 are converted to NHIA - non-Hispanic.

A comparison of the direct and indirect categorization of Hispanic ethnicity is provided in the Appendix F available from the NAACCR Website, https://www.naaccr.org/cancer-in-north-america-cina-volumes/.

The NHIAv2 method is described in detail elsewhere. (NAACCR Race and Ethnicity Work Group, 2011; NAACCR Guideline for Enhancing Hispanic-Latino Identification: Revised NAACCR Hispanic/Latino Identification Algorithm [NHIA v2.2.1], Revised September 12, 2011.)
[URL: https://20tqtx36s1la18rvn82wcmpn-wpengine.netdna-ssl.com/wp-content/uploads/2016/11/NHIA_v2_2_1_09122011.pdf Last Accessed, May 11, 2017]

Asian/Pacific Islander Identification. The information on Asian and Pacific Islander populations is also enhanced by submitting registries using a NAACCR developed algorithm (NAACCR, 2010). The NAACCR Asian Pacific Islander Identification Algorithm version 1 (NAPIIA v1.2.1) uses a combination of NAACCR variables to classify cases directly or indirectly as Asian/Pacific Islander for analytic purposes. It is focused on coding cases with a race code of Asian NOS (race code 96) or Pacific Islander NOS (race code 97) to a more

specific Asian or Pacific Islander race category, using the birthplace and name fields (first, last, and maiden names). Birthplace can be used to indirectly assign a specific race to one of eight Asian groups (Chinese, Japanese, Vietnamese, Korean, Asian Indian, Filipino, Thai, and Cambodian) and three Pacific Islander groups (Samoan, Micronesian, and Polynesian). Names can be used to indirectly assign a specific race to one of seven Asian groups (Chinese, Japanese, Vietnamese, Korean, Asian Indian, Filipino, and Hmong) and three Pacific Islander groups (Hawaiian, Guamanian, and Samoan). The algorithm uses the following NAACCR standard variables: Race 1 through Race 5 (NAACCR Items 160 through 164), Name – Last (NAACCR Item 2230), Name – First (NAACCR Item 2240), Name – Maiden (NAACCR Item 2390), Birthplace (NAACCR Item 250), Sex (NAACCR Item 220).

Misclassification of American Indian and Alaskan Native Populations. The collection of accurate information on the cancer incidence in American Indian and Alaskan Native populations is hampered by misclassification of this population within cancer case reports. In order to address this problem, cancer registries in the United States coordinate with the Centers for Disease Control and the Bureau of Indian Affairs to link their case files to the enrollment files of the Indian Health Service (Espey, et al., 2008). This process identifies significant numbers of American Indians and Alaskan Natives that were otherwise classified within the cancer case report. The improved information on this population enables calculation of much more accurate cancer incidence rates. As presented in the incidence volumes, the data on American Indians and Alaskan natives are restricted to those counties where Indian Health Service Clinics are located, namely Contract Health Service Delivery Area (CHSDA) counties. This restriction assures that the incidence data presented is the best available data for this population.

DATA QUALITY INDICATORS

NAACCR assesses the quality of cancer incidence data from individual registries for a number of data quality indicators, which are described in detail below. Results for these indicators can be found in the *Cancer in North America, 2010-2014* Appendices on the NAACCR website. A dash in the tables indicates that data were not submitted for the year or were not able to be calculated.

In order to be included in the NAACCR Combined rates presented in Volume One, the data had to meet the criteria for high quality incidence data. These criteria were applied to each year of data individually, except for the estimate of duplicate reports, which was calculated for the years 2010 to 2014 as a whole.

- Data for 2010 through 2014 had to be submitted to NAACCR by December 1, 2016.
- The estimate of duplicate case reports had to be less than 2 duplicate reports per 1000 cases.
- All cases had to pass the edits on variables needed to compute cancer incidence rates by site, sex, race, and tumor descriptions on the NAACCR Call for Data EDITS metafile.
- Fewer than 3% of cases had missing information for sex, county of residence at diagnosis, and age at diagnosis, and fewer than 5% of the cases had an unknown race.
- Death clearance must be completed for 2010 through 2014 mortality with the percent of all cases derived from death certificates only (DCOs) of less than 5%.
- The NAACCR method to estimate completeness of case ascertainment yielded an estimate of 90% or higher.
- All cases had to pass the inter-record edits on cases reporting multiple primary cancers, as determined by the standard Inter-record EDITS program developed for the NAACCR Call for Data submissions.

To be included in the combined stage data, the registry must meet the requirements for being included in the combined rates (above), have submitted stage data for all years 2010 through 2014 and provided the derived summary stage field in their submission.

To be included in the delay adjusted data, the registry must meet the requirements for being included in the combined rates, have submitted data for all years 2010 through 2014 and have provided the data needed to calculate the delay factors.

The selection criteria for including registry data in Volume Four on cancer survival included meeting the criteria for inclusion in the combined incidence rates and providing the follow up fields required for the survival calculations. In addition, the data either needed to meet the SEER standards for follow-up or the registry needed

to have ascertained deaths through the study cutoff date (December 31, 2013). To meet the SEER standard for follow-up, a minimum of 90% of patients needed to have follow-up dates on or after January 1, 2014, or be deceased. For U.S. registries that did not meet the SEER standard for follow-up, it was necessary to conduct state death linkages and linkage with the National Death Index. For Canadian registries, it was necessary to conduct death linkages within the province or territory. Registry data also were required to have no irregularities in the variables that are key to the survival calculations.

Duplicate Case Records. Most central cancer registries rely on multiple reporting sources for cancer case reports. At the central cancer registry, multiple reports for the same patient must then be matched and the information from all records consolidated. In addition to determining whether multiple reports refer to the same individual, central cancer registry staff must also determine whether the tumor represents a new primary tumor or a duplicate report for a tumor already recorded. Failure to eliminate duplicate cases and duplicate tumors results in over-counting cancer cases. As a part of routine cancer registry operations, a variety of tools are used to ensure accurate case linkage and case consolidation. As part of the preparation of the data submission to NAACCR, each registry uses the NAACCR protocol to determine the adequacy of case linkage and consolidation operations in identifying duplicate records.

The NAACCR protocol for assessing duplicate cases can be found on the NAACCR website, http://www.naaccr.org.

Completeness of Case Ascertainment. In order to evaluate case completeness for all geographic areas included in this monograph, the NAACCR Method to Estimate Completeness is used. The method is described in detail elsewhere (Wu, et al., 2002; Howe, 2007). A data analysis tool is available on the NAACCR website, https://www.naaccr.org/analysis-and-data-improvement-tools/#COMPLETENESS that documents and calculates sex-, race-, and site- specific estimates based on observed incidence counts and rates, observed local cancer death rates, and a standard rate ratio of incidence to mortality. In addition, adjustments are made to the calculations to account for some variation in sex-, race- and site-specific variation in case fatality. The method assumes that the relationship between incidence rates and mortality rates are stable. However, with the recent annual decreases in cancer death rates, and the awareness that the decreases are not the same in all parts of the United States and Canada, it appears that this assumption is being challenged with a noticeable bias in an upward migration of completeness estimates for many locales. The completeness estimates provided in this publication used mortality data for the years 2010-2014 to determine completeness in U.S. registries and Canadian registries. The percent completeness presented for each registry and used to determine inclusion of individual registry data in the combined rates presented in Volume One were based upon rate ratios of 2010-2014 SEER 11 incidence rates divided by 2010-2014 U.S. mortality rates.

Missing Case Information. NAACCR has developed standards for completeness of data on key data items that are needed to produce meaningful cancer incidence statistics. These key data items include race, sex, county of residence at diagnosis, and age at diagnosis. Cases with unknown sex or age are omitted from all calculations. Cases with unknown race and county of residence were included in the cancer counts and rates for all races and/or all counties combined.

Death Certificate Only Cases. The proportion of cases identified by death certificate only (DCO) is an indicator of data quality and completeness. Central cancer registries use death certificates to identify potentially missed cases and to conduct follow back on cases that have cancer on the death certificate but who are not incident cases in the registry. Cases without follow back information are considered to be DCO cases and may have incomplete or missing information, including date of diagnosis and stage at diagnosis. For DCO cases, the date of death is used as the date of diagnosis. Registries that did not use death certificates as a source of case ascertainment have "na" listed in the death certificate only row on the registry description page.

Data Reliability and Accuracy. All data submitted to NAACCR are evaluated for reliability and accuracy using the EDITS program and a specific set of edits incorporated into a Call for Data Edits Metafile. An edit reviews the internal consistency between and among data elements, such as anatomic site and morphology. Cases that are identified as having errors are reviewed by registry staff and resolved prior to their NAACCR submission. The NAACCR Call for Data EDITS metafile is available on the NAACCR website.
[URL http://www.naaccr.org/DataandPublications/CallforData.aspx, Last accessed April 24, 2017].

An inter-record edits program, developed by the Centers for Disease Control and Prevention, is used to identify errors among the records reported for persons who have had multiple tumors diagnosed. For example, an inter-record edit will identify whether birthplace is the same on all reports of multiple tumors that are reported for one individual.

Timeliness of Data. The NAACCR standard defines timely data as data that are available for use in incidence statistics within 23 months of the close of a diagnostic year (i.e., December 1, 2016 for all cases diagnosed in 2014).

Site-Specific Microscopic Confirmation. This criterion is not used by NAACCR to determine high quality for the purposes of this publication. However, it is a useful indicator of quality of data collection. A proportion of microscopically confirmed cases that is too high or low may suggest problems in case ascertainment or abstracting. However, this proportion varies by cancer site. For sites that are more likely to rely on a clinical or radiological diagnosis, e.g., cancers of the pancreas and brain, confirmation rates that are too high may suggest that some clinically diagnosed cases are missing. Also, registries that do not use death certificates for case finding have an artificially high proportion of microscopically confirmed cases, because DCO cases by definition do not have information on whether the tumor was microscopically confirmed. While no NAACCR standard has been determined for microscopic confirmation, the guideline we use is that the proportion should fall between 92 and 96 percent of all cancer cases, based on the experience of the SEER program.

CALCULATION OF STATISTICS

Rates. Rates are per 100,000 population and are age-adjusted by five-year age groups to the 2000 U.S. standard population based on single years of age (Ries, et al. 2005), the 1996 Canadian population standard (Statistics Canada, 2003), and to the World population standard (Parkin, et al., 2002). Rates for children and adolescents in Volume One were expressed per million population. The incidence and death rates in this monograph are annual averages for the period 2010 through 2014. The age distributions of the three population standards are as follows:

AGE GROUP	2000 U.S.	1996 CDN.	WORLD
00 years	3,794,901	12,342	24,000
01-04 years	15,191,619	53,893	96,000
05-09 years	19,919,840	67,985	100,000
10-14 years	20,056,779	67,716	90,000
15-19 years	19,819,518	67,841	90,000
20-24 years	18,257,225	67,761	80,000
25-29 years	17,722,067	72,914	80,000
30-34 years	19,511,370	87,030	60,000
35-39 years	22,179,956	88,510	60,000
40-44 years	22,479,229	80,055	60,000
45-49 years	19,805,793	71,847	60,000
50-54 years	17,224,359	55,812	50,000
55-59 years	13,307,234	44,869	40,000
60-64 years	10,654,272	40,705	40,000
65-69 years	9,409,940	37,858	30,000
70-74 years	8,725,574	32,589	20,000
75-79 years	7,414,559	23,232	10,000
80-84 years	4,900,234	15,424	5,000
85+ years	4,259,173	11,617	5,000
Total	**274,633,642**	**1,000,000**	**1,000,000**

Standard Errors. Standard errors (S.E.) of the rates were calculated using the formula:

$$S.E. = \sqrt{\sum \frac{w_j^2 n_j}{p_j^2}}$$

where w_j = the fraction of the standard population in age group j, n_j = number of cases or deaths in that age group, and p_j = person years denominator (Breslow and Day, 1987). For many registries, the standard errors of the rates are small, as the population covered is large. However, for registries that cover a small population, the standard error may be substantial.

Confidence Intervals. Although not reproduced in the printed monograph, bar charts of the registry-specific, age-adjusted rates with 95 percent confidence intervals (Tiwari, et al., 2006) are available on the NAACCR website (https://www.naaccr.org/cancer-in-north-america-cina-volumes/). The confidence intervals allow the user to assess the precision of the estimate and is an approximate and conservative indicator of whether a registry's rate is statistically higher or lower than the rate for the combined United States, based on whether or not the upper or lower limit of the confidence interval overlaps the 95% confidence interval for the United States.

Comparison of Rates. In addition to true regional variation in cancer risk, differences in cancer incidence or mortality rates between areas may be due to either differences in the demographic makeup of the population or differences in data quality. In making valid comparisons of cancer incidence rates among registries, it is important to review the data quality indicators for each registry and consider differences in the racial composition of the populations being compared before conclusions are drawn about variations in regional rates. Interpretation without consideration of these factors may contribute to misleading or inaccurate conclusions.

The standard error of adjusted rates can be used to evaluate the statistical significance of rate differences among comparable populations. For example, if the adjusted rates in two populations are R_1 and R_2 and their standard errors are $S.E._1$ and $S.E._2$, an approximate confidence interval for the rate ratio can be calculated using the following formula:

$$(R_1/R_2)^{1 \pm z/x}$$

where $x = (R_1 \text{ B } R_2) / \sqrt{(S.E._1^2 + S.E._2^2)}$ and z = 1.96 for 95% limits (Parkin, et al., 1992). If this interval does not include one, the two rates are statistically significantly different at a p value of 0.05. This test can be inaccurate for rates based on fewer than 16 cases or deaths, and it should not be used for rates based on fewer than six cases or deaths. It should be emphasized that this kind of comparison of adjusted rates must be undertaken with caution as misleading conclusions may be drawn if the ratios of the age specific rates in the two populations are not constant in all age groups. In these circumstances, the ratios of the adjusted rates will vary according to the standard populations used (Esteve, et al., 1994).

Cell Suppression. Counts and rates were suppressed (indicated by "-") in the tables if the race, gender, and site-specific number of cases for incidence was less than six or when the number of deaths was less than ten. Complementary cell suppression was employed, when necessary, to prevent the contents of a single suppressed cell from being calculated. These counts are included in the calculation of all sites combined. A dash is also used to indicate not applicable, as in the gender specific cancers. If the rate was less than 0.05 per 100,000 then the rate is listed as 0.0.

REFERENCES

Anderson RN, Minino AM, Hoyert DL, Rosenberg HM. Comparability of cause of death between ICD-9 and ICD-10: Preliminary estimates. *National Vital Statistics Reports 2001*;49:1-32.

Breslow NE and Day NE. Statistical Methods in Cancer Research. Vol. II. 1987. International Agency for Research on Cancer: Lyon, France.

Clegg LX, Feuer EJ, Midthune D, Fay MP, Hankey BF. Impact of reporting delay and reporting error on cancer incidence rates and trends. J Natl Cancer Inst 2002;94:1537-45.

Espey DK, Wiggins CL, Jim MA, Miller BA, Johnson CJ, Becker TM. Methods for improving cancer surveillance data in American Indian and Alaska Native populations. Cancer. 2008 Sep 1;113(5 Suppl):1120-30.

Espey DK, Wu XC, Swan J, Wiggins C, Jim MA, Ward E, Wingo PA, Howe HL, Ries LA, Miller BA, Jemal A, Ahmed F, Cobb N, Kaur JS, Edwards BK. Annual report to the nation on the status of cancer, 1975-2004, featuring cancer in American Indians and Alaska Natives. Cancer. 2007 Nov 15;110(10):2119-52.

Esteve J, Benhamou E, and Raymond L. Statistical Methods in Cancer Research. Vol. V. 1994. International Agency for Research on Cancer: Lyon, France.

Horner MJ, Ries LAG, Krapcho M, Neyman N, Aminou R, Howlader N, Altekruse SF, Feuer EJ, Huang L, Mariotto A, Miller BA, Lewis DR, Eisner MP, Stinchcomb DG, Edwards BK (eds). SEER Cancer Statistics Review, 1975-2006, National Cancer Institute. Bethesda, MD
[URL: http://www.seer.cancer.gov/csr/1975_2006/, based on November 2008 SEER data submission, posted to the SEER web site, 2009.]

Howe HL, Carozza S, O'Malley C, Dolecek TA, Finch JL, Kohler B, Wet D, Liu L, Schymura MJ, Williams M, Abe T, Agovino P, Chen VW, Firth R, Harkins D, Hotes, J, Lake A, Roney D, Suarez L (eds). Cancer in U.S. Hispanics/Latinos, 1995-2000. Springfield (IL): North American Association of Central Cancer Registries, December 2003.

Johnson CH, Peace S, Adamo P, Fritz A, Percy-Laurry A, Edwards BK. The 2007 Multiple Primary and Histology Coding Rules. National Cancer Institute, Surveillance, Epidemiology and End Results Program. Bethesda, MD, 2007.

Midthune DN, Fay MP, Clegg LX, Feuer EJ. Modeling reporting delays and reporting corrections in cancer registry data. J Am Stat Assoc 2005;100(469):61-70.

NAACCR Expert Panel in Hispanic Identification. *Report of the NAACCR Expert Panel on Hispanic Identification 2003*. Springfield (IL): North American Association of Central Cancer Registries, October 2003.

NAACCR Latino Research Work Group. *NAACCR Guideline for Enhancing Hispanic/Latino Identification: Revised NAACCR Hispanic/Latino Identification Algorithm [NHIA v2]*. Springfield (IL): North American Association of Central Cancer Registries. September 2005.
[URL:https://20tqtx36s1la18rvn82wcmpn-wpengine.netdna-ssl.com/wp-content/uploads/2016/11/NHIA_v2_2_1_09122011.pdfNAACCR Method to Estimate Completeness. A Data Analysis Tool for Calculations.
[URL: https://www.naaccr.org/analysis-and-data-improvement-tools/#COMPLETENESS Last accessed May 11, 2017].

NAACCR Race and Ethnicity Work Group. *NAACCR Asian Pacific Islander Identification Algorithm [NAPIIA v1.2.1]*. Springfield, IL: North American Association of Central Cancer Registries, September 2010.

National Center for Health Statistics. U.S. Census Populations with Bridged Race Categories. September 2003. [URL: http://www.cdc.gov/nchs/nvss/bridged_race.htm. Last Accessed, April 24, 2017].

Parkin D.M., Whelan S.L., Ferlay J., Teppo L., Thomas D., Thomas D.B. Cancer Incidence in Five Continents. Vol. VIII. 2002. International Agency for Research on Cancer: Lyon, France.

SEER. 2000 U.S. Standard Population vs. Standard Million. 13 December 2004. http://seer.cancer.gov/stdpopulations/single_age.html
Steliarova-Foucher E, Stiller C, Lacour B, Kaatsch P. International Classification of Childhood Cancer, third edition. Cancer 2005;103:1457-67.

Tiwari RC, Clegg LX, Zou Z. Efficient interval estimation for age-adjusted cancer rates. Stat Methods Med Res 2006;15:547-569.

Weir HK, Johnson CJ, Mariotto AB, Turner D, Wilson RJ, Nishri D, Ward KC. Evaluation of North American Association of Central Cancer Registries' (NAACCR) data for use in population-based cancer survival studies. J Natl Cancer Inst Monogr. 2014 Nov;2014(49):198-209.

Wu XC, Hotes J, Fulton JP, Cormier M, Correa C, McLaughlin CC, Kosary C, Howe HL, Chen VW (Eds.) Cancer in North America, 1996-2000, Volume One: Introduction and Technical Notes, Incidence. April 2002. [URL: https://www.naaccr.org/previous-volumes-of-cina/ Last accessed May 11, 2017].

Zou J, Huang L, Midthune D, Horner MJ, Krapcho M, Feuer EJ. Effect of reporting year on delay modeling. Statistical Research and Applications Branch, National Cancer Institute; 2009. Technical Report #2009-01
Clegg LX, Feuer EJ, Midthune D, Fay MP, Hankey BF. Impact of reporting delay and reporting error on cancer incidence rates and trends. J Natl Cancer Inst 2002;94:1537-45. [Full Text]

Midthune DN, Fay MP, Clegg LX, Feuer EJ. Modeling reporting delays and reporting corrections in cancer registry data. J Am Stat Assoc 2005;100(469):61-70. [View Abstract]

Zou J, Huang L, Midthune D, Horner MJ, Krapcho M, Feuer EJ. Effect of reporting year on delay modeling. Statistical Research and Applications Branch, National Cancer Institute; 2009. Technical Report #2009-01. Available from: http://surveillance.cancer.gov/reports/

Section II

Top Five Most Commonly Diagnosed Cancers in North America

Top Five Most Commonly Diagnosed Cancers[1] in North America
by Sex, All Races
2010-2014

Males/All Races

United States

Cancer Sites	Count	Percent
All Sites	3,894,209	100.0
Prostate	943,855	24.2
Lung & Bronchus	555,126	14.3
Colon & Rectum	350,832	9.0
Bladder	262,379	6.7
Melanomas of the Skin	201,442	5.2

Canada

Cancer Sites	Count	Percent
All Sites	176,555	100.0
Prostate	42,600	24.1
Colon & Rectum	24,708	14.0
Lung & Bronchus	22,999	13.0
Bladder	12,665	7.2
Non-Hodgkin Lymphoma	8,025	4.5

North America

Cancer Sites	Count	Percent
All Sites	4,070,764	100.0
Prostate	986,455	24.2
Lung & Bronchus	578,125	14.2
Colon & Rectum	375,540	9.2
Bladder	275,044	6.8
Melanomas of the Skin	208,393	5.1

Females/All Races

United States

Cancer Sites	Count	Percent
All Sites	3,781,169	100.0
Breast	1,107,649	29.3
Lung & Bronchus	493,029	13.0
Colon & Rectum	323,272	8.5
Uterus Corpus & Uterus NOS	241,508	6.4
Thyroid	169,974	4.5

Canada

Cancer Sites	Count	Percent
All Sites	164,882	100.0
Breast	46,172	28.0
Lung & Bronchus	22,022	13.4
Colon & Rectum	19,829	12.0
Uterus Corpus & Uterus NOS	10,414	6.3
Non-Hodgkin Lymphoma	6,448	3.9

North America

Cancer Sites	Count	Percent
All Sites	3,946,051	100.0
Breast	1,153,821	29.2
Lung & Bronchus	515,051	13.1
Colon & Rectum	343,101	8.7
Uterus Corpus & Uterus NOS	251,922	6.4
Thyroid	175,611	4.5

[1] Compiled by the North American Association of Central Cancer Registries, based on cases included in the combined incidence for North America.

Top Five Most Commonly Diagnosed Cancers[1] in the U.S. by Race/Ethnicity, Males 2010-2014

Hispanic/Latino, All Races

Cancer Sites	Count	Percent
All Sites	267,627	100.0
Prostate	63,511	23.7
Colon & Rectum	28,612	10.7
Lung & Bronchus	23,424	8.8
Kidney & Renal Pelvis	15,570	5.8
Liver & Intrahepatic Bile Duct	14,558	5.4

White

Cancer Sites	Count	Percent
All Sites	3,265,680	100.0
Prostate	735,242	22.5
Lung & Bronchus	472,759	14.5
Colon & Rectum	289,746	8.9
Bladder	240,261	7.4
Melanomas of the Skin	190,947	5.8

Black

Cancer Sites	Count	Percent
All Sites	432,925	100.0
Prostate	150,352	34.7
Lung & Bronchus	62,845	14.5
Colon & Rectum	42,108	9.7
Kidney & Renal Pelvis	19,982	4.6
Liver & Intrahepatic Bile Duct	15,442	3.6

American Indian/Alaskan Native[2]

Cancer Sites	Count	Percent
All Sites	12,813	100.0
Prostate	2,311	18.0
Lung & Bronchus	1,911	14.9
Colon & Rectum	1,513	11.8
Kidney & Renal Pelvis	994	7.8
Liver & Intrahepatic Bile Duct	687	5.4

Chinese

Cancer Sites	Count	Percent
All Sites	20,869	100.0
Lung & Bronchus	3,508	16.8
Prostate	3,424	16.4
Colon & Rectum	2,729	13.1
Liver & Intrahepatic Bile Duct	1,552	7.4
Stomach	1,195	5.7

Japanese

Cancer Sites	Count	Percent
All Sites	7,781	100.0
Prostate	1,653	21.2
Colon & Rectum	1,178	15.1
Lung & Bronchus	983	12.6
Bladder	486	6.2
Stomach	409	5.3

Filipino

Cancer Sites	Count	Percent
All Sites	15,402	100.0
Prostate	3,698	24.0
Lung & Bronchus	2,553	16.6
Colon & Rectum	1,855	12.0
Non-Hodgkin Lymphoma	850	5.5
Liver & Intrahepatic Bile Duct	772	5.0

Vietnamese

Cancer Sites	Count	Percent
All Sites	9,167	100.0
Lung & Bronchus	1,709	18.6
Liver & Intrahepatic Bile Duct	1,454	15.9
Colon & Rectum	1,163	12.7
Prostate	1,053	11.5
Non-Hodgkin Lymphoma	490	5.3

Korean

Cancer Sites	Count	Percent
All Sites	7,238	100.0
Colon & Rectum	1,037	14.3
Lung & Bronchus	1,036	14.3
Prostate	881	12.2
Stomach	841	11.6
Liver & Intrahepatic Bile Duct	597	8.2

[1] Compiled by the North American Association of Central Cancer Registries, based on cases included in the combined incidence for the United States.

[2] Counts and percentages are based upon cancer cases diagnosed among the AI/AN population in CHSDA counties. Statistics exclude data for KS for all years. See Introduction for more information.

Top Five Most Commonly Diagnosed Cancers[1] in the U.S. by Race/Ethnicity, Females 2010-2014

Hispanic/Latino, All Races

Cancer Sites	Count	Percent
All Sites	292,872	100.0
Breast	83,873	28.6
Colon & Rectum	24,205	8.3
Thyroid	21,544	7.4
Uterus Corpus & Uterus NOS	20,579	7.0
Lung and Bronchus	19,375	6.6

White

Cancer Sites	Count	Percent
All Sites	3,168,903	100.0
Breast	919,063	29.0
Lung & Bronchus	426,857	13.5
Colon & Rectum	264,042	8.3
Uterus Corpus & Uterus NOS	201,669	6.4
Thyroid	140,681	4.4

Black

Cancer Sites	Count	Percent
All Sites	418,011	100.0
Breast	130,551	31.2
Lung & Bronchus	50,098	12.0
Colon & Rectum	42,301	10.1
Uterus Corpus & Uterus NOS	27,125	6.5
Thyroid	14,716	3.5

American Indian/Alaskan Native[2]

Cancer Sites	Count	Percent
All Sites	14,277	100.0
Breast	3,839	26.9
Lung & Bronchus	1,875	13.1
Colon & Rectum	1,463	10.2
Uterus Corpus & Uterus NOS	896	6.3
Kidney & Renal Pelvis	670	4.7

Chinese

Cancer Sites	Count	Percent
All Sites	23,512	100.0
Breast	7,129	30.3
Lung & Bronchus	2,831	12.0
Colon & Rectum	2,313	9.8
Thyroid	1,745	7.4
Uterus Corpus & Uterus NOS	1,314	5.6

Japanese

Cancer Sites	Count	Percent
All Sites	11,228	100.0
Breast	3,684	32.8
Colon & Rectum	1,318	11.7
Lung & Bronchus	1,287	11.5
Uterus Corpus & Uterus NOS	583	5.2
Pancreas	528	4.7

Filipino

Cancer Sites	Count	Percent
All Sites	22,652	100.0
Breast	8,202	36.2
Lung & Bronchus	1,945	8.6
Colon & Rectum	1,860	8.2
Uterus Corpus & Uterus NOS	1,810	8.0
Thyroid	1,787	7.9

Vietnamese

Cancer Sites	Count	Percent
All Sites	8,730	100.0
Breast	2,398	27.5
Lung & Bronchus	996	11.4
Colon & Rectum	915	10.5
Thyroid	590	6.8
Liver & Intrahepatic Bile Duct	444	5.1

Korean

Cancer Sites	Count	Percent
All Sites	9,235	100.0
Breast	2,631	28.5
Colon & Rectum	1,026	11.1
Lung & Bronchus	916	9.9
Thyroid	764	8.3
Stomach	554	6.0

[1] Compiled by the North American Association of Central Cancer Registries, based on cases included in the combined incidence for the United States.

[2] Counts and percentages are based upon cancer cases diagnosed among the AI/AN population in CHSDA counties. Statistics exclude data for KS for all years. See Introduction for more information.

Section III

Average-annual Age-adjusted NAACCR Combined Cancer Incidence Rates for the United States, Canada and North America

Race-Specific Population Counts by Year[1]

Race	2010	2011	2012	2013	2014
All Races	299,267,779	301,573,693	303,882,514	306,129,598	308,526,428
White	237,094,784	238,318,010	239,517,600	240,627,110	241,822,462
Black	41,571,351	42,077,600	42,585,600	43,086,124	43,608,506
American Indian/Alaskan Native (AI/AN)	3,924,008	3,978,135	4,034,826	4,088,627	4,147,709
Asian/Pacific Islander (API)	16,677,636	17,199,948	17,744,488	18,327,737	18,947,751

All Races Population 2010-2014 1,519,380,012
Distribution of Population by Race

Hispanic/Latino Population Counts by Year[1]

Ethnicity	2010	2011	2012	2013	2014
Hispanic/Latino, All Races	48,823,013	49,987,124	51,074,401	52,170,009	53,326,735
Non-Hispanic White	193,333,519	193,529,098	193,781,823	193,937,678	194,125,809
Non-Hispanic Black	38,977,483	39,402,283	39,827,659	40,243,595	40,676,408

Hispanic/Latino Population, Census 2010: 48,557,432
Distribution of Hispanic/Latino Populations[2]

% Hispanic/Latino of Total Population
By Census Year[2]

[1] Source: NCI SEER Program, U.S. Population Data
[2] Source: U.S. Census Bureau, 2010 Census

Race-Specific Population Indicators (2010-2014)

Combined Population:	
All Races	1,519,380,012
White	1,197,379,966
Black	212,929,181
American Indian/Alaskan Native	20,173,305
Asian/Pacific Islander	88,897,560
Hispanic/All Races	255,381,282
Non-Hispanic White	968,707,927
Non-Hispanic Black	199,127,428

Percentage of U.S. Population Reflected in the Combined Cancer Statistics:	
All Races	96.7%
White	96.5%
Black	98.5%
American Indian/Alaskan Native	91.8%
Asian/Pacific Islander	97.0%
Hispanic/All Races	96.2%
Non-Hispanic White	96.6%
Non-Hispanic Black	98.5%

Number of Cases in Combined Cancer Statistics:	
All Races	7,675,378
White	6,434,583
Black	850,948
American Indian/Alaskan Native	39,645
Asian/Pacific Islander	230,920
Hispanic/All Races	560,499
Non-Hispanic White	5,917,355
Non-Hispanic Black	833,513

Data Quality Indicators (2010-2014)

Indicators (%)	2010	2011	2012	2013	2014
Missing Race	0.9	1.0	1.0	1.1	1.3
Missing Sex	0.0	0.0	0.0	0.0	0.0
Missing County	0.0	0.0	0.0	0.0	0.0
Missing Age	0.0	0.0	0.0	0.0	0.0
Death Certificate Only Cases	1.8	1.7	1.7	1.7	1.9

Site-Specific Microscopic Confirmation (2010-2014)

Cancer Site	% Micro. Confirm
All Sites	93.1
Oral Cavity & Pharynx	97.5
Esophagus	94.6
Stomach	96.2
Colon & Rectum	96.2
Liver	57.9
Pancreas	81.9
Lung & Bronchus	88.1
Melanomas of the Skin	99.1
Female Breast (excl. *in situ*)	98.4
Cervix uteri	97.1
Uterus Corpus & Uterus NOS[1]	98.5
Ovary	92.1

Cancer Site	% Micro. Confirm
Prostate	96.7
Testis	99.0
Bladder (incl. *in situ*)	97.8
Kidney & Renal Pelvis	89.8
Brain & Other Nervous System	84.7
Thyroid Gland	99.4
Hodgkin Lymphoma	98.3
Non-Hodgkin Lymphoma	96.3
Multiple Myeloma	88.4
Leukemias	88.7
Brain & ONS[1] *benign/borderline*	44.2
Breast *in situ*	100.0

[1] Abbreviations: NOS - Not Otherwise Specified; ONS - Other Nervous System

NAACCR U.S. Combined Incidence 2010-2014, All Races Combined

Total Cancer Cases and Average Annual Age-adjusted Cancer Incidence Rates*
NAACCR United States Combined, 2010-2014, All Races

	MALES (ALL RACES)					FEMALES (ALL RACES)				
SITES	*Count*	*Adj. Rate 2000 U.S.*	*S.E.*	*Adj. Rate 1996 Cdn*	*S.E.*	*Count*	*Adj. Rate 2000 U.S.*	*S.E.*	*Adj. Rate 1996 Cdn*	*S.E.*
All Sites (invasive & in situ bladder)	3,894,209	502.1	0.26	479.8	0.25	3,781,169	418.0	0.22	403.7	0.21
Oral Cavity & Pharynx	141,503	17.4	0.05	16.8	0.05	58,235	6.4	0.03	6.1	0.03
Lip	6,952	0.9	0.01	0.9	0.01	2,650	0.3	0.01	0.3	0.01
Tongue	43,558	5.3	0.03	5.1	0.02	17,207	1.9	0.01	1.8	0.01
Major salivary gland	12,352	1.7	0.02	1.6	0.01	8,676	1.0	0.01	1.0	0.01
Floor of mouth	6,735	0.8	0.01	0.8	0.01	3,042	0.3	0.01	0.3	0.01
Gum & other mouth	14,865	1.9	0.02	1.8	0.02	11,788	1.3	0.01	1.2	0.01
Nasopharynx	6,244	0.8	0.01	0.8	0.01	2,529	0.3	0.01	0.3	0.01
Oropharynx	7,038	0.8	0.01	0.8	0.01	2,192	0.2	0.01	0.2	0.00
Hypopharynx	9,028	1.1	0.01	1.1	0.01	2,149	0.2	0.01	0.2	0.00
Digestive System	756,357	97.6	0.11	92.5	0.11	609,032	65.3	0.09	61.6	0.08
Esophagus	62,815	8.0	0.03	7.6	0.03	16,881	1.8	0.01	1.7	0.01
Stomach	69,817	9.2	0.04	8.7	0.03	42,821	4.7	0.02	4.4	0.02
Small intestine	20,905	2.7	0.02	2.6	0.02	18,746	2.0	0.02	2.0	0.01
Colon excl. rectum	238,274	31.6	0.07	29.7	0.06	242,580	26.0	0.05	24.3	0.05
Rectum & rectosigmoid	112,558	14.3	0.04	13.7	0.04	80,692	8.8	0.03	8.5	0.03
Anus, anal canal & anorectum	11,666	1.5	0.01	1.4	0.01	20,005	2.2	0.02	2.1	0.02
Liver	92,538	11.0	0.04	10.6	0.04	31,101	3.3	0.02	3.1	0.02
Gallbladder	6,079	0.8	0.01	0.8	0.01	13,019	1.4	0.01	1.3	0.01
Pancreas	109,099	14.2	0.04	13.4	0.04	104,962	11.1	0.03	10.4	0.03
Respiratory System	612,819	80.3	0.10	75.9	0.10	511,256	54.8	0.08	52.2	0.07
Larynx	48,658	6.0	0.03	5.8	0.03	12,639	1.4	0.01	1.3	0.01
Lung & bronchus	555,126	73.0	0.10	69.0	0.09	493,029	52.8	0.08	50.3	0.07
Bones & Joints	8,221	1.1	0.01	1.1	0.01	6,572	0.8	0.01	0.8	0.01
Soft Tissue (incl. heart)	29,672	4.0	0.02	3.8	0.02	23,897	2.8	0.02	2.7	0.02
Skin (excl. basal & sq.)	217,979	28.9	0.06	27.4	0.06	152,026	17.5	0.05	17.2	0.05
Melanomas of the skin	201,442	26.6	0.06	25.3	0.06	141,306	16.3	0.04	16.1	0.04
Breast (excl. *in situ*)	9,909	1.3	0.01	1.2	0.01	1,107,649	123.6	0.12	120.4	0.12
Female Genital System	-	-	-	-	-	445,619	49.3	0.08	48.3	0.07
Cervix uteri	-	-	-	-	-	60,334	7.6	0.03	7.7	0.03
Uterus Corpus	-	-	-	-	-	233,530	25.0	0.05	24.5	0.05
Uterus, NOS[1]	-	-	-	-	-	7,978	0.9	0.01	0.8	0.01
Ovary	-	-	-	-	-	103,334	11.5	0.04	11.1	0.04
Male Genital System	992,481	121.6	0.12	119.1	0.12	-	-	-	-	-
Prostate	943,855	115.0	0.12	112.2	0.12	-	-	-	-	-
Testis	40,376	5.5	0.03	5.9	0.03	-	-	-	-	-
Penis	6,501	0.9	0.01	0.8	0.01	-	-	-	-	-
Urinary System	443,681	59.0	0.09	55.4	0.08	191,265	20.7	0.05	19.7	0.05
Bladder (incl. *in situ*)	262,379	35.8	0.07	33.1	0.07	83,602	8.8	0.03	8.3	0.03
Kidney & renal pelvis	171,900	21.8	0.05	21.1	0.05	102,450	11.3	0.04	11.0	0.04
Ureter	5,942	0.8	0.01	0.8	0.01	3,579	0.4	0.01	0.4	0.01
Eye & Orbit	7,406	1.0	0.01	0.9	0.01	6,267	0.7	0.01	0.7	0.01
Brain & Other Nervous System	59,227	7.7	0.03	7.6	0.03	47,419	5.6	0.03	5.4	0.03
Brain	55,970	7.3	0.03	7.1	0.03	43,939	5.1	0.02	5.0	0.02
Endocrine System	61,840	8.0	0.03	8.0	0.03	175,634	21.8	0.05	22.2	0.05
Thyroid gland	55,668	7.2	0.03	7.2	0.03	169,974	21.1	0.05	21.5	0.05
Lymphomas	196,106	26.0	0.06	24.9	0.06	162,518	18.1	0.05	17.5	0.04
Hodgkin Lymphoma	22,955	3.1	0.02	3.1	0.02	18,446	2.4	0.02	2.4	0.02
Non-Hodgkin Lymphoma	173,151	22.9	0.06	21.8	0.05	144,072	15.8	0.04	15.1	0.04
Multiple Myeloma	61,561	8.1	0.03	7.6	0.03	49,548	5.3	0.02	5.0	0.02
Leukemias	130,643	17.6	0.05	16.6	0.05	95,521	10.7	0.04	10.2	0.03
Acute lymphocytic	13,417	1.8	0.02	1.8	0.02	10,505	1.4	0.01	1.4	0.01
Chronic lymphocytic	46,590	6.2	0.03	5.8	0.03	30,523	3.2	0.02	3.0	0.02
Acute myeloid	38,098	5.2	0.03	4.9	0.03	31,157	3.5	0.02	3.4	0.02
Chronic myeloid	16,765	2.3	0.02	2.2	0.02	12,390	1.4	0.01	1.3	0.01
Other leukemias	15,773	2.2	0.02	2.0	0.02	10,946	1.2	0.01	1.1	0.01
Mesothelioma	12,045	1.7	0.02	1.6	0.01	3,808	0.4	0.01	0.4	0.01
Kaposi Sarcoma	5,042	0.7	0.01	0.7	0.01	651	0.1	0.00	0.1	0.00
Ill-defined & Unknown	147,717	20.3	0.05	18.6	0.05	134,252	14.2	0.04	13.1	0.04
Selected tumors (non-invasive)										
Brain & ONS[1] (*benign/borderline*)	63,464	8.4	0.03	8.1	0.03	129,269	14.6	0.04	14.2	0.04
Breast (*in situ*)	1,122	0.1	0.00	0.1	0.00	271,007	30.6	0.06	29.9	0.06

* Rates are per 100,000 population and are age-adjusted by five-year age groups to the 2000 U.S. standard population based on single years of age and to the 1996 Canadian population standard.

- Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for All Sites combined.

[1] Abbreviations: NOS - Not Otherwise Specified; ONS - Other Nervous System

Total Cancer Cases and Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates[1] for the Five Most Commonly Diagnosed Cancers
Selected Areas in the NAACCR United States Combined[2], 2010-2014, All Races, Males

	All Sites			Prostate			Lung & Bronchus			Colon & Rectum			Bladder			Melanoma of the Skin		
	Count	*Rate*	*S.E.*[3]	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
NAACCR U.S. Combined	**3,894,209**	**502.1**	**0.26**	**943,855**	**115.0**	**0.12**	**555,126**	**73.0**	**0.10**	**350,832**	**45.9**	**0.08**	**262,379**	**35.8**	**0.07**	**201,442**	**26.6**	**0.06**
Alabama	67,581	533.2	2.11	17,371	129.3	1.01	11,675	92.5	0.88	6,498	52.2	0.67	4,058	34.0	0.55	3,304	27.2	0.49
Alaska	6,949	447.7	6.08	1,548	91.4	2.63	937	67.5	2.49	707	46.6	1.99	451	33.8	1.78	248	15.2	1.07
Arizona	72,737	413.8	1.56	14,985	80.8	0.67	9,842	56.0	0.57	6,778	39.2	0.48	5,534	32.2	0.44	4,429	25.7	0.39
Arkansas	42,168	528.0	2.63	10,204	120.7	1.22	7,946	99.0	1.13	3,937	50.1	0.82	2,745	36.0	0.70	1,921	25.1	0.59
California	407,224	459.3	0.74	100,652	109.3	0.35	42,877	51.0	0.25	37,576	42.5	0.22	25,904	31.4	0.20	25,399	28.9	0.19
CA: Greater California	231,016	468.0	1.00	55,682	108.4	0.47	25,441	54.1	0.35	20,841	42.4	0.30	15,128	32.9	0.27	15,910	32.6	0.26
CA: Greater Bay Area	80,433	465.2	1.68	20,916	115.8	0.82	7,967	49.0	0.56	7,006	40.6	0.50	4,915	30.4	0.44	5,443	31.8	0.44
CA: Los Angeles	95,589	434.0	1.44	23,993	105.9	0.70	9,449	45.6	0.48	9,717	44.2	0.46	5,855	28.7	0.38	4,043	18.5	0.30
Colorado	55,158	445.4	1.98	14,649	110.9	0.96	5,581	48.3	0.67	4,577	37.9	0.58	3,696	32.7	0.56	3,422	27.8	0.49
Connecticut	50,632	526.7	2.40	12,169	118.8	1.10	6,578	69.7	0.88	4,205	44.4	0.70	4,342	46.9	0.73	2,566	27.4	0.55
Delaware	14,378	570.7	4.87	3,837	141.6	2.33	2,025	81.3	1.85	1,069	43.5	1.36	1,017	42.2	1.35	969	39.5	1.30
District of Columbia	7,245	534.4	6.49	2,188	159.7	3.52	883	67.4	2.34	641	48.7	1.99	297	23.3	1.39	168	12.4	0.99
Florida	287,296	479.4	0.91	65,177	103.6	0.41	43,754	71.4	0.34	25,244	42.5	0.27	20,090	33.2	0.24	17,089	29.2	0.23
Georgia	118,328	534.1	1.63	30,481	129.3	0.77	17,952	84.6	0.66	10,783	48.9	0.49	6,580	33.3	0.43	7,314	33.9	0.42
Hawaii	17,112	444.1	3.48	3,582	90.1	1.55	2,178	56.9	1.25	2,003	51.3	1.18	893	23.7	0.81	1,023	26.5	0.85
Idaho	19,579	477.5	3.50	5,187	119.7	1.70	2,219	55.7	1.21	1,612	39.8	1.01	1,479	37.9	1.01	1,311	32.2	0.91
Illinois	164,223	520.1	1.32	39,845	119.4	0.61	24,277	79.2	0.52	16,315	52.5	0.42	11,139	37.5	0.36	7,142	23.0	0.28
Indiana	81,526	494.3	1.78	16,713	95.7	0.76	14,434	88.9	0.76	7,932	48.8	0.56	5,747	36.7	0.50	3,539	21.8	0.38
Iowa	44,898	529.6	2.54	10,064	112.2	1.14	6,611	78.3	0.98	4,310	51.7	0.80	3,190	38.4	0.69	2,394	29.1	0.61
Kansas	38,405	521.8	2.72	9,518	124.1	1.30	5,306	73.1	1.02	3,439	47.4	0.82	2,748	38.8	0.75	2,167	30.2	0.66
Kentucky	67,517	590.8	2.35	13,778	113.0	0.99	13,203	116.3	1.04	6,687	59.3	0.75	4,304	40.2	0.63	3,403	30.7	0.54
Louisiana	65,307	576.5	2.33	17,311	144.4	1.13	10,066	90.8	0.94	6,303	56.0	0.73	3,530	33.5	0.58	2,482	23.0	0.48
Maine	21,320	517.6	3.65	4,491	99.8	1.53	3,464	83.9	1.46	1,717	42.7	1.06	1,892	47.5	1.12	1,134	28.1	0.86
Maryland	72,723	502.5	1.92	20,406	131.5	0.95	9,265	66.6	0.71	6,025	42.6	0.56	4,916	37.0	0.54	4,032	28.5	0.46
Massachusetts	87,754	504.1	1.74	21,277	114.2	0.80	11,876	70.3	0.66	7,327	42.9	0.51	6,718	40.5	0.50	4,332	25.4	0.40
Michigan	138,639	516.4	1.42	36,195	126.3	0.68	20,448	77.2	0.55	11,508	44.0	0.42	9,972	39.1	0.40	5,904	22.7	0.30
MI: Metropolitan Detroit	57,850	583.8	2.50	16,298	155.2	1.25	8,152	85.0	0.97	4,722	48.5	0.73	3,965	42.4	0.69	2,041	21.2	0.48
Mississippi	41,771	558.7	2.82	10,748	135.5	1.34	7,466	101.4	1.21	4,300	58.2	0.92	2,145	30.9	0.69	1,662	23.1	0.59
Missouri	80,984	502.0	1.80	17,428	101.0	0.78	14,218	89.0	0.76	7,820	49.7	0.57	5,198	33.8	0.48	3,831	24.3	0.40

Total Cancer Cases and Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates[1] for the Five Most Commonly Diagnosed Cancers
Selected Areas in the NAACCR United States Combined[2], 2010-2014, All Races, Males

	All Sites			Prostate			Lung & Bronchus			Colon & Rectum			Bladder			Melanoma of the Skin		
	Count	*Rate*	*S.E.*[3]	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
Montana	14,771	485.9	4.13	3,856	116.6	1.93	1,810	60.9	1.47	1,334	44.2	1.25	1,056	36.0	1.13	897	30.3	1.05
Nebraska	24,322	507.5	3.31	5,977	119.6	1.58	3,374	71.8	1.26	2,375	49.9	1.04	1,704	37.1	0.91	1,122	23.5	0.71
New Hampshire	19,813	527.5	3.88	5,094	123.5	1.79	2,581	71.2	1.45	1,504	40.9	1.09	1,714	48.1	1.20	1,218	33.3	0.99
New Jersey	122,895	543.0	1.58	33,215	139.4	0.78	14,382	65.6	0.56	10,728	47.9	0.47	8,887	41.5	0.45	6,416	28.7	0.37
New York	270,901	546.4	1.07	71,412	136.8	0.52	33,864	69.9	0.39	22,839	46.9	0.32	19,314	41.0	0.30	11,099	22.8	0.22
North Carolina	130,183	527.9	1.51	32,878	125.0	0.71	21,415	88.5	0.62	10,565	43.7	0.44	8,158	35.5	0.40	7,449	31.4	0.37
North Dakota	9,651	510.4	5.31	2,446	123.2	2.55	1,308	70.8	1.99	1,000	53.8	1.74	683	37.0	1.44	466	24.9	1.18
Ohio	157,260	504.6	1.30	37,060	111.8	0.59	25,857	83.7	0.53	14,626	47.9	0.41	11,494	38.6	0.37	7,519	24.9	0.29
Oklahoma	49,300	502.0	2.31	11,133	108.3	1.05	8,448	86.6	0.96	4,705	48.6	0.72	3,158	33.6	0.61	2,276	23.9	0.51
Oregon	50,882	467.9	2.13	11,791	101.1	0.96	6,714	63.1	0.79	4,313	40.8	0.64	3,946	38.0	0.62	3,324	31.1	0.55
Pennsylvania	196,353	543.9	1.25	45,104	117.5	0.56	28,146	78.3	0.47	17,716	49.9	0.38	15,434	43.9	0.36	10,098	28.7	0.29
Rhode Island	14,563	518.5	4.40	3,240	108.5	1.95	2,139	77.7	1.72	1,135	41.2	1.25	1,234	45.1	1.31	781	28.5	1.04
South Carolina	65,603	521.3	2.11	16,502	121.4	0.98	10,738	85.7	0.86	5,531	44.7	0.62	4,026	34.7	0.57	3,466	28.4	0.50
South Dakota	11,379	498.4	4.77	2,815	116.8	2.24	1,568	69.7	1.79	1,165	51.6	1.54	771	35.1	1.28	558	24.9	1.08
Tennessee	89,477	529.2	1.83	21,512	119.1	0.84	16,028	95.7	0.78	7,724	46.8	0.55	5,466	34.5	0.48	4,232	26.0	0.41
Texas	262,427	465.1	0.94	58,205	99.4	0.43	36,319	68.0	0.37	26,047	46.3	0.30	13,966	27.4	0.24	9,825	17.7	0.19
Utah	25,518	457.4	2.94	7,404	130.6	1.56	1,717	32.7	0.81	1,942	35.5	0.82	1,515	29.8	0.78	2,643	47.7	0.95
Vermont	9,122	493.3	5.36	1,989	98.4	2.28	1,325	71.8	2.04	721	39.3	1.52	691	38.9	1.52	715	39.5	1.54
Virginia	93,384	460.8	1.56	23,473	107.6	0.72	14,291	73.1	0.63	8,205	41.1	0.47	5,791	31.1	0.42	4,861	24.6	0.36
Washington	88,331	502.1	1.74	21,829	116.3	0.81	11,011	65.5	0.64	7,123	40.9	0.50	6,175	37.6	0.49	5,344	30.5	0.43
WA: Seattle/Puget Sound	62,082	525.1	2.18	15,504	123.3	1.03	7,398	66.1	0.79	4,930	41.8	0.61	4,246	38.9	0.62	3,978	33.7	0.55
West Virginia	29,544	523.7	3.14	6,057	99.6	1.31	5,639	98.7	1.35	2,939	53.1	1.01	2,175	39.9	0.88	1,350	25.3	0.71
Wisconsin	80,116	515.8	1.86	19,236	116.0	0.86	10,633	69.5	0.69	6,648	43.6	0.55	5,912	39.9	0.53	4,173	27.3	0.43
Wyoming	6,960	447.2	5.57	1,823	108.5	2.64	738	49.4	1.89	634	41.7	1.72	524	36.0	1.63	425	28.5	1.43

1. Rates are per 100,000 population and are age-adjusted by five-year age groups to the 2000 U.S. standard population based on single years of age.
2. See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.
3. Abbreviations: S.E. - Standard Error

\- Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for the U.S. combined.

Total Cancer Cases and Average Annual Age-adjusted (1996 Canadian Standard) Cancer Incidence Rates[1] for the Five Most Commonly Diagnosed Cancers
Selected Areas in the NAACCR United States Combined[2], 2010-2014, All Races, Males

	All Sites			Prostate			Lung & Bronchus			Colon & Rectum			Bladder			Melanoma of the Skin		
	Count	*Rate*	*S.E.*[3]	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
NAACCR U.S. Combined	**3,894,209**	**479.8**	**0.25**	**943,855**	**112.2**	**0.12**	**555,126**	**69.0**	**0.09**	**350,832**	**43.3**	**0.07**	**262,379**	**33.1**	**0.07**	**201,442**	**25.3**	**0.06**
Alabama	67,581	510.1	2.00	17,371	125.4	0.96	11,675	87.9	0.82	6,498	49.5	0.63	4,058	31.7	0.50	3,304	25.9	0.46
Alaska	6,949	426.5	5.58	1,548	88.7	2.46	937	63.3	2.25	707	43.9	1.81	451	31.7	1.61	248	14.8	1.01
Arizona	72,737	394.6	1.48	14,985	78.5	0.64	9,842	52.7	0.53	6,778	37.1	0.45	5,534	29.7	0.40	4,429	24.4	0.37
Arkansas	42,168	506.4	2.50	10,204	117.8	1.18	7,946	94.4	1.07	3,937	47.6	0.77	2,745	33.4	0.64	1,921	23.8	0.55
California	407,224	438.0	0.70	100,652	106.6	0.34	42,877	47.7	0.23	37,576	40.1	0.21	25,904	28.9	0.18	25,399	27.4	0.17
CA: Greater California	231,016	446.1	0.94	55,682	105.6	0.45	25,441	50.6	0.32	20,841	40.1	0.28	15,128	30.2	0.25	15,910	30.9	0.25
CA: Greater Bay Area	80,433	443.5	1.59	20,916	113.0	0.80	7,967	45.6	0.52	7,006	38.2	0.47	4,915	27.9	0.40	5,443	30.2	0.42
CA: Los Angeles	95,589	414.7	1.37	23,993	103.5	0.68	9,449	42.7	0.45	9,717	41.9	0.43	5,855	26.4	0.35	4,043	17.5	0.28
Colorado	55,158	425.0	1.86	14,649	108.0	0.92	5,581	45.2	0.62	4,577	35.6	0.54	3,696	30.2	0.51	3,422	26.6	0.47
Connecticut	50,632	502.9	2.29	12,169	116.5	1.08	6,578	65.8	0.83	4,205	41.7	0.66	4,342	43.3	0.67	2,566	25.9	0.52
Delaware	14,378	543.8	4.60	3,837	138.1	2.25	2,025	76.4	1.71	1,069	40.8	1.27	1,017	38.7	1.22	969	37.5	1.23
District of Columbia	7,245	516.0	6.21	2,188	155.6	3.40	883	64.6	2.22	641	46.4	1.88	297	21.8	1.29	168	12.0	0.94
Florida	287,296	459.4	0.87	65,177	101.5	0.40	43,754	67.7	0.33	25,244	40.4	0.26	20,090	30.8	0.22	17,089	27.6	0.22
Georgia	118,328	510.9	1.53	30,481	126.1	0.74	17,952	80.0	0.61	10,783	46.4	0.46	6,580	30.8	0.39	7,314	32.2	0.39
Hawaii	17,112	424.6	3.32	3,582	87.1	1.48	2,178	53.5	1.17	2,003	49.1	1.12	893	22.0	0.75	1,023	25.7	0.82
Idaho	19,579	455.2	3.30	5,187	115.8	1.63	2,219	52.1	1.12	1,612	37.6	0.95	1,479	35.1	0.92	1,311	31.1	0.87
Illinois	164,223	496.5	1.25	39,845	116.5	0.59	24,277	74.7	0.49	16,315	49.4	0.39	11,139	34.8	0.33	7,142	21.9	0.26
Indiana	81,526	472.0	1.68	16,713	92.9	0.73	14,434	84.3	0.71	7,932	46.1	0.53	5,747	34.0	0.45	3,539	20.8	0.36
Iowa	44,898	505.2	2.43	10,064	109.2	1.10	6,611	73.9	0.92	4,310	48.5	0.75	3,190	35.5	0.64	2,394	27.9	0.59
Kansas	38,405	499.3	2.59	9,518	120.7	1.26	5,306	69.1	0.96	3,439	44.8	0.78	2,748	36.0	0.70	2,167	28.9	0.63
Kentucky	67,517	564.7	2.21	13,778	109.9	0.95	13,203	110.3	0.98	6,687	56.2	0.70	4,304	37.3	0.58	3,403	29.4	0.51
Louisiana	65,307	553.2	2.21	17,311	141.1	1.09	10,066	86.3	0.88	6,303	53.3	0.69	3,530	31.3	0.54	2,482	21.8	0.45
Maine	21,320	493.4	3.47	4,491	97.3	1.47	3,464	79.2	1.37	1,717	40.2	1.00	1,892	44.1	1.03	1,134	26.8	0.82
Maryland	72,723	480.4	1.82	20,406	128.8	0.92	9,265	62.8	0.67	6,025	40.1	0.53	4,916	34.0	0.49	4,032	27.1	0.44
Massachusetts	87,754	482.1	1.66	21,277	111.8	0.78	11,876	66.2	0.62	7,327	40.3	0.48	6,718	37.6	0.47	4,332	24.2	0.38
Michigan	138,639	493.8	1.35	36,195	123.4	0.66	20,448	73.0	0.52	11,508	41.4	0.39	9,972	36.3	0.37	5,904	21.7	0.29
MI: Metropolitan Detroit	57,850	558.5	2.38	16,298	151.7	1.22	8,152	80.2	0.91	4,722	45.6	0.68	3,965	39.3	0.64	2,041	20.1	0.46
Mississippi	41,771	536.4	2.67	10,748	132.5	1.30	7,466	96.4	1.13	4,300	55.5	0.86	2,145	28.7	0.63	1,662	22.1	0.55
Missouri	80,984	480.3	1.71	17,428	98.8	0.76	14,218	84.4	0.72	7,820	46.8	0.54	5,198	31.4	0.44	3,831	23.3	0.38

Total Cancer Cases and Average Annual Age-adjusted (1996 Canadian Standard) Cancer Incidence Rates[1] for the Five Most Commonly Diagnosed Cancers
Selected Areas in the NAACCR United States Combined[2], 2010-2014, All Races, Males

	All Sites			Prostate			Lung & Bronchus			Colon & Rectum			Bladder			Melanoma of the Skin		
	Count	*Rate*	*S.E.*[3]	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
Montana	14,771	465.5	3.93	3,856	113.7	1.86	1,810	57.0	1.36	1,334	42.2	1.18	1,056	33.3	1.04	897	29.3	1.02
Nebraska	24,322	485.3	3.17	5,977	116.1	1.53	3,374	67.9	1.19	2,375	47.2	0.99	1,704	34.4	0.84	1,122	22.7	0.69
New Hampshire	19,813	504.2	3.69	5,094	121.1	1.74	2,581	66.7	1.34	1,504	38.3	1.02	1,714	44.7	1.10	1,218	31.7	0.93
New Jersey	122,895	518.3	1.50	33,215	136.0	0.76	14,382	61.8	0.52	10,728	45.1	0.44	8,887	38.3	0.41	6,416	27.2	0.35
New York	270,901	522.2	1.02	71,412	133.6	0.51	33,864	65.9	0.36	22,839	44.2	0.30	19,314	37.9	0.28	11,099	21.7	0.21
North Carolina	130,183	505.3	1.42	32,878	121.7	0.68	21,415	83.9	0.58	10,565	41.4	0.41	8,158	32.9	0.37	7,449	29.9	0.35
North Dakota	9,651	488.6	5.09	2,446	120.1	2.48	1,308	66.5	1.87	1,000	51.3	1.66	683	34.1	1.33	466	24.3	1.16
Ohio	157,260	482.8	1.24	37,060	109.1	0.58	25,857	79.3	0.50	14,626	45.2	0.38	11,494	35.8	0.34	7,519	23.8	0.28
Oklahoma	49,300	480.0	2.19	11,133	104.9	1.00	8,448	82.1	0.90	4,705	46.2	0.68	3,158	31.3	0.56	2,276	22.7	0.48
Oregon	50,882	447.1	2.02	11,791	98.6	0.92	6,714	59.3	0.73	4,313	38.4	0.60	3,946	35.1	0.57	3,324	29.9	0.53
Pennsylvania	196,353	519.9	1.20	45,104	114.7	0.55	28,146	74.0	0.45	17,716	46.9	0.36	15,434	40.7	0.33	10,098	27.3	0.28
Rhode Island	14,563	494.6	4.20	3,240	106.1	1.90	2,139	73.0	1.61	1,135	38.5	1.17	1,234	41.9	1.22	781	27.3	1.00
South Carolina	65,603	497.9	1.98	16,502	118.3	0.93	10,738	81.1	0.80	5,531	42.4	0.58	4,026	31.9	0.51	3,466	27.1	0.47
South Dakota	11,379	476.4	4.56	2,815	114.0	2.18	1,568	65.1	1.67	1,165	48.7	1.46	771	32.4	1.18	558	24.1	1.05
Tennessee	89,477	507.1	1.73	21,512	116.2	0.80	16,028	90.9	0.73	7,724	44.4	0.51	5,466	32.1	0.44	4,232	24.8	0.39
Texas	262,427	443.5	0.88	58,205	96.2	0.41	36,319	64.1	0.34	26,047	43.9	0.28	13,966	25.3	0.22	9,825	16.8	0.17
Utah	25,518	438.2	2.79	7,404	127.2	1.50	1,717	30.8	0.75	1,942	33.6	0.77	1,515	27.6	0.72	2,643	45.5	0.90
Vermont	9,122	471.7	5.09	1,989	95.6	2.19	1,325	67.9	1.91	721	37.0	1.42	691	36.3	1.40	715	37.8	1.46
Virginia	93,384	439.9	1.47	23,473	105.1	0.70	14,291	68.9	0.59	8,205	38.8	0.44	5,791	28.7	0.38	4,861	23.3	0.34
Washington	88,331	478.4	1.64	21,829	113.4	0.78	11,011	61.3	0.59	7,123	38.5	0.46	6,175	34.6	0.45	5,344	29.3	0.41
WA: Seattle/Puget Sound	62,082	499.7	2.05	15,504	120.1	0.99	7,398	61.8	0.73	4,930	39.3	0.57	4,246	35.6	0.56	3,978	32.3	0.52
West Virginia	29,544	501.0	2.98	6,057	96.8	1.26	5,639	93.9	1.27	2,939	50.3	0.95	2,175	37.3	0.81	1,350	24.2	0.68
Wisconsin	80,116	492.5	1.77	19,236	113.1	0.83	10,633	65.6	0.65	6,648	41.0	0.51	5,912	36.9	0.49	4,173	26.1	0.41
Wyoming	6,960	429.0	5.28	1,823	106.2	2.55	738	46.5	1.75	634	39.5	1.61	524	33.7	1.50	425	27.3	1.36

1. Rates are per 100,000 and are age-adjusted to the 1996 Canadian population standard.
2. See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.
3. Abbreviations: S.E. - Standard Error

\- Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for the U.S. combined.

Total Cancer Cases and Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates[1] for the Five Most Commonly Diagnosed Cancers
Selected Areas in the NAACCR United States Combined[2], 2010-2014, All Races, Females

	All Sites			Breast			Lung & Bronchus			Colon & Rectum			Uterus Corpus & Uterus, NOS[3]			Thyroid		
	Count	*Rate*	*S.E.*[3]	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
NAACCR U.S. Combined	**3,781,169**	**418.0**	**0.22**	**1,107,649**	**123.6**	**0.12**	**493,029**	**52.8**	**0.08**	**323,272**	**34.8**	**0.06**	**241,508**	**25.9**	**0.05**	**169,974**	**21.1**	**0.05**
Alabama	59,329	394.9	1.66	17,811	119.6	0.92	8,235	52.1	0.58	5,651	36.9	0.50	2,968	19.1	0.36	1,712	13.0	0.32
Alaska	6,568	411.9	5.38	2,085	125.4	2.90	773	53.3	2.04	620	40.2	1.71	431	24.6	1.26	299	17.5	1.05
Arizona	72,444	375.0	1.42	21,537	112.4	0.79	9,264	45.4	0.48	5,908	29.8	0.40	4,260	21.6	0.34	3,979	23.7	0.39
Arkansas	36,111	398.1	2.15	10,122	112.7	1.15	5,818	60.7	0.81	3,456	37.1	0.64	1,997	21.6	0.49	1,180	15.1	0.45
California	404,208	388.5	0.62	125,201	120.7	0.35	41,600	39.9	0.20	34,469	32.6	0.18	26,391	24.7	0.15	18,747	19.1	0.14
CA: Greater California	225,193	395.6	0.85	69,313	122.1	0.47	24,875	43.4	0.28	18,789	32.6	0.24	13,933	23.7	0.20	10,305	19.4	0.19
CA: Greater Bay Area	79,420	392.4	1.42	25,087	124.6	0.80	7,912	39.0	0.45	6,667	32.3	0.40	5,363	25.7	0.36	3,212	16.9	0.30
CA: Los Angeles	99,469	370.0	1.19	30,772	114.6	0.66	8,801	32.9	0.36	9,004	32.9	0.35	7,095	25.8	0.31	5,229	20.0	0.28
Colorado	54,644	390.2	1.71	17,541	123.7	0.95	5,751	41.6	0.56	4,259	30.8	0.48	3,321	22.3	0.40	3,154	23.5	0.43
Connecticut	52,718	458.4	2.06	15,767	139.2	1.14	6,767	56.8	0.71	4,159	34.3	0.55	3,483	29.2	0.50	2,827	29.1	0.56
Delaware	13,375	456.4	4.05	3,886	133.1	2.20	1,930	62.9	1.45	986	32.9	1.07	890	29.0	1.00	578	23.6	1.01
District of Columbia	7,606	446.1	5.24	2,389	143.5	3.00	845	49.5	1.74	690	40.3	1.57	549	31.2	1.36	396	22.5	1.18
Florida	266,096	396.9	0.80	75,212	115.5	0.44	38,674	53.1	0.27	22,825	32.3	0.22	15,788	23.1	0.19	9,412	17.6	0.19
Georgia	110,666	410.9	1.25	33,614	123.5	0.68	14,072	52.2	0.45	9,578	35.6	0.37	6,133	21.6	0.28	4,359	16.6	0.26
Hawaii	17,260	407.4	3.21	5,670	136.0	1.87	1,717	38.1	0.94	1,599	36.5	0.94	1,237	30.0	0.88	831	23.2	0.83
Idaho	18,042	410.5	3.12	5,295	120.5	1.69	2,117	46.8	1.03	1,437	32.5	0.88	1,111	24.2	0.74	980	24.9	0.81
Illinois	165,887	437.9	1.10	48,665	130.0	0.60	22,372	57.8	0.39	14,948	38.1	0.32	11,336	29.0	0.28	7,370	21.7	0.26
Indiana	82,508	425.6	1.51	23,084	120.1	0.81	12,158	60.8	0.56	7,666	38.5	0.45	5,390	27.0	0.38	3,015	17.8	0.33
Iowa	42,359	439.5	2.21	11,520	122.8	1.19	5,377	53.0	0.74	4,114	39.8	0.64	3,042	31.2	0.58	1,575	20.1	0.52
Kansas	35,882	430.0	2.33	10,182	123.5	1.26	4,596	52.8	0.79	3,129	36.1	0.66	2,104	24.7	0.55	1,833	25.3	0.61
Kentucky	63,000	472.7	1.92	16,413	123.3	0.98	11,053	79.7	0.77	5,774	42.4	0.57	3,521	25.8	0.44	2,526	21.6	0.44
Louisiana	55,904	420.8	1.81	16,330	123.2	0.99	7,513	54.9	0.64	5,551	41.4	0.57	2,610	19.1	0.38	2,371	19.4	0.41
Maine	21,056	454.0	3.25	5,743	125.5	1.72	3,189	65.3	1.18	1,638	34.3	0.88	1,506	30.9	0.82	822	22.3	0.82
Maryland	74,065	421.0	1.58	23,092	131.0	0.88	9,245	52.0	0.55	5,920	33.3	0.44	4,867	26.4	0.39	3,554	22.0	0.38
Massachusetts	94,120	449.3	1.50	28,058	136.1	0.83	13,038	60.5	0.54	7,338	33.7	0.40	6,255	28.9	0.37	5,322	29.1	0.41
Michigan	131,536	426.3	1.21	37,301	122.2	0.65	18,930	58.8	0.43	10,849	34.0	0.33	9,267	28.9	0.31	5,218	19.8	0.28
MI: Metropolitan Detroit	55,534	458.5	2.00	15,788	131.6	1.07	8,076	64.8	0.73	4,704	37.5	0.56	3,787	30.3	0.50	2,511	23.6	0.48
Mississippi	35,981	407.4	2.19	10,104	115.5	1.17	5,172	56.3	0.79	3,786	41.8	0.69	1,778	19.6	0.47	1,261	15.9	0.46
Missouri	79,928	427.6	1.55	23,218	125.9	0.85	12,540	64.3	0.58	7,077	36.6	0.45	4,997	26.1	0.38	2,866	17.8	0.34

Total Cancer Cases and Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates[1] for the Five Most Commonly Diagnosed Cancers
Selected Areas in the NAACCR United States Combined[2], 2010-2014, All Races, Females

	All Sites			Breast			Lung & Bronchus			Colon & Rectum			Uterus Corpus & Uterus, NOS[3]			Thyroid		
	Count	*Rate*	*S.E.*[3]	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
Montana	13,450	424.7	3.80	3,893	123.1	2.06	1,791	53.8	1.30	1,117	34.1	1.05	860	25.9	0.92	561	21.6	0.95
Nebraska	22,633	416.0	2.84	6,538	121.8	1.55	2,834	50.2	0.96	2,170	38.2	0.84	1,470	26.6	0.71	975	21.0	0.69
New Hampshire	19,517	463.1	3.42	5,914	140.4	1.88	2,726	63.3	1.24	1,481	34.2	0.91	1,466	32.8	0.88	845	23.3	0.83
New Jersey	125,460	452.9	1.31	36,075	132.0	0.71	15,020	52.5	0.44	10,947	37.8	0.37	8,874	31.1	0.34	6,800	27.9	0.35
New York	273,599	451.0	0.88	76,791	129.0	0.48	34,081	54.2	0.30	22,514	35.7	0.24	19,341	31.0	0.23	15,178	28.3	0.23
North Carolina	124,067	421.4	1.22	37,914	129.4	0.68	17,109	56.1	0.43	9,839	33.0	0.34	7,459	24.4	0.29	4,854	18.3	0.27
North Dakota	8,513	414.5	4.67	2,397	121.4	2.58	1,048	48.9	1.54	872	40.0	1.40	527	25.4	1.14	426	24.4	1.23
Ohio	154,863	424.4	1.11	44,328	122.9	0.60	22,438	59.0	0.40	13,624	35.8	0.31	10,697	28.4	0.28	6,693	21.7	0.27
Oklahoma	46,231	412.3	1.96	13,103	117.8	1.05	6,873	58.7	0.72	4,236	36.9	0.58	2,496	21.7	0.45	1,731	17.5	0.43
Oregon	50,591	419.2	1.92	15,188	126.0	1.06	6,696	53.4	0.67	3,881	31.2	0.52	3,342	26.5	0.47	2,010	19.6	0.45
Pennsylvania	195,598	461.3	1.08	53,851	129.8	0.58	25,224	56.1	0.36	16,964	37.5	0.30	14,004	32.1	0.28	10,515	30.5	0.31
Rhode Island	15,625	457.9	3.80	4,331	130.3	2.04	2,265	63.7	1.37	1,271	34.7	1.01	1,117	31.9	0.98	784	27.1	1.00
South Carolina	60,486	411.9	1.72	18,597	127.2	0.96	8,268	53.7	0.60	5,051	33.8	0.49	3,505	22.7	0.39	2,222	17.2	0.38
South Dakota	10,661	428.1	4.31	3,167	130.7	2.41	1,324	50.8	1.42	1,041	38.8	1.25	708	28.2	1.09	412	19.8	1.01
Tennessee	82,826	418.7	1.49	23,909	121.1	0.80	12,555	61.0	0.55	7,307	36.4	0.43	4,633	22.7	0.34	3,231	18.5	0.33
Texas	251,958	380.2	0.77	74,269	111.5	0.41	29,302	44.7	0.26	21,373	32.2	0.22	14,447	21.1	0.18	11,944	18.3	0.17
Utah	23,175	373.9	2.49	7,012	114.7	1.39	1,449	24.1	0.64	1,686	27.7	0.68	1,643	25.9	0.65	1,856	28.8	0.68
Vermont	9,030	437.4	4.78	2,662	130.1	2.62	1,285	59.7	1.71	710	33.0	1.28	710	32.8	1.27	316	19.0	1.13
Virginia	94,849	398.7	1.32	30,271	126.9	0.74	12,257	50.8	0.47	7,934	33.1	0.38	5,915	23.8	0.31	4,277	19.6	0.31
Washington	86,489	437.9	1.53	26,686	134.9	0.85	10,713	53.7	0.53	6,534	32.9	0.42	5,342	25.6	0.36	3,679	20.4	0.34
WA: Seattle/Puget Sound	60,793	450.7	1.87	18,947	139.9	1.04	7,271	54.0	0.65	4,519	33.4	0.51	3,756	26.3	0.44	2,586	20.6	0.41
West Virginia	27,673	443.8	2.77	7,112	114.8	1.42	4,416	66.2	1.01	2,664	40.9	0.81	1,935	30.4	0.72	1,226	24.3	0.72
Wisconsin	76,465	437.3	1.63	21,974	127.9	0.89	9,884	54.8	0.56	6,139	33.6	0.44	5,428	29.9	0.41	2,947	19.7	0.37
Wyoming	6,117	382.5	5.04	1,827	113.5	2.74	725	44.1	1.67	490	30.5	1.41	357	21.5	1.18	305	21.3	1.26

1. Rates are per 100,000 population and are age-adjusted by five-year age groups to the 2000 U.S. standard population based on single years of age.
2. See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.
3. Abbreviations: NOS - Not Otherwise Specified; S.E. - Standard Error

\- Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for the U.S. combined.

Total Cancer Cases and Average Annual Age-adjusted (1996 Canadian Standard) Cancer Incidence Rates[1] for the Five Most Commonly Diagnosed Cancers Selected Areas in the NAACCR United States Combined[2], 2010-2014, All Races, Females

	All Sites			Breast			Lung & Bronchus			Colon & Rectum			Uterus Corpus & Uterus, NOS[3]			Thyroid		
	Count	*Rate*	*S.E.*[3]	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
NAACCR U.S. Combined	**3,781,169**	**403.7**	**0.21**	**1,107,649**	**120.4**	**0.12**	**493,029**	**50.3**	**0.07**	**323,272**	**32.8**	**0.06**	**241,508**	**25.4**	**0.05**	**169,974**	**21.5**	**0.05**
Alabama	59,329	382.0	1.62	17,811	116.6	0.90	8,235	50.1	0.56	5,651	35.0	0.48	2,968	18.6	0.35	1,712	13.2	0.33
Alaska	6,568	397.8	5.14	2,085	122.0	2.80	773	50.6	1.91	620	38.3	1.61	431	24.2	1.23	299	17.9	1.07
Arizona	72,444	361.6	1.38	21,537	109.4	0.77	9,264	43.0	0.45	5,908	28.0	0.37	4,260	21.2	0.33	3,979	24.2	0.40
Arkansas	36,111	385.4	2.09	10,122	109.6	1.13	5,818	58.4	0.78	3,456	35.3	0.62	1,997	21.1	0.49	1,180	15.4	0.47
California	404,208	374.3	0.60	125,201	117.5	0.34	41,600	37.6	0.19	34,469	30.6	0.17	26,391	24.2	0.15	18,747	19.4	0.14
CA: Greater California	225,193	381.1	0.82	69,313	118.8	0.46	24,875	41.0	0.26	18,789	30.6	0.23	13,933	23.3	0.20	10,305	19.7	0.20
CA: Greater Bay Area	79,420	377.6	1.37	25,087	121.5	0.78	7,912	36.5	0.42	6,667	30.3	0.38	5,363	25.0	0.35	3,212	17.2	0.31
CA: Los Angeles	99,469	356.7	1.15	30,772	111.6	0.65	8,801	31.1	0.34	9,004	31.0	0.33	7,095	25.4	0.31	5,229	20.3	0.29
Colorado	54,644	376.2	1.65	17,541	120.4	0.93	5,751	39.3	0.53	4,259	28.8	0.45	3,321	21.8	0.39	3,154	24.0	0.44
Connecticut	52,718	442.6	2.02	15,767	135.6	1.12	6,767	54.0	0.68	4,159	32.1	0.53	3,483	28.6	0.50	2,827	29.6	0.59
Delaware	13,375	441.3	3.96	3,886	129.5	2.15	1,930	59.9	1.39	986	30.9	1.02	890	28.6	0.99	578	24.1	1.05
District of Columbia	7,606	431.5	5.08	2,389	139.6	2.93	845	47.0	1.65	690	38.3	1.50	549	30.4	1.32	396	23.0	1.19
Florida	266,096	383.3	0.78	75,212	112.5	0.43	38,674	50.7	0.26	22,825	30.5	0.21	15,788	22.7	0.19	9,412	17.9	0.19
Georgia	110,666	397.0	1.21	33,614	120.3	0.67	14,072	49.8	0.42	9,578	33.8	0.35	6,133	21.2	0.28	4,359	16.9	0.26
Hawaii	17,260	395.1	3.15	5,670	133.0	1.84	1,717	36.2	0.90	1,599	34.6	0.91	1,237	29.5	0.87	831	23.8	0.86
Idaho	18,042	396.2	3.03	5,295	117.1	1.65	2,117	44.5	0.98	1,437	30.5	0.83	1,111	23.7	0.73	980	25.5	0.84
Illinois	165,887	422.6	1.07	48,665	126.6	0.59	22,372	55.0	0.38	14,948	35.8	0.30	11,336	28.5	0.27	7,370	22.1	0.26
Indiana	82,508	411.5	1.48	23,084	117.1	0.79	12,158	58.3	0.54	7,666	36.2	0.43	5,390	26.4	0.37	3,015	18.2	0.34
Iowa	42,359	424.3	2.18	11,520	119.4	1.17	5,377	50.6	0.71	4,114	37.1	0.61	3,042	30.5	0.57	1,575	20.5	0.54
Kansas	35,882	416.3	2.29	10,182	120.3	1.24	4,596	50.5	0.76	3,129	34.1	0.64	2,104	24.2	0.54	1,833	25.7	0.62
Kentucky	63,000	457.5	1.88	16,413	120.3	0.97	11,053	76.6	0.74	5,774	40.1	0.54	3,521	25.3	0.44	2,526	22.1	0.46
Louisiana	55,904	407.2	1.77	16,330	120.1	0.97	7,513	52.6	0.62	5,551	39.4	0.54	2,610	18.8	0.38	2,371	19.6	0.42
Maine	21,056	438.7	3.19	5,743	122.4	1.70	3,189	62.3	1.13	1,638	32.2	0.84	1,506	30.3	0.81	822	22.8	0.85
Maryland	74,065	406.4	1.54	23,092	127.6	0.86	9,245	49.3	0.52	5,920	31.3	0.42	4,867	25.9	0.38	3,554	22.3	0.39
Massachusetts	94,120	434.0	1.47	28,058	132.7	0.82	13,038	57.6	0.52	7,338	31.5	0.39	6,255	28.4	0.37	5,322	29.7	0.42
Michigan	131,536	412.0	1.18	37,301	119.1	0.64	18,930	56.2	0.42	10,849	32.0	0.32	9,267	28.4	0.30	5,218	20.3	0.29
MI: Metropolitan Detroit	55,534	442.1	1.95	15,788	128.1	1.05	8,076	61.6	0.70	4,704	35.2	0.53	3,787	29.7	0.50	2,511	24.0	0.50
Mississippi	35,981	394.4	2.14	10,104	112.8	1.15	5,172	54.1	0.76	3,786	39.8	0.66	1,778	19.2	0.47	1,261	16.2	0.47
Missouri	79,928	413.2	1.52	23,218	122.6	0.83	12,540	61.5	0.56	7,077	34.4	0.42	4,997	25.6	0.37	2,866	18.1	0.35

Total Cancer Cases and Average Annual Age-adjusted (1996 Canadian Standard) Cancer Incidence Rates[1] for the Five Most Commonly Diagnosed Cancers
Selected Areas in the NAACCR United States Combined[2], 2010-2014, All Races, Females

	All Sites			Breast			Lung & Bronchus			Colon & Rectum			Uterus Corpus & Uterus, NOS[3]			Thyroid		
	Count	*Rate*	*S.E.*[3]	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
Montana	13,450	410.6	3.72	3,893	119.7	2.02	1,791	51.0	1.23	1,117	32.2	1.01	860	25.3	0.90	561	22.1	0.99
Nebraska	22,633	402.0	2.79	6,538	118.6	1.53	2,834	48.1	0.93	2,170	35.8	0.80	1,470	26.1	0.70	975	21.4	0.71
New Hampshire	19,517	447.8	3.35	5,914	136.9	1.85	2,726	60.4	1.19	1,481	31.9	0.87	1,466	32.3	0.87	845	23.7	0.86
New Jersey	125,460	437.0	1.28	36,075	128.6	0.70	15,020	49.9	0.42	10,947	35.4	0.35	8,874	30.5	0.33	6,800	28.4	0.36
New York	273,599	435.3	0.86	76,791	125.6	0.47	34,081	51.6	0.29	22,514	33.5	0.23	19,341	30.4	0.22	15,178	28.8	0.24
North Carolina	124,067	407.4	1.19	37,914	126.0	0.66	17,109	53.7	0.42	9,839	31.2	0.32	7,459	23.9	0.28	4,854	18.6	0.27
North Dakota	8,513	401.9	4.61	2,397	118.1	2.54	1,048	47.1	1.50	872	37.8	1.35	527	24.9	1.13	426	24.9	1.27
Ohio	154,863	410.5	1.09	44,328	119.6	0.59	22,438	56.4	0.39	13,624	33.6	0.30	10,697	27.8	0.28	6,693	22.2	0.28
Oklahoma	46,231	399.2	1.91	13,103	114.7	1.03	6,873	56.3	0.69	4,236	35.0	0.55	2,496	21.3	0.44	1,731	17.8	0.44
Oregon	50,591	404.8	1.87	15,188	122.7	1.03	6,696	50.6	0.63	3,881	29.2	0.49	3,342	26.0	0.46	2,010	20.0	0.46
Pennsylvania	195,598	446.0	1.06	53,851	126.4	0.57	25,224	53.6	0.35	16,964	35.2	0.29	14,004	31.5	0.28	10,515	31.1	0.32
Rhode Island	15,625	443.3	3.74	4,331	127.1	2.02	2,265	60.7	1.32	1,271	32.5	0.97	1,117	31.3	0.97	784	27.7	1.04
South Carolina	60,486	398.2	1.67	18,597	123.8	0.94	8,268	51.3	0.57	5,051	32.0	0.46	3,505	22.3	0.39	2,222	17.5	0.39
South Dakota	10,661	413.3	4.23	3,167	127.4	2.38	1,324	48.4	1.37	1,041	36.3	1.19	708	27.5	1.08	412	20.2	1.04
Tennessee	82,826	405.5	1.45	23,909	118.2	0.79	12,555	58.6	0.53	7,307	34.5	0.41	4,633	22.3	0.34	3,231	18.9	0.34
Texas	251,958	367.0	0.74	74,269	108.6	0.40	29,302	42.4	0.25	21,373	30.4	0.21	14,447	20.8	0.18	11,944	18.6	0.17
Utah	23,175	362.2	2.41	7,012	111.5	1.35	1,449	22.9	0.61	1,686	26.2	0.65	1,643	25.5	0.64	1,856	29.5	0.69
Vermont	9,030	422.3	4.68	2,662	126.9	2.58	1,285	57.0	1.63	710	30.5	1.20	710	32.0	1.25	316	19.7	1.19
Virginia	94,849	384.8	1.28	30,271	123.6	0.73	12,257	48.3	0.44	7,934	31.2	0.36	5,915	23.3	0.31	4,277	20.0	0.31
Washington	86,489	421.8	1.48	26,686	131.2	0.83	10,713	50.7	0.50	6,534	30.9	0.39	5,342	25.0	0.35	3,679	20.7	0.35
WA: Seattle/Puget Sound	60,793	433.8	1.81	18,947	136.2	1.02	7,271	50.8	0.61	4,519	31.4	0.48	3,756	25.7	0.43	2,586	20.9	0.42
West Virginia	27,673	430.3	2.72	7,112	111.9	1.39	4,416	63.5	0.98	2,664	38.5	0.78	1,935	30.2	0.72	1,226	24.8	0.75
Wisconsin	76,465	422.5	1.59	21,974	124.7	0.87	9,884	52.5	0.54	6,139	31.3	0.42	5,428	29.2	0.41	2,947	20.1	0.39
Wyoming	6,117	371.2	4.92	1,827	110.4	2.68	725	42.0	1.60	490	29.0	1.35	357	21.1	1.16	305	21.9	1.31

1. Rates are per 100,000 and are age-adjusted to the 1996 Canadian population standard.
2. See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.
3. Abbreviations: NOS - Not Otherwise Specified; S.E. - Standard Error

\- Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for the U.S. combined.

NAACCR U.S. Combined Incidence 2010-2014, The White Population

Total Cancer Cases and Average Annual Age-adjusted Cancer Incidence Rates*
NAACCR United States Combined, 2010-2014, White

	MALES (WHITE)					FEMALES (WHITE)				
SITES	*Count*	*Adj. Rate 2000 U.S.*	*S.E.*	*Adj. Rate 1996 Cdn*	*S.E.*	*Count*	*Adj. Rate 2000 U.S.*	*S.E.*	*Adj. Rate 1996 Cdn*	*S.E.*
All Sites (invasive & in situ bladder)	3,265,680	497.5	0.28	475.1	0.27	3,168,903	423.8	0.24	409.4	0.24
Oral Cavity & Pharynx	122,398	17.9	0.05	17.2	0.05	49,388	6.5	0.03	6.2	0.03
Lip	6,579	1.0	0.01	1.0	0.01	2,453	0.3	0.01	0.3	0.01
Tongue	39,127	5.6	0.03	5.5	0.03	15,104	2.0	0.02	1.9	0.02
Major salivary gland	10,782	1.7	0.02	1.6	0.02	6,992	1.0	0.01	0.9	0.01
Floor of mouth	5,804	0.8	0.01	0.8	0.01	2,628	0.3	0.01	0.3	0.01
Gum & other mouth	12,421	1.9	0.02	1.8	0.02	10,097	1.3	0.01	1.2	0.01
Nasopharynx	3,795	0.6	0.01	0.6	0.01	1,529	0.2	0.01	0.2	0.01
Oropharynx	5,858	0.8	0.01	0.8	0.01	1,819	0.2	0.01	0.2	0.01
Hypopharynx	7,183	1.0	0.01	1.0	0.01	1,777	0.2	0.01	0.2	0.01
Digestive System	618,964	94.3	0.12	89.3	0.12	492,188	63.0	0.09	59.3	0.09
Esophagus	55,310	8.3	0.04	7.9	0.03	13,892	1.7	0.02	1.6	0.01
Stomach	53,959	8.4	0.04	7.9	0.03	30,863	4.0	0.02	3.8	0.02
Small intestine	16,836	2.6	0.02	2.5	0.02	14,738	1.9	0.02	1.9	0.02
Colon excl. rectum	196,571	30.7	0.07	28.8	0.07	198,649	25.3	0.06	23.6	0.06
Rectum & rectosigmoid	93,175	14.1	0.05	13.5	0.04	65,393	8.7	0.03	8.3	0.03
Anus, anal canal & anorectum	9,610	1.4	0.02	1.4	0.01	17,685	2.3	0.02	2.2	0.02
Liver	69,332	9.9	0.04	9.4	0.04	22,992	2.9	0.02	2.8	0.02
Gallbladder	4,718	0.7	0.01	0.7	0.01	10,080	1.3	0.01	1.2	0.01
Pancreas	92,145	14.1	0.05	13.3	0.04	85,916	10.7	0.04	10.0	0.04
Respiratory System	520,770	79.8	0.11	75.5	0.11	442,050	56.4	0.09	53.8	0.08
Larynx	40,378	5.9	0.03	5.7	0.03	10,553	1.4	0.01	1.3	0.01
Lung & bronchus	472,759	72.7	0.11	68.7	0.10	426,857	54.4	0.08	51.9	0.08
Bones & Joints	6,872	1.1	0.01	1.1	0.01	5,413	0.9	0.01	0.8	0.01
Soft Tissue (incl. heart)	24,929	4.0	0.03	3.8	0.02	19,122	2.7	0.02	2.7	0.02
Skin (excl. basal & sq.)	205,618	32.1	0.07	30.5	0.07	141,052	20.0	0.05	19.7	0.06
Melanomas of the skin	190,947	29.7	0.07	28.3	0.07	132,012	18.8	0.05	18.6	0.05
Breast (excl. *in situ*)	8,197	1.3	0.01	1.2	0.01	919,063	124.5	0.13	121.2	0.13
Female Genital System	-	-	-	-	-	370,689	50.0	0.08	49.0	0.08
Cervix uteri	-	-	-	-	-	46,054	7.3	0.04	7.5	0.04
Uterus Corpus	-	-	-	-	-	195,873	25.5	0.06	25.0	0.06
Uterus, NOS[1]	-	-	-	-	-	5,796	0.7	0.01	0.7	0.01
Ovary	-	-	-	-	-	88,008	11.8	0.04	11.4	0.04
Male Genital System	778,905	113.0	0.13	110.7	0.13	-	-	-	-	-
Prostate	735,242	105.5	0.13	102.8	0.12	-	-	-	-	-
Testis	36,726	6.4	0.03	6.8	0.04	-	-	-	-	-
Penis	5,531	0.9	0.01	0.8	0.01	-	-	-	-	-
Urinary System	393,862	61.3	0.10	57.7	0.09	163,874	21.2	0.05	20.3	0.05
Bladder (incl. *in situ*)	240,261	38.0	0.08	35.2	0.07	74,103	9.3	0.03	8.7	0.03
Kidney & renal pelvis	145,125	21.9	0.06	21.2	0.06	85,287	11.4	0.04	11.1	0.04
Ureter	5,509	0.9	0.01	0.8	0.01	3,222	0.4	0.01	0.4	0.01
Eye & Orbit	6,794	1.1	0.01	1.0	0.01	5,717	0.8	0.01	0.8	0.01
Brain & Other Nervous System	52,376	8.3	0.04	8.1	0.04	41,495	6.0	0.03	5.9	0.03
Brain	49,683	7.8	0.04	7.7	0.04	38,641	5.5	0.03	5.4	0.03
Endocrine System	53,202	8.4	0.04	8.3	0.04	144,903	22.8	0.06	23.2	0.06
Thyroid gland	48,427	7.6	0.04	7.6	0.04	140,681	22.1	0.06	22.6	0.06
Lymphomas	169,252	26.7	0.07	25.5	0.06	139,101	18.7	0.05	18.0	0.05
Hodgkin Lymphoma	18,811	3.1	0.02	3.2	0.02	15,072	2.5	0.02	2.5	0.02
Non-Hodgkin Lymphoma	150,441	23.5	0.06	22.4	0.06	124,029	16.2	0.05	15.5	0.05
Multiple Myeloma	47,663	7.4	0.03	6.9	0.03	35,869	4.6	0.02	4.3	0.02
Leukemias	113,460	18.1	0.05	17.1	0.05	81,458	11.0	0.04	10.5	0.04
Acute lymphocytic	11,199	2.0	0.02	1.9	0.02	8,761	1.5	0.02	1.5	0.02
Chronic lymphocytic	41,273	6.4	0.03	6.0	0.03	26,870	3.3	0.02	3.1	0.02
Acute myeloid	33,106	5.3	0.03	5.0	0.03	26,249	3.6	0.02	3.4	0.02
Chronic myeloid	14,120	2.3	0.02	2.2	0.02	10,230	1.4	0.01	1.3	0.01
Other leukemias	13,762	2.2	0.02	2.1	0.02	9,348	1.2	0.01	1.1	0.01
Mesothelioma	11,222	1.8	0.02	1.7	0.02	3,476	0.4	0.01	0.4	0.01
Kaposi Sarcoma	3,109	0.5	0.01	0.5	0.01	420	0.1	0.00	0.0	0.00
Ill-defined & Unknown	128,087	20.5	0.06	18.8	0.05	113,625	14.2	0.04	13.1	0.04
Selected tumors (non-invasive)										
Brain & ONS[1] (*benign/borderline*)	53,052	8.4	0.04	8.1	0.04	106,108	14.6	0.05	14.2	0.05
Breast (*in situ*)	856	0.1	0.00	0.1	0.00	219,011	30.3	0.07	29.7	0.07

* Rates are per 100,000 population and are age-adjusted by five-year age groups to the 2000 U.S. standard population based on single years of age and to the 1996 Canadian population standard.

\- Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for All Sites combined.

[1] Abbreviations: NOS - Not Otherwise Specified; ONS - Other Nervous System

Total Cancer Cases and Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates[1] for the Five Most Commonly Diagnosed Cancers
Selected Areas in the NAACCR United States Combined[2], 2010-2014, White, Males

	All Sites			Prostate			Lung & Bronchus			Colon & Rectum			Bladder			Melanoma of the Skin		
	Count	*Rate*	*S.E.*[3]	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
NAACCR U.S. Combined	**3,265,680**	**497.5**	**0.28**	**735,242**	**105.5**	**0.13**	**472,759**	**72.7**	**0.11**	**289,746**	**44.8**	**0.08**	**240,261**	**38.0**	**0.08**	**190,947**	**29.7**	**0.07**
Alabama	51,642	517.8	2.34	11,431	107.3	1.03	9,280	92.2	0.98	4,866	49.6	0.73	3,577	37.2	0.64	3,275	34.1	0.61
Alaska	5,415	440.1	6.77	1,288	94.3	2.97	692	61.1	2.62	461	39.0	2.07	394	36.8	2.07	239	18.8	1.34
Arizona	66,555	413.2	1.63	13,279	77.8	0.68	9,274	56.9	0.60	6,171	38.9	0.50	5,345	33.4	0.46	4,112	26.1	0.41
Arkansas	35,733	510.8	2.76	7,998	107.7	1.22	6,865	96.7	1.19	3,339	48.5	0.86	2,544	37.4	0.75	1,832	27.3	0.65
California	325,476	470.1	0.84	76,018	105.4	0.39	33,462	50.8	0.28	28,912	42.0	0.25	22,430	34.5	0.23	23,188	33.8	0.23
CA: Greater California	197,609	474.9	1.09	45,347	104.3	0.50	21,845	54.6	0.38	17,503	42.3	0.33	13,706	34.9	0.30	14,427	35.1	0.30
CA: Greater Bay Area	58,344	494.8	2.10	14,812	119.4	1.01	5,124	46.2	0.66	4,657	39.7	0.60	3,988	35.9	0.58	5,104	43.8	0.63
CA: Los Angeles	69,366	437.9	1.70	15,809	97.6	0.80	6,476	43.7	0.56	6,743	42.7	0.53	4,731	32.3	0.48	3,653	23.2	0.39
Colorado	49,927	434.6	2.02	12,129	98.6	0.93	5,183	47.9	0.69	4,217	37.6	0.60	3,447	32.6	0.58	3,391	29.6	0.53
Connecticut	45,050	523.8	2.53	10,261	111.7	1.13	5,968	69.9	0.92	3,709	43.8	0.74	4,125	48.9	0.78	2,456	29.3	0.61
Delaware	11,741	572.1	5.40	2,837	128.1	2.45	1,699	82.2	2.03	863	43.0	1.49	924	45.8	1.53	959	47.9	1.59
District of Columbia	2,093	380.6	8.76	539	97.6	4.35	173	35.6	2.81	148	26.6	2.30	124	24.4	2.28	122	22.2	2.12
Florida	249,665	476.2	0.97	52,312	94.6	0.42	39,120	72.2	0.37	21,501	41.3	0.29	18,793	34.7	0.26	16,379	32.0	0.26
Georgia	85,652	527.0	1.86	18,972	109.0	0.82	13,582	85.4	0.76	7,439	46.2	0.55	5,651	37.2	0.51	7,091	44.2	0.54
Hawaii	6,123	519.7	7.07	1,145	88.4	2.78	635	54.2	2.28	502	42.1	2.00	422	38.3	1.96	875	77.1	2.76
Idaho	18,766	470.5	3.52	4,820	114.2	1.68	2,159	55.4	1.22	1,547	39.2	1.02	1,437	37.7	1.02	1,303	33.0	0.94
Illinois	136,045	514.8	1.43	30,909	110.5	0.64	20,202	78.2	0.56	13,241	50.7	0.45	10,174	40.3	0.41	6,768	26.0	0.32
Indiana	74,239	489.3	1.84	14,560	90.3	0.77	13,369	88.9	0.79	7,241	48.4	0.58	5,443	37.4	0.52	3,328	22.3	0.40
Iowa	43,350	526.4	2.58	9,613	109.9	1.14	6,437	78.1	0.99	4,177	51.6	0.81	3,114	38.3	0.70	2,325	29.2	0.62
Kansas	34,756	508.0	2.78	8,261	115.3	1.29	4,864	71.7	1.05	3,125	46.3	0.84	2,608	39.1	0.78	1,979	29.8	0.68
Kentucky	62,328	586.3	2.42	11,986	105.6	0.99	12,432	116.9	1.08	6,202	59.1	0.77	4,125	41.1	0.66	3,339	32.4	0.58
Louisiana	46,473	563.5	2.68	11,177	127.1	1.23	6,992	85.5	1.05	4,295	52.5	0.82	3,017	38.1	0.71	2,436	30.6	0.64
Maine	20,908	516.7	3.68	4,345	98.1	1.53	3,441	84.7	1.48	1,694	42.8	1.07	1,869	47.6	1.12	1,106	28.0	0.87
Maryland	50,790	499.2	2.27	12,310	112.6	1.04	6,628	66.6	0.84	4,077	40.8	0.65	4,070	41.8	0.67	3,911	39.0	0.64
Massachusetts	78,379	499.9	1.83	17,890	106.0	0.81	10,935	71.1	0.69	6,549	42.6	0.54	6,383	42.1	0.54	4,072	26.6	0.43
Michigan	115,827	496.9	1.50	27,917	111.6	0.68	17,453	75.2	0.58	9,560	42.0	0.44	9,118	40.6	0.43	5,473	24.3	0.34
MI: Metropolitan Detroit	43,185	572.8	2.83	11,099	138.6	1.35	6,096	82.5	1.08	3,393	45.6	0.80	3,468	47.5	0.83	1,978	26.8	0.62
Mississippi	28,608	538.6	3.26	6,278	110.4	1.42	5,169	96.9	1.38	2,787	53.2	1.03	1,826	35.6	0.85	1,633	31.8	0.81
Missouri	72,020	495.6	1.88	14,865	95.4	0.80	12,714	87.7	0.79	6,934	48.8	0.60	4,929	35.1	0.51	3,766	26.6	0.44

Total Cancer Cases and Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates[1] for the Five Most Commonly Diagnosed Cancers
Selected Areas in the NAACCR United States Combined[2], 2010-2014, White, Males

	All Sites			Prostate			Lung & Bronchus			Colon & Rectum			Bladder			Melanoma of the Skin		
	Count	*Rate*	*S.E.*[3]	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
Montana	13,880	477.4	4.19	3,598	113.5	1.94	1,679	58.8	1.47	1,248	43.1	1.26	1,017	35.9	1.15	853	30.2	1.08
Nebraska	22,810	500.2	3.37	5,514	115.5	1.58	3,199	71.1	1.28	2,223	49.0	1.06	1,661	37.7	0.94	1,055	23.3	0.73
New Hampshire	19,244	525.2	3.92	4,909	121.6	1.79	2,539	71.5	1.46	1,461	40.7	1.10	1,681	48.2	1.21	1,149	32.3	0.99
New Jersey	101,292	549.5	1.76	25,269	130.0	0.83	12,201	67.2	0.62	8,838	48.4	0.52	8,005	44.8	0.51	6,053	33.2	0.44
New York	216,063	554.0	1.21	51,944	126.4	0.56	27,825	71.9	0.44	17,872	46.5	0.35	17,475	46.0	0.35	10,686	27.9	0.27
North Carolina	101,152	513.4	1.65	22,783	107.9	0.73	16,911	86.3	0.68	7,997	41.4	0.47	7,216	38.2	0.46	7,191	37.6	0.45
North Dakota	9,219	507.4	5.40	2,357	123.3	2.59	1,238	69.3	2.00	949	53.5	1.78	667	37.3	1.47	461	25.9	1.24
Ohio	136,148	491.0	1.36	29,179	98.3	0.59	22,969	83.0	0.56	12,822	47.1	0.43	10,594	39.5	0.39	7,012	26.1	0.32
Oklahoma	40,691	479.9	2.43	8,690	97.1	1.06	7,147	83.8	1.01	3,827	45.9	0.76	2,839	34.4	0.66	2,071	25.1	0.56
Oregon	47,711	465.7	2.19	10,983	99.5	0.97	6,278	62.3	0.80	4,003	40.2	0.65	3,765	38.1	0.64	3,267	32.6	0.59
Pennsylvania	172,302	530.4	1.30	36,818	106.0	0.56	25,243	77.2	0.49	15,804	49.3	0.40	14,561	45.4	0.38	9,156	28.9	0.31
Rhode Island	13,456	517.2	4.57	2,837	102.3	1.97	2,029	78.8	1.79	1,052	41.1	1.29	1,199	46.7	1.38	753	29.7	1.11
South Carolina	49,089	507.4	2.36	10,631	101.1	1.01	8,286	84.8	0.96	3,996	42.2	0.69	3,626	39.4	0.67	3,393	35.9	0.64
South Dakota	10,720	498.1	4.91	2,680	117.4	2.31	1,459	67.9	1.80	1,079	50.7	1.58	756	36.0	1.33	553	26.4	1.15
Tennessee	77,057	524.5	1.94	17,485	110.8	0.86	14,007	95.2	0.83	6,513	45.5	0.58	5,152	36.6	0.53	4,166	29.4	0.47
Texas	224,338	463.5	1.01	47,193	93.8	0.44	30,883	66.8	0.39	21,976	45.6	0.32	12,834	28.8	0.26	9,431	19.7	0.21
Utah	24,457	458.2	3.00	7,076	130.0	1.58	1,631	32.3	0.82	1,840	35.2	0.84	1,492	30.5	0.80	2,585	48.8	0.98
Vermont	8,917	489.7	5.38	1,940	97.2	2.28	1,291	71.0	2.04	706	39.1	1.53	685	39.0	1.53	702	39.4	1.55
Virginia	70,955	445.2	1.72	15,834	92.1	0.75	11,222	71.9	0.70	6,215	39.4	0.51	4,984	33.3	0.48	4,263	27.3	0.43
Washington	79,228	503.8	1.84	19,294	114.4	0.85	10,010	65.8	0.68	6,233	40.1	0.52	5,771	38.8	0.52	5,115	32.8	0.47
WA: Seattle/Puget Sound	54,837	535.5	2.36	13,649	124.6	1.10	6,542	66.6	0.85	4,175	41.0	0.65	3,947	41.1	0.67	3,854	37.9	0.63
West Virginia	28,458	523.3	3.20	5,725	97.7	1.32	5,471	99.0	1.38	2,820	52.8	1.02	2,134	40.5	0.90	1,339	26.0	0.73
Wisconsin	74,129	505.0	1.90	17,573	111.8	0.86	9,791	67.2	0.69	6,101	42.4	0.55	5,738	40.6	0.55	3,912	27.1	0.44
Wyoming	6,803	452.4	5.69	1,790	110.1	2.70	722	49.8	1.92	614	41.8	1.75	520	36.8	1.67	423	29.4	1.48

1. Rates are per 100,000 population and are age-adjusted by five-year age groups to the 2000 U.S. standard population based on single years of age.
2. See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.
3. Abbreviations: S.E. - Standard Error

\- Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for the U.S. combined.

Total Cancer Cases and Average Annual Age-adjusted (1996 Canadian Standard) Cancer Incidence Rates[1] for the Five Most Commonly Diagnosed Cancers
Selected Areas in the NAACCR United States Combined[2], 2010-2014, White, Males

	All Sites			Prostate			Lung & Bronchus			Colon & Rectum			Bladder			Melanoma of the Skin		
	Count	*Rate*	*S.E.*[3]	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
NAACCR U.S. Combined	**3,265,680**	**475.1**	**0.27**	**735,242**	**102.8**	**0.12**	**472,759**	**68.7**	**0.10**	**289,746**	**42.3**	**0.08**	**240,261**	**35.2**	**0.07**	**190,947**	**28.3**	**0.07**
Alabama	51,642	494.9	2.21	11,431	104.1	0.98	9,280	87.5	0.92	4,866	47.0	0.69	3,577	34.5	0.58	3,275	32.6	0.58
Alaska	5,415	419.8	6.22	1,288	91.7	2.79	692	57.6	2.38	461	36.4	1.86	394	34.5	1.88	239	18.3	1.27
Arizona	66,555	393.9	1.54	13,279	75.5	0.66	9,274	53.5	0.56	6,171	36.8	0.47	5,345	30.9	0.42	4,112	24.7	0.39
Arkansas	35,733	489.3	2.62	7,998	105.0	1.18	6,865	92.1	1.12	3,339	46.1	0.81	2,544	34.8	0.69	1,832	25.9	0.62
California	325,476	448.1	0.80	76,018	102.9	0.38	33,462	47.5	0.26	28,912	39.6	0.24	22,430	31.7	0.21	23,188	32.1	0.21
CA: Greater California	197,609	452.6	1.03	45,347	101.6	0.48	21,845	51.2	0.35	17,503	39.9	0.31	13,706	32.1	0.28	14,427	33.2	0.28
CA: Greater Bay Area	58,344	471.5	1.99	14,812	116.5	0.98	5,124	43.0	0.61	4,657	37.3	0.56	3,988	32.9	0.53	5,104	41.6	0.59
CA: Los Angeles	69,366	418.1	1.62	15,809	95.4	0.77	6,476	41.0	0.52	6,743	40.4	0.50	4,731	29.7	0.44	3,653	22.0	0.37
Colorado	49,927	414.3	1.90	12,129	96.0	0.89	5,183	44.8	0.64	4,217	35.3	0.56	3,447	30.1	0.52	3,391	28.4	0.50
Connecticut	45,050	499.8	2.42	10,261	109.4	1.10	5,968	65.8	0.87	3,709	41.0	0.69	4,125	45.3	0.72	2,456	27.8	0.58
Delaware	11,741	545.0	5.13	2,837	124.8	2.36	1,699	77.3	1.89	863	40.3	1.40	924	42.0	1.39	959	45.7	1.52
District of Columbia	2,093	364.4	8.24	539	95.0	4.18	173	33.5	2.60	148	24.9	2.12	124	22.7	2.09	122	21.0	1.97
Florida	249,665	456.2	0.94	52,312	92.7	0.41	39,120	68.4	0.35	21,501	39.2	0.27	18,793	32.2	0.24	16,379	30.4	0.25
Georgia	85,652	503.8	1.75	18,972	106.4	0.79	13,582	80.8	0.70	7,439	43.7	0.52	5,651	34.6	0.47	7,091	42.3	0.51
Hawaii	6,123	496.9	6.67	1,145	85.6	2.63	635	50.9	2.10	502	40.3	1.89	422	35.3	1.78	875	73.5	2.61
Idaho	18,766	448.2	3.32	4,820	110.4	1.61	2,159	51.8	1.13	1,547	37.1	0.96	1,437	34.9	0.93	1,303	31.8	0.90
Illinois	136,045	491.2	1.36	30,909	107.8	0.62	20,202	73.7	0.53	13,241	47.8	0.42	10,174	37.4	0.38	6,768	24.8	0.31
Indiana	74,239	467.2	1.75	14,560	87.7	0.74	13,369	84.4	0.74	7,241	45.7	0.55	5,443	34.7	0.48	3,328	21.3	0.38
Iowa	43,350	502.2	2.46	9,613	107.0	1.11	6,437	73.8	0.93	4,177	48.4	0.77	3,114	35.5	0.65	2,325	28.0	0.60
Kansas	34,756	485.9	2.66	8,261	112.3	1.25	4,864	67.8	0.99	3,125	43.8	0.80	2,608	36.3	0.72	1,979	28.5	0.66
Kentucky	62,328	560.2	2.29	11,986	102.6	0.95	12,432	110.9	1.01	6,202	56.0	0.72	4,125	38.1	0.60	3,339	31.0	0.55
Louisiana	46,473	540.0	2.55	11,177	124.3	1.19	6,992	81.1	0.98	4,295	49.9	0.77	3,017	35.6	0.66	2,436	29.2	0.60
Maine	20,908	492.6	3.50	4,345	95.6	1.47	3,441	79.9	1.38	1,694	40.3	1.01	1,869	44.2	1.04	1,106	26.7	0.83
Maryland	50,790	476.4	2.16	12,310	110.4	1.01	6,628	62.7	0.78	4,077	38.4	0.61	4,070	38.5	0.61	3,911	37.3	0.61
Massachusetts	78,379	477.9	1.75	17,890	103.8	0.79	10,935	66.9	0.65	6,549	40.0	0.50	6,383	39.1	0.50	4,072	25.4	0.41
Michigan	115,827	474.4	1.42	27,917	108.9	0.66	17,453	71.0	0.54	9,560	39.5	0.41	9,118	37.6	0.40	5,473	23.1	0.32
MI: Metropolitan Detroit	43,185	546.5	2.70	11,099	135.4	1.31	6,096	77.6	1.01	3,393	42.8	0.75	3,468	44.1	0.76	1,978	25.5	0.59
Mississippi	28,608	516.4	3.10	6,278	108.0	1.37	5,169	91.9	1.29	2,787	50.6	0.98	1,826	33.2	0.78	1,633	30.5	0.77
Missouri	72,020	474.1	1.80	14,865	93.3	0.77	12,714	83.1	0.74	6,934	46.0	0.56	4,929	32.7	0.47	3,766	25.5	0.42

Total Cancer Cases and Average Annual Age-adjusted (1996 Canadian Standard) Cancer Incidence Rates[1] for the Five Most Commonly Diagnosed Cancers
Selected Areas in the NAACCR United States Combined[2], 2010-2014, White, Males

	All Sites			Prostate			Lung & Bronchus			Colon & Rectum			Bladder			Melanoma of the Skin		
	Count	*Rate*	*S.E.*[3]	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
Montana	13,880	457.1	3.99	3,598	110.5	1.87	1,679	55.0	1.36	1,248	41.2	1.20	1,017	33.2	1.06	853	29.2	1.05
Nebraska	22,810	478.4	3.23	5,514	112.2	1.53	3,199	67.3	1.21	2,223	46.3	1.00	1,661	35.0	0.87	1,055	22.6	0.71
New Hampshire	19,244	502.3	3.73	4,909	119.2	1.74	2,539	67.0	1.36	1,461	38.2	1.03	1,681	44.8	1.12	1,149	30.7	0.93
New Jersey	101,292	524.3	1.68	25,269	126.8	0.81	12,201	63.3	0.58	8,838	45.5	0.49	8,005	41.4	0.47	6,053	31.6	0.42
New York	216,063	529.0	1.16	51,944	123.3	0.55	27,825	67.8	0.41	17,872	43.7	0.33	17,475	42.6	0.33	10,686	26.6	0.26
North Carolina	101,152	490.8	1.57	22,783	105.0	0.70	16,911	81.7	0.63	7,997	39.2	0.44	7,216	35.5	0.42	7,191	36.0	0.43
North Dakota	9,219	485.8	5.19	2,357	120.1	2.52	1,238	64.9	1.88	949	51.0	1.70	667	34.4	1.36	461	25.3	1.22
Ohio	136,148	469.4	1.30	29,179	96.0	0.57	22,969	78.5	0.53	12,822	44.5	0.40	10,594	36.7	0.36	7,012	25.0	0.31
Oklahoma	40,691	458.6	2.31	8,690	94.1	1.02	7,147	79.5	0.95	3,827	43.6	0.72	2,839	32.1	0.61	2,071	23.9	0.53
Oregon	47,711	444.9	2.08	10,983	97.1	0.94	6,278	58.5	0.75	4,003	37.8	0.61	3,765	35.2	0.58	3,267	31.3	0.56
Pennsylvania	172,302	506.6	1.25	36,818	103.5	0.55	25,243	72.9	0.47	15,804	46.4	0.38	14,561	42.1	0.35	9,156	27.6	0.30
Rhode Island	13,456	493.4	4.37	2,837	100.1	1.92	2,029	74.1	1.68	1,052	38.3	1.21	1,199	43.3	1.28	753	28.5	1.07
South Carolina	49,089	483.5	2.22	10,631	98.3	0.96	8,286	80.1	0.89	3,996	39.9	0.64	3,626	36.3	0.61	3,393	34.4	0.60
South Dakota	10,720	476.0	4.71	2,680	114.6	2.25	1,459	63.4	1.68	1,079	47.8	1.49	756	33.3	1.23	553	25.7	1.13
Tennessee	77,057	502.6	1.84	17,485	108.1	0.83	14,007	90.4	0.77	6,513	43.2	0.54	5,152	34.1	0.48	4,166	28.1	0.44
Texas	224,338	441.9	0.95	47,193	90.8	0.43	30,883	62.9	0.36	21,976	43.2	0.30	12,834	26.6	0.24	9,431	18.8	0.20
Utah	24,457	439.2	2.85	7,076	126.6	1.53	1,631	30.5	0.76	1,840	33.4	0.79	1,492	28.2	0.74	2,585	46.6	0.93
Vermont	8,917	468.6	5.12	1,940	94.6	2.19	1,291	67.2	1.91	706	36.8	1.43	685	36.5	1.42	702	37.7	1.47
Virginia	70,955	424.4	1.62	15,834	89.9	0.73	11,222	67.7	0.65	6,215	37.3	0.48	4,984	30.7	0.44	4,263	25.9	0.40
Washington	79,228	480.1	1.74	19,294	111.5	0.82	10,010	61.6	0.63	6,233	37.7	0.49	5,771	35.7	0.48	5,115	31.6	0.45
WA: Seattle/Puget Sound	54,837	509.9	2.23	13,649	121.4	1.06	6,542	62.3	0.78	4,175	38.6	0.61	3,947	37.6	0.61	3,854	36.4	0.60
West Virginia	28,458	500.5	3.04	5,725	94.8	1.27	5,471	94.2	1.29	2,820	50.1	0.96	2,134	37.7	0.83	1,339	24.9	0.70
Wisconsin	74,129	482.2	1.81	17,573	109.1	0.84	9,791	63.4	0.65	6,101	39.8	0.52	5,738	37.5	0.50	3,912	26.0	0.43
Wyoming	6,803	434.1	5.41	1,790	107.8	2.61	722	46.8	1.78	614	39.7	1.65	520	34.5	1.54	423	28.2	1.41

1. Rates are per 100,000 and are age-adjusted to the 1996 Canadian population standard.
2. See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.
3. Abbreviations: S.E. - Standard Error

\- Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for the U.S. combined.

Total Cancer Cases and Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates[1] for the Five Most Commonly Diagnosed Cancers
Selected Areas in the NAACCR United States Combined[2], 2010-2014, White, Females

	All Sites			Breast			Lung & Bronchus			Colon & Rectum			Uterus Corpus & Uterus, NOS[3]			Thyroid		
	Count	*Rate*	*S.E.*[3]	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
NAACCR U.S. Combined	**3,168,903**	**423.8**	**0.24**	**919,063**	**124.5**	**0.13**	**426,857**	**54.4**	**0.08**	**264,042**	**34.0**	**0.07**	**201,669**	**26.2**	**0.06**	**140,681**	**22.1**	**0.06**
Alabama	45,391	400.0	1.94	13,184	117.1	1.05	6,850	56.3	0.69	4,052	34.7	0.56	2,187	18.8	0.41	1,331	14.2	0.40
Alaska	4,827	403.7	6.19	1,558	124.9	3.37	573	52.6	2.35	392	33.3	1.79	348	26.4	1.51	200	15.6	1.14
Arizona	65,962	377.9	1.51	19,729	114.4	0.84	8,716	46.4	0.50	5,372	29.7	0.42	3,778	21.1	0.35	3,520	24.2	0.42
Arkansas	30,737	394.7	2.32	8,495	110.1	1.23	5,180	62.0	0.88	2,858	35.4	0.68	1,687	21.5	0.54	1,014	15.9	0.52
California	321,201	404.6	0.73	98,648	125.3	0.41	33,786	42.0	0.23	26,344	32.4	0.20	20,725	25.5	0.18	14,394	19.7	0.17
CA: Greater California	192,617	407.8	0.95	59,165	126.2	0.53	21,946	45.5	0.31	15,712	32.5	0.26	11,829	24.4	0.23	8,449	19.8	0.22
CA: Greater Bay Area	56,383	423.5	1.83	17,509	133.3	1.03	5,664	41.6	0.57	4,489	32.7	0.50	3,748	27.2	0.45	2,112	17.6	0.39
CA: Los Angeles	72,089	382.5	1.44	21,949	117.1	0.80	6,166	33.0	0.43	6,134	31.9	0.41	5,148	26.9	0.38	3,832	20.8	0.34
Colorado	51,019	394.5	1.79	16,336	124.9	1.00	5,397	41.9	0.58	3,947	30.7	0.50	3,100	22.5	0.41	2,884	23.8	0.45
Connecticut	47,094	465.2	2.23	14,030	141.8	1.24	6,255	58.6	0.76	3,662	33.8	0.58	3,108	29.5	0.54	2,427	29.8	0.63
Delaware	10,845	469.0	4.70	3,091	134.4	2.52	1,626	65.3	1.65	788	32.6	1.20	714	29.4	1.14	444	24.9	1.24
District of Columbia	2,211	378.1	8.46	816	143.7	5.22	191	34.9	2.63	145	25.7	2.24	130	22.4	2.03	165	23.1	2.00
Florida	230,155	401.3	0.88	64,149	115.6	0.48	35,346	55.8	0.30	19,377	31.5	0.24	13,217	22.9	0.21	7,948	18.6	0.22
Georgia	78,573	422.4	1.54	23,143	123.9	0.83	10,946	56.7	0.55	6,273	33.3	0.43	4,112	21.2	0.34	3,212	19.2	0.35
Hawaii	4,842	461.5	7.08	1,523	145.2	3.98	488	44.8	2.13	380	36.0	1.96	275	24.0	1.54	184	20.7	1.62
Idaho	17,446	409.5	3.17	5,131	120.5	1.72	2,074	47.0	1.05	1,377	32.1	0.88	1,068	24.1	0.75	934	24.8	0.83
Illinois	136,743	442.4	1.23	39,801	130.9	0.67	18,722	58.6	0.44	11,910	36.7	0.34	9,343	29.5	0.31	6,171	23.1	0.30
Indiana	75,131	426.9	1.60	20,890	119.8	0.85	11,212	61.3	0.59	6,930	38.1	0.47	4,983	27.5	0.40	2,778	18.6	0.36
Iowa	41,119	440.6	2.26	11,233	123.8	1.22	5,236	53.0	0.75	4,009	39.8	0.65	2,954	31.2	0.59	1,513	20.3	0.54
Kansas	32,702	423.6	2.42	9,341	122.5	1.31	4,233	52.1	0.82	2,863	35.3	0.68	1,928	24.5	0.57	1,630	25.0	0.64
Kentucky	58,525	475.4	2.01	15,170	123.4	1.03	10,362	80.4	0.80	5,293	42.0	0.59	3,280	26.1	0.47	2,349	22.2	0.47
Louisiana	39,443	425.1	2.20	11,185	121.0	1.18	5,592	57.2	0.78	3,625	38.2	0.65	1,724	18.3	0.45	1,786	22.2	0.54
Maine	20,686	454.1	3.29	5,655	126.0	1.75	3,146	65.3	1.19	1,612	34.2	0.88	1,488	31.0	0.83	787	22.0	0.83
Maryland	50,740	434.2	1.99	15,248	131.4	1.10	6,781	55.6	0.69	3,831	31.6	0.53	3,299	27.1	0.48	2,419	24.7	0.52
Massachusetts	85,309	455.5	1.61	25,550	139.2	0.90	12,209	62.5	0.58	6,590	33.4	0.43	5,736	29.7	0.40	4,530	29.1	0.45
Michigan	110,879	422.7	1.31	31,272	120.8	0.71	16,130	58.3	0.47	8,910	32.5	0.35	7,957	29.3	0.34	4,451	20.8	0.33
MI: Metropolitan Detroit	41,541	468.4	2.38	11,601	132.5	1.27	6,128	66.0	0.86	3,322	35.3	0.63	2,897	31.9	0.61	1,965	26.9	0.63
Mississippi	24,435	411.3	2.72	6,624	112.5	1.43	3,868	60.6	0.99	2,318	37.3	0.80	1,117	18.7	0.58	884	18.3	0.64
Missouri	70,484	426.6	1.66	20,383	125.1	0.90	11,178	64.2	0.62	6,183	35.9	0.47	4,417	26.2	0.40	2,539	18.5	0.38

Total Cancer Cases and Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates[1] for the Five Most Commonly Diagnosed Cancers
Selected Areas in the NAACCR United States Combined[2], 2010-2014, White, Females

	All Sites			Breast			Lung & Bronchus			Colon & Rectum			Uterus Corpus & Uterus, NOS[3]			Thyroid		
	Count	*Rate*	*S.E.*[3]	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
Montana	12,598	419.4	3.90	3,674	122.9	2.12	1,660	52.0	1.30	1,026	33.0	1.07	805	25.4	0.93	523	21.8	1.00
Nebraska	21,321	414.3	2.93	6,197	122.3	1.60	2,683	49.8	0.98	2,051	37.8	0.86	1,406	27.0	0.74	897	21.0	0.72
New Hampshire	19,044	463.6	3.47	5,809	141.7	1.92	2,689	63.8	1.26	1,447	34.2	0.93	1,432	32.9	0.90	804	23.0	0.85
New Jersey	103,380	468.5	1.51	29,252	135.9	0.82	12,939	55.5	0.50	8,916	37.6	0.41	7,387	32.6	0.39	5,534	30.4	0.42
New York	218,024	472.3	1.05	60,373	134.4	0.56	28,724	58.9	0.35	17,454	35.6	0.28	15,073	32.0	0.27	11,540	30.2	0.29
North Carolina	96,327	423.8	1.40	28,898	128.0	0.77	14,046	58.5	0.50	7,335	31.6	0.38	5,696	24.2	0.33	3,771	19.4	0.33
North Dakota	8,055	410.7	4.78	2,281	121.5	2.67	969	46.8	1.54	823	39.2	1.42	502	25.2	1.16	398	24.6	1.28
Ohio	136,031	424.8	1.19	38,759	122.6	0.64	19,820	58.9	0.43	11,855	35.1	0.33	9,611	29.2	0.31	5,832	22.2	0.30
Oklahoma	38,075	398.8	2.10	10,765	113.6	1.13	5,755	56.9	0.76	3,438	34.7	0.61	2,021	20.8	0.48	1,414	17.7	0.49
Oregon	47,571	421.2	2.00	14,307	126.9	1.10	6,366	53.7	0.69	3,615	30.8	0.53	3,141	26.6	0.49	1,818	19.5	0.47
Pennsylvania	172,614	459.6	1.16	47,785	130.3	0.62	22,299	55.3	0.38	15,001	36.9	0.31	12,590	32.8	0.30	9,230	31.5	0.34
Rhode Island	14,656	466.8	4.03	4,066	133.5	2.18	2,155	64.9	1.44	1,185	34.6	1.06	1,054	32.6	1.04	723	28.4	1.10
South Carolina	45,440	420.9	2.04	13,779	127.8	1.13	6,664	57.7	0.72	3,606	32.6	0.56	2,449	21.9	0.46	1,709	19.1	0.48
South Dakota	9,947	428.2	4.50	2,951	131.0	2.52	1,222	49.7	1.46	970	38.1	1.28	676	28.8	1.15	386	20.7	1.10
Tennessee	70,715	422.8	1.63	20,059	120.3	0.87	11,096	62.8	0.60	6,085	35.4	0.46	3,959	23.2	0.38	2,852	20.1	0.39
Texas	214,004	383.8	0.84	62,605	112.2	0.45	25,241	45.0	0.29	17,579	31.3	0.24	12,211	21.3	0.20	10,162	19.0	0.19
Utah	22,134	375.7	2.56	6,730	115.9	1.43	1,375	23.8	0.65	1,583	27.3	0.69	1,545	25.6	0.66	1,781	29.5	0.71
Vermont	8,845	436.8	4.84	2,614	130.2	2.65	1,250	59.0	1.71	701	33.1	1.29	708	33.3	1.30	308	19.2	1.16
Virginia	72,949	401.9	1.52	23,025	127.3	0.86	9,913	53.0	0.54	5,839	31.5	0.42	4,663	24.5	0.37	3,238	20.5	0.37
Washington	77,210	444.9	1.65	23,797	137.1	0.92	9,767	54.9	0.57	5,767	32.8	0.44	4,782	26.0	0.39	3,089	20.4	0.38
WA: Seattle/Puget Sound	53,018	464.2	2.08	16,502	144.2	1.16	6,476	55.7	0.71	3,853	33.3	0.55	3,287	27.1	0.49	2,080	20.6	0.46
West Virginia	26,775	445.4	2.83	6,848	114.5	1.44	4,305	66.8	1.04	2,578	41.0	0.83	1,870	30.6	0.74	1,184	24.5	0.74
Wisconsin	70,744	432.2	1.68	20,523	127.9	0.92	9,116	53.4	0.57	5,673	32.8	0.45	5,071	29.7	0.43	2,695	19.9	0.40
Wyoming	5,945	386.8	5.18	1,781	115.0	2.82	706	44.6	1.71	474	30.6	1.44	342	21.5	1.20	299	21.9	1.32

1. Rates are per 100,000 population and are age-adjusted by five-year age groups to the 2000 U.S. standard population based on single years of age.
2. See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.
3. Abbreviations: NOS - Not Otherwise Specified; S.E. - Standard Error

\- Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for the U.S. combined.

Total Cancer Cases and Average Annual Age-adjusted (1996 Canadian Standard) Cancer Incidence Rates[1] for the Five Most Commonly Diagnosed Cancers
Selected Areas in the NAACCR United States Combined[2], 2010-2014, White, Females

	All Sites			Breast			Lung & Bronchus			Colon & Rectum			Uterus Corpus & Uterus, NOS[3]			Thyroid		
	Count	*Rate*	*S.E.*[3]	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
NAACCR U.S. Combined	**3,168,903**	**409.4**	**0.24**	**919,063**	**121.2**	**0.13**	**426,857**	**51.9**	**0.08**	**264,042**	**31.9**	**0.06**	**201,669**	**25.7**	**0.06**	**140,681**	**22.6**	**0.06**
Alabama	45,391	387.2	1.90	13,184	114.0	1.03	6,850	54.1	0.66	4,052	32.9	0.54	2,187	18.4	0.41	1,331	14.5	0.42
Alaska	4,827	389.5	5.91	1,558	121.0	3.24	573	49.9	2.19	392	31.6	1.68	348	25.9	1.47	200	16.1	1.18
Arizona	65,962	364.6	1.47	19,729	111.4	0.82	8,716	44.0	0.48	5,372	27.9	0.39	3,778	20.7	0.35	3,520	24.7	0.44
Arkansas	30,737	382.1	2.27	8,495	107.1	1.21	5,180	59.8	0.84	2,858	33.6	0.65	1,687	21.0	0.53	1,014	16.2	0.53
California	321,201	389.8	0.71	98,648	121.9	0.40	33,786	39.6	0.22	26,344	30.4	0.19	20,725	25.0	0.18	14,394	20.0	0.17
CA: Greater California	192,617	393.0	0.92	59,165	122.8	0.52	21,946	43.0	0.30	15,712	30.5	0.25	11,829	23.9	0.22	8,449	20.1	0.23
CA: Greater Bay Area	56,383	407.6	1.78	17,509	129.8	1.01	5,664	39.0	0.53	4,489	30.6	0.48	3,748	26.6	0.44	2,112	18.0	0.40
CA: Los Angeles	72,089	368.6	1.40	21,949	113.9	0.78	6,166	31.1	0.41	6,134	30.0	0.39	5,148	26.5	0.37	3,832	21.2	0.35
Colorado	51,019	380.3	1.73	16,336	121.5	0.98	5,397	39.5	0.55	3,947	28.7	0.47	3,100	22.0	0.41	2,884	24.2	0.46
Connecticut	47,094	449.4	2.20	14,030	138.2	1.23	6,255	55.8	0.73	3,662	31.6	0.56	3,108	29.0	0.54	2,427	30.4	0.66
Delaware	10,845	454.1	4.61	3,091	130.7	2.48	1,626	62.3	1.58	788	30.6	1.14	714	28.9	1.12	444	25.6	1.29
District of Columbia	2,211	366.4	8.11	816	140.6	5.06	191	32.8	2.46	145	24.4	2.11	130	21.8	1.96	165	23.6	1.99
Florida	230,155	387.6	0.87	64,149	112.5	0.47	35,346	53.2	0.29	19,377	29.7	0.23	13,217	22.4	0.20	7,948	19.0	0.23
Georgia	78,573	408.7	1.49	23,143	120.6	0.81	10,946	54.2	0.52	6,273	31.5	0.41	4,112	20.7	0.33	3,212	19.6	0.36
Hawaii	4,842	445.7	6.87	1,523	141.2	3.88	488	42.4	2.00	380	33.8	1.85	275	23.3	1.49	184	21.4	1.68
Idaho	17,446	395.2	3.08	5,131	117.0	1.68	2,074	44.7	1.00	1,377	30.1	0.84	1,068	23.6	0.74	934	25.4	0.86
Illinois	136,743	427.0	1.20	39,801	127.4	0.66	18,722	55.8	0.42	11,910	34.4	0.33	9,343	28.9	0.31	6,171	23.5	0.31
Indiana	75,131	413.0	1.56	20,890	116.9	0.84	11,212	58.8	0.57	6,930	35.8	0.45	4,983	27.0	0.39	2,778	19.0	0.38
Iowa	41,119	425.5	2.23	11,233	120.5	1.20	5,236	50.6	0.72	4,009	37.0	0.62	2,954	30.6	0.58	1,513	20.8	0.56
Kansas	32,702	410.1	2.38	9,341	119.3	1.29	4,233	49.8	0.79	2,863	33.4	0.65	1,928	24.0	0.57	1,630	25.4	0.66
Kentucky	58,525	460.4	1.97	15,170	120.4	1.01	10,362	77.4	0.77	5,293	39.7	0.56	3,280	25.6	0.46	2,349	22.7	0.49
Louisiana	39,443	411.6	2.15	11,185	117.8	1.15	5,592	54.8	0.75	3,625	36.3	0.62	1,724	17.9	0.45	1,786	22.4	0.55
Maine	20,686	438.8	3.23	5,655	122.9	1.72	3,146	62.3	1.14	1,612	32.1	0.85	1,488	30.4	0.82	787	22.5	0.87
Maryland	50,740	419.5	1.95	15,248	127.9	1.08	6,781	52.8	0.66	3,831	29.8	0.50	3,299	26.6	0.48	2,419	25.2	0.54
Massachusetts	85,309	440.0	1.58	25,550	135.6	0.89	12,209	59.5	0.55	6,590	31.3	0.41	5,736	29.1	0.40	4,530	29.7	0.46
Michigan	110,879	408.8	1.29	31,272	117.7	0.70	16,130	55.6	0.45	8,910	30.5	0.34	7,957	28.8	0.34	4,451	21.3	0.34
MI: Metropolitan Detroit	41,541	452.2	2.34	11,601	128.9	1.25	6,128	62.8	0.83	3,322	33.0	0.60	2,897	31.3	0.60	1,965	27.4	0.65
Mississippi	24,435	398.6	2.67	6,624	109.6	1.41	3,868	58.4	0.96	2,318	35.5	0.77	1,117	18.3	0.57	884	18.6	0.66
Missouri	70,484	412.4	1.62	20,383	121.9	0.89	11,178	61.5	0.59	6,183	33.7	0.45	4,417	25.7	0.40	2,539	18.8	0.39

Total Cancer Cases and Average Annual Age-adjusted (1996 Canadian Standard) Cancer Incidence Rates[1] for the Five Most Commonly Diagnosed Cancers
Selected Areas in the NAACCR United States Combined[2], 2010-2014, White, Females

	All Sites			Breast			Lung & Bronchus			Colon & Rectum			Uterus Corpus & Uterus, NOS[3]			Thyroid		
	Count	*Rate*	*S.E.*[3]	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
Montana	12,598	405.5	3.82	3,674	119.5	2.08	1,660	49.3	1.24	1,026	31.0	1.02	805	24.8	0.92	523	22.3	1.04
Nebraska	21,321	400.6	2.88	6,197	119.2	1.58	2,683	47.7	0.95	2,051	35.4	0.82	1,406	26.5	0.73	897	21.5	0.75
New Hampshire	19,044	448.5	3.40	5,809	138.2	1.89	2,689	60.9	1.20	1,447	31.9	0.88	1,432	32.4	0.89	804	23.5	0.88
New Jersey	103,380	452.5	1.48	29,252	132.4	0.81	12,939	52.9	0.48	8,916	35.2	0.39	7,387	32.0	0.38	5,534	30.9	0.44
New York	218,024	456.0	1.02	60,373	130.9	0.56	28,724	56.2	0.34	17,454	33.3	0.27	15,073	31.3	0.26	11,540	30.7	0.30
North Carolina	96,327	410.0	1.37	28,898	124.7	0.76	14,046	56.0	0.48	7,335	29.9	0.36	5,696	23.7	0.32	3,771	19.8	0.34
North Dakota	8,055	398.2	4.73	2,281	118.3	2.63	969	45.0	1.49	823	37.0	1.37	502	24.7	1.15	398	25.1	1.33
Ohio	136,031	411.1	1.17	38,759	119.3	0.63	19,820	56.3	0.41	11,855	32.9	0.32	9,611	28.6	0.30	5,832	22.8	0.31
Oklahoma	38,075	386.0	2.06	10,765	110.5	1.11	5,755	54.6	0.73	3,438	32.9	0.58	2,021	20.4	0.47	1,414	18.0	0.50
Oregon	47,571	406.9	1.95	14,307	123.5	1.08	6,366	50.9	0.65	3,615	28.9	0.50	3,141	26.1	0.48	1,818	19.9	0.49
Pennsylvania	172,614	444.5	1.14	47,785	126.9	0.61	22,299	52.8	0.37	15,001	34.6	0.30	12,590	32.1	0.30	9,230	32.1	0.36
Rhode Island	14,656	452.2	3.98	4,066	130.3	2.16	2,155	61.9	1.38	1,185	32.4	1.02	1,054	31.9	1.03	723	29.0	1.14
South Carolina	45,440	407.0	1.99	13,779	124.3	1.10	6,664	55.2	0.69	3,606	30.8	0.53	2,449	21.6	0.45	1,709	19.6	0.50
South Dakota	9,947	413.5	4.43	2,951	127.6	2.49	1,222	47.4	1.41	970	35.6	1.23	676	28.1	1.13	386	21.1	1.14
Tennessee	70,715	409.7	1.60	20,059	117.4	0.86	11,096	60.4	0.58	6,085	33.6	0.44	3,959	22.7	0.37	2,852	20.6	0.40
Texas	214,004	370.6	0.82	62,605	109.3	0.44	25,241	42.7	0.27	17,579	29.5	0.23	12,211	21.0	0.19	10,162	19.4	0.20
Utah	22,134	364.1	2.48	6,730	112.6	1.39	1,375	22.7	0.62	1,583	25.8	0.66	1,545	25.1	0.65	1,781	30.2	0.73
Vermont	8,845	422.1	4.74	2,614	127.0	2.61	1,250	56.3	1.64	701	30.6	1.21	708	32.5	1.27	308	19.9	1.22
Virginia	72,949	387.9	1.48	23,025	123.8	0.84	9,913	50.3	0.51	5,839	29.6	0.40	4,663	24.0	0.36	3,238	20.9	0.38
Washington	77,210	428.8	1.60	23,797	133.3	0.89	9,767	51.8	0.54	5,767	30.8	0.42	4,782	25.4	0.38	3,089	20.7	0.39
WA: Seattle/Puget Sound	53,018	447.2	2.01	16,502	140.3	1.13	6,476	52.5	0.67	3,853	31.3	0.52	3,287	26.4	0.47	2,080	20.9	0.47
West Virginia	26,775	432.0	2.78	6,848	111.6	1.42	4,305	64.0	1.00	2,578	38.6	0.80	1,870	30.3	0.74	1,184	25.1	0.78
Wisconsin	70,744	417.7	1.65	20,523	124.6	0.91	9,116	51.1	0.55	5,673	30.5	0.43	5,071	29.1	0.42	2,695	20.4	0.41
Wyoming	5,945	375.2	5.06	1,781	111.9	2.76	706	42.4	1.63	474	29.1	1.38	342	21.1	1.19	299	22.5	1.36

1. Rates are per 100,000 and are age-adjusted to the 1996 Canadian population standard.
2. See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.
3. Abbreviations: NOS - Not Otherwise Specified; S.E. - Standard Error

\- Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for the U.S. combined.

NAACCR U.S. Combined Incidence 2010-2014, The Black Population

Total Cancer Cases and Average Annual Age-adjusted Cancer Incidence Rates*
NAACCR United States Combined, 2010-2014, Black

	MALES (BLACK)					FEMALES (BLACK)				
SITES	*Count*	*Adj. Rate 2000 U.S.*	*S.E.*	*Adj. Rate 1996 Cdn*	*S.E.*	*Count*	*Adj. Rate 2000 U.S.*	*S.E.*	*Adj. Rate 1996 Cdn*	*S.E.*
All Sites (invasive & in situ bladder)	432,930	550.2	0.90	529.9	0.85	418,018	400.2	0.63	386.4	0.61
Oral Cavity & Pharynx	12,501	14.5	0.14	14.2	0.13	5,489	5.1	0.07	4.9	0.07
Lip	92	0.1	0.01	0.1	0.01	67	0.1	0.01	0.1	0.01
Tongue	2,875	3.3	0.07	3.2	0.06	1,160	1.1	0.03	1.0	0.03
Major salivary gland	960	1.2	0.04	1.1	0.04	1,113	1.0	0.03	1.0	0.03
Floor of mouth	743	0.9	0.03	0.8	0.03	302	0.3	0.02	0.3	0.02
Gum & other mouth	1,478	1.8	0.05	1.7	0.05	1,079	1.0	0.03	1.0	0.03
Nasopharynx	898	1.0	0.03	1.0	0.03	380	0.4	0.02	0.3	0.02
Oropharynx	999	1.2	0.04	1.1	0.04	318	0.3	0.02	0.3	0.02
Hypopharynx	1,513	1.8	0.05	1.8	0.05	303	0.3	0.02	0.3	0.02
Digestive System	94,069	121.0	0.42	115.5	0.40	82,760	80.5	0.29	76.3	0.27
Esophagus	5,558	7.1	0.10	6.8	0.10	2,380	2.3	0.05	2.2	0.05
Stomach	10,118	14.1	0.15	13.2	0.14	7,724	7.7	0.09	7.3	0.08
Small intestine	3,318	4.3	0.08	4.1	0.08	3,347	3.2	0.06	3.1	0.05
Colon excl. rectum	30,187	40.6	0.25	38.4	0.23	32,498	31.7	0.18	30.0	0.17
Rectum & rectosigmoid	11,921	14.7	0.14	14.1	0.14	9,803	9.2	0.09	8.8	0.09
Anus, anal canal & anorectum	1,694	1.9	0.05	1.9	0.05	1,845	1.7	0.04	1.7	0.04
Liver	14,693	16.0	0.14	15.7	0.14	4,700	4.3	0.06	4.1	0.06
Gallbladder	926	1.4	0.05	1.3	0.04	2,066	2.0	0.05	1.9	0.04
Pancreas	12,593	16.9	0.16	16.0	0.15	14,447	14.4	0.12	13.5	0.11
Respiratory System	70,577	95.5	0.39	91.0	0.36	52,541	51.4	0.23	49.0	0.22
Larynx	6,871	8.5	0.11	8.2	0.10	1,828	1.7	0.04	1.6	0.04
Lung & bronchus	62,847	85.9	0.37	81.7	0.34	50,098	49.1	0.22	46.8	0.21
Bones & Joints	831	0.9	0.03	0.8	0.03	761	0.7	0.03	0.7	0.03
Soft Tissue (incl. heart)	3,078	3.6	0.07	3.5	0.07	3,371	3.2	0.06	3.1	0.05
Skin (excl. basal & sq.)	1,655	2.1	0.06	2.0	0.05	1,862	1.8	0.04	1.8	0.04
Melanomas of the skin	828	1.1	0.04	1.1	0.04	981	1.0	0.03	0.9	0.03
Breast (excl. *in situ*)	1,387	1.9	0.05	1.8	0.05	130,554	123.0	0.35	119.9	0.34
Female Genital System	-	-	-	-	-	50,442	47.4	0.22	46.5	0.21
Cervix uteri	-	-	-	-	-	9,792	9.3	0.10	9.3	0.10
Uterus Corpus	-	-	-	-	-	25,323	23.5	0.15	23.2	0.15
Uterus, NOS[1]	-	-	-	-	-	1,802	1.7	0.04	1.6	0.04
Ovary	-	-	-	-	-	9,641	9.2	0.10	8.9	0.09
Male Genital System	152,688	186.3	0.51	182.0	0.49	-	-	-	-	-
Prostate	150,353	183.6	0.50	179.3	0.48	-	-	-	-	-
Testis	1,464	1.5	0.04	1.6	0.04	-	-	-	-	-
Penis	690	1.0	0.04	0.9	0.04	-	-	-	-	-
Urinary System	33,598	44.5	0.26	42.4	0.24	20,140	19.7	0.14	18.8	0.13
Bladder (incl. *in situ*)	13,090	19.5	0.18	18.1	0.17	6,440	6.6	0.08	6.1	0.08
Kidney & renal pelvis	19,982	24.2	0.18	23.6	0.17	13,230	12.6	0.11	12.2	0.11
Ureter	207	0.3	0.02	0.3	0.02	187	0.2	0.01	0.2	0.01
Eye & Orbit	250	0.3	0.02	0.3	0.02	252	0.2	0.01	0.2	0.01
Brain & Other Nervous System	4,269	4.8	0.08	4.7	0.08	3,769	3.5	0.06	3.4	0.06
Brain	3,928	4.4	0.08	4.3	0.07	3,359	3.1	0.05	3.1	0.05
Endocrine System	4,177	4.8	0.08	4.7	0.08	15,690	14.5	0.12	14.6	0.12
Thyroid gland	3,272	3.8	0.07	3.7	0.07	14,716	13.6	0.11	13.8	0.12
Lymphomas	16,841	20.0	0.16	19.5	0.16	15,024	14.3	0.12	14.0	0.12
Hodgkin Lymphoma	3,042	3.1	0.06	3.2	0.06	2,542	2.3	0.05	2.4	0.05
Non-Hodgkin Lymphoma	13,799	16.9	0.15	16.4	0.15	12,482	12.0	0.11	11.6	0.11
Multiple Myeloma	11,309	15.4	0.16	14.6	0.14	11,556	11.4	0.11	10.8	0.10
Leukemias	10,423	13.5	0.14	12.8	0.13	8,983	8.7	0.09	8.4	0.09
Acute lymphocytic	1,191	1.1	0.03	1.1	0.03	925	0.8	0.03	0.8	0.03
Chronic lymphocytic	3,176	4.4	0.08	4.2	0.08	2,327	2.3	0.05	2.2	0.05
Acute myeloid	3,142	4.1	0.08	3.9	0.07	3,189	3.1	0.06	3.0	0.05
Chronic myeloid	1,638	2.0	0.05	2.0	0.05	1,432	1.4	0.04	1.3	0.04
Other leukemias	1,276	1.7	0.05	1.6	0.05	1,110	1.1	0.03	1.0	0.03
Mesothelioma	564	0.9	0.04	0.8	0.04	232	0.2	0.02	0.2	0.01
Kaposi Sarcoma	1,497	1.6	0.04	1.7	0.04	165	0.2	0.01	0.2	0.01
Ill-defined & Unknown	13,216	18.8	0.18	17.5	0.16	14,427	14.5	0.12	13.5	0.11
Selected tumors (non-invasive)										
Brain & ONS[1] (*benign/borderline*)	6,525	8.1	0.11	7.8	0.10	15,224	14.6	0.12	14.1	0.12
Breast (*in situ*)	202	0.2	0.02	0.2	0.02	34,037	31.7	0.17	31.0	0.17

* Rates are per 100,000 population and are age-adjusted by five-year age groups to the 2000 U.S. standard population based on single years of age and to the 1996 Canadian population standard.

- Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for All Sites combined.

[1] Abbreviations: NOS - Not Otherwise Specified; ONS - Other Nervous System

Total Cancer Cases and Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates[1] for the Five Most Commonly Diagnosed Cancers
Selected Areas in the NAACCR United States Combined[2], 2010-2014, Black, Males

	All Sites			Prostate			Lung & Bronchus			Colon & Rectum			Kidney & Renal Pelvis			Liver & IBD[3]		
	Count	*Rate*	*S.E.*[3]	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
NAACCR U.S. Combined	**432,930**	**550.2**	**0.90**	**150,353**	**183.6**	**0.50**	**62,847**	**85.9**	**0.37**	**42,108**	**55.2**	**0.29**	**19,982**	**24.2**	**0.18**	**15,443**	**16.9**	**0.14**
Alabama	14,445	568.4	5.08	5,176	196.7	2.93	2,327	95.4	2.13	1,544	63.6	1.74	620	23.8	1.02	365	12.4	0.69
Alaska	219	465.4	39.21	88	164.7	21.37	27	78.5	17.89	21	43.3	11.58	8	12.4	5.02	9	16.0	6.24
Arizona	2,124	390.2	9.47	663	115.1	4.91	297	63.0	4.05	197	38.2	3.10	128	22.2	2.20	84	14.0	1.66
Arkansas	5,114	582.0	8.79	1,672	178.9	4.70	969	115.7	4.02	510	59.8	2.85	208	23.6	1.75	138	13.4	1.22
California	27,798	506.6	3.22	9,685	166.5	1.78	3,411	67.9	1.23	2,764	52.4	1.06	1,339	23.3	0.67	1,090	17.7	0.56
CA: Greater California	11,078	490.8	5.09	3,891	163.6	2.83	1,355	67.6	2.00	1,116	51.9	1.70	553	22.7	1.04	442	17.4	0.89
CA: Greater Bay Area	6,045	540.8	7.36	2,071	173.0	4.00	773	74.3	2.83	531	49.6	2.28	293	26.2	1.61	291	22.4	1.38
CA: Los Angeles	10,663	505.9	5.08	3,717	166.3	2.82	1,280	65.0	1.89	1,116	54.6	1.70	492	22.4	1.04	357	15.5	0.85
Colorado	2,021	455.0	11.32	696	151.8	6.44	223	58.6	4.32	201	47.4	3.75	105	22.2	2.33	100	18.4	2.00
Connecticut	3,779	524.1	9.16	1,351	179.0	5.18	465	71.7	3.57	347	50.3	2.92	172	20.3	1.63	111	13.9	1.40
Delaware	2,361	592.6	13.08	926	214.9	7.45	294	79.4	4.97	184	49.9	3.98	124	30.8	2.96	84	18.5	2.16
District of Columbia	4,621	601.2	9.12	1,461	186.0	5.02	685	89.4	3.52	454	61.4	2.97	198	25.5	1.87	255	29.6	1.91
Florida	29,403	479.6	2.96	10,680	169.7	1.73	3,787	65.5	1.13	2,940	49.5	0.97	1,098	17.0	0.54	838	12.0	0.44
Georgia	29,883	567.8	3.70	10,901	201.0	2.14	4,073	84.8	1.50	3,016	58.0	1.19	1,302	23.0	0.71	813	12.9	0.50
Hawaii	258	464.2	34.13	104	162.8	17.98	21	48.6	12.61	26	47.6	11.43	10	15.3	5.46	12	17.0	5.22
Idaho	70	356.0	52.18	26	118.8	26.76	6	44.1	23.49	-	-	-	-	-	-	8	50.6	24.40
Illinois	21,655	580.2	4.15	6,996	179.3	2.24	3,485	98.5	1.76	2,406	66.8	1.43	1,043	26.8	0.87	661	15.9	0.65
Indiana	5,702	518.4	7.35	1,616	140.2	3.71	982	96.8	3.30	577	53.6	2.39	322	26.6	1.57	223	16.9	1.18
Iowa	758	544.7	22.83	220	159.3	12.06	124	94.6	9.66	75	49.8	6.80	45	31.6	5.47	35	20.7	3.96
Kansas	1,901	574.0	14.29	630	186.4	8.01	317	100.0	6.09	190	61.6	4.86	84	23.6	2.75	65	16.5	2.17
Kentucky	4,072	605.5	10.31	1,158	165.2	5.24	733	118.4	4.75	415	62.8	3.36	236	31.6	2.19	142	17.8	1.61
Louisiana	18,001	625.5	5.01	5,907	199.7	2.78	2,978	109.5	2.16	1,914	66.7	1.64	798	27.5	1.03	676	19.7	0.80
Maine	114	370.7	38.86	36	121.0	22.09	8	31.9	12.31	7	17.8	6.96	-	-	-	7	25.3	11.19
Maryland	18,761	535.7	4.22	7,160	192.7	2.43	2,365	74.2	1.65	1,660	50.1	1.33	805	22.2	0.84	759	18.6	0.72
Massachusetts	4,798	493.8	7.69	1,853	183.3	4.54	526	62.0	2.91	408	43.2	2.32	192	19.0	1.47	204	18.8	1.42
Michigan	17,191	592.1	4.76	5,903	194.9	2.67	2,615	96.5	1.99	1,551	55.1	1.48	927	31.2	1.08	657	19.6	0.80
MI: Metropolitan Detroit	13,084	639.9	5.87	4,594	214.8	3.32	1,934	101.4	2.42	1,191	60.2	1.83	705	33.8	1.33	510	21.5	1.00
Mississippi	12,815	613.7	5.86	4,356	202.1	3.30	2,264	113.8	2.59	1,479	72.6	2.05	601	28.5	1.25	229	9.4	0.66
Missouri	7,608	568.9	6.96	2,237	158.5	3.55	1,342	105.4	3.06	748	60.0	2.34	362	25.8	1.44	369	23.6	1.29

Total Cancer Cases and Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates[1] for the Five Most Commonly Diagnosed Cancers
Selected Areas in the NAACCR United States Combined[2], 2010-2014, Black, Males

	All Sites			Prostate			Lung & Bronchus			Colon & Rectum			Kidney & Renal Pelvis			Liver & IBD[3]		
	Count	*Rate*	*S.E.*[3]	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
Montana	50	554.1	90.53	17	182.5	47.17	9	119.2	43.10	-	-	-	-	-	-	-	-	-
Nebraska	909	619.0	22.63	310	204.1	12.62	125	96.3	9.36	107	78.6	8.45	54	35.5	5.34	36	21.6	3.87
New Hampshire	138	407.9	38.82	52	151.0	23.13	12	35.8	11.09	15	43.4	12.22	6	17.1	7.78	7	23.2	9.74
New Jersey	13,468	533.6	4.88	5,172	199.5	2.92	1,622	70.5	1.86	1,253	50.7	1.52	610	22.7	0.97	409	14.1	0.74
New York	38,712	560.3	3.00	15,490	216.8	1.82	4,217	65.2	1.06	3,371	51.3	0.93	1,574	21.8	0.57	1,419	18.6	0.52
North Carolina	24,679	570.3	3.91	8,576	189.5	2.19	4,030	99.4	1.69	2,232	52.8	1.21	1,120	25.1	0.80	737	14.5	0.57
North Dakota	48	432.0	87.51	14	116.8	41.09	7	98.0	45.95	6	58.4	32.24	-	-	-	-	-	-
Ohio	15,808	533.4	4.45	5,272	170.5	2.46	2,660	95.3	1.93	1,459	50.6	1.39	770	24.9	0.93	607	17.6	0.75
Oklahoma	3,182	591.4	11.47	1,082	199.1	6.59	527	102.5	4.85	298	54.1	3.43	144	26.7	2.41	99	15.5	1.70
Oregon	819	500.7	19.85	263	156.4	10.87	119	81.9	8.55	75	50.4	6.51	44	23.2	3.93	50	29.2	4.64
Pennsylvania	16,633	588.9	4.82	5,209	176.7	2.58	2,506	97.0	2.04	1,475	53.9	1.48	905	30.0	1.04	915	28.2	0.98
Rhode Island	578	437.0	19.93	181	124.5	10.02	75	65.2	8.22	43	37.0	6.12	24	17.8	4.01	22	11.9	2.68
South Carolina	15,464	563.2	4.87	5,431	188.1	2.73	2,373	90.4	2.00	1,464	54.8	1.54	641	22.7	0.95	388	12.3	0.67
South Dakota	65	261.1	36.95	10	32.6	10.55	8	39.6	17.91	7	28.5	11.20	6	21.3	8.93	-	-	-
Tennessee	11,230	564.8	5.87	3,743	182.1	3.26	1,900	103.3	2.61	1,094	57.8	1.92	504	24.5	1.19	384	16.0	0.88
Texas	29,896	537.3	3.45	9,029	157.5	1.83	4,573	91.3	1.49	3,236	58.4	1.14	1,514	25.5	0.72	1,516	22.4	0.62
Utah	245	505.4	39.92	89	194.2	24.50	24	55.6	13.51	29	57.5	13.48	14	28.4	9.25	16	30.1	8.68
Vermont	35	270.5	52.15	14	108.5	33.13	-	-	-	-	-	-	-	-	-	-	-	-
Virginia	17,922	538.0	4.30	6,428	180.7	2.39	2,676	88.0	1.82	1,601	49.6	1.33	902	25.7	0.90	545	13.8	0.62
Washington	2,772	522.2	11.39	936	165.7	6.11	315	72.7	4.62	225	43.0	3.27	165	27.2	2.37	140	23.7	2.26
WA: Seattle/Puget Sound	2,520	539.5	12.33	865	171.9	6.57	286	76.6	5.08	199	43.9	3.56	145	27.2	2.53	134	26.3	2.57
West Virginia	859	522.5	19.47	282	159.1	10.23	134	92.4	8.71	92	55.8	6.37	46	25.3	3.96	38	18.7	3.23
Wisconsin	3,914	670.9	12.02	1,243	207.0	6.51	606	116.7	5.28	375	66.2	3.85	193	29.4	2.35	172	24.5	1.98
Wyoming	32	179.2	32.84	9	47.9	16.53	-	-	-	6	40.8	16.67	-	-	-	-	-	-

1. Rates are per 100,000 population and are age-adjusted by five-year age groups to the 2000 U.S. standard population based on single years of age.
2. See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.
3. Abbreviations: IBD - Intrahepatic Bile Duct; S.E. - Standard Error

\- Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for the U.S. combined.

Total Cancer Cases and Average Annual Age-adjusted (1996 Canadian Standard) Cancer Incidence Rates[1] for the Five Most Commonly Diagnosed Cancers
Selected Areas in the NAACCR United States Combined[2], 2010-2014, Black, Males

	All Sites			Prostate			Lung & Bronchus			Colon & Rectum			Kidney & Renal Pelvis			Liver & IBD[3]		
	Count	*Rate*	*S.E.*[3]	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
NAACCR U.S. Combined	**432,930**	**529.9**	**0.85**	**150,353**	**179.3**	**0.48**	**62,847**	**81.7**	**0.34**	**42,108**	**52.5**	**0.27**	**19,982**	**23.6**	**0.17**	**15,443**	**16.6**	**0.14**
Alabama	14,445	546.4	4.78	5,176	190.5	2.79	2,327	91.1	1.99	1,544	60.4	1.61	620	23.0	0.97	365	12.1	0.67
Alaska	219	436.5	35.24	88	157.5	19.49	27	72.2	15.97	21	41.5	10.76	8	12.3	5.05	9	15.6	6.27
Arizona	2,124	374.7	8.80	663	112.1	4.66	297	59.6	3.71	197	35.8	2.79	128	21.7	2.07	84	13.8	1.60
Arkansas	5,114	562.5	8.34	1,672	175.9	4.56	969	110.9	3.78	510	57.5	2.69	208	22.8	1.67	138	13.2	1.19
California	27,798	487.6	3.05	9,685	163.0	1.72	3,411	64.4	1.14	2,764	49.7	0.99	1,339	22.6	0.64	1,090	17.4	0.55
CA: Greater California	11,078	471.2	4.77	3,891	159.4	2.71	1,355	64.0	1.84	1,116	49.1	1.57	553	22.1	1.00	442	17.0	0.86
CA: Greater Bay Area	6,045	521.1	6.98	2,071	169.8	3.89	773	70.4	2.64	531	46.9	2.13	293	25.2	1.53	291	22.3	1.36
CA: Los Angeles	10,663	488.0	4.85	3,717	163.3	2.75	1,280	61.6	1.76	1,116	52.1	1.60	492	21.9	1.01	357	15.3	0.83
Colorado	2,021	437.9	10.57	696	146.7	6.04	223	55.6	3.98	201	45.1	3.45	105	21.9	2.28	100	18.2	1.95
Connecticut	3,779	506.1	8.66	1,351	175.8	5.01	465	68.0	3.30	347	47.6	2.70	172	20.2	1.60	111	13.7	1.36
Delaware	2,361	568.4	12.24	926	210.6	7.19	294	75.1	4.58	184	46.7	3.61	124	29.6	2.78	84	17.9	2.04
District of Columbia	4,621	583.5	8.82	1,461	181.7	4.88	685	86.3	3.38	454	58.8	2.83	198	24.7	1.80	255	29.3	1.89
Florida	29,403	462.7	2.80	10,680	166.4	1.67	3,787	62.3	1.05	2,940	47.1	0.90	1,098	16.6	0.52	838	11.8	0.42
Georgia	29,883	546.6	3.44	10,901	195.8	2.03	4,073	80.7	1.37	3,016	55.2	1.10	1,302	22.3	0.67	813	12.7	0.48
Hawaii	258	444.3	31.58	104	160.5	17.53	21	43.5	10.88	26	45.1	10.36	10	15.4	5.56	12	16.9	5.27
Idaho	70	350.1	50.00	26	116.9	26.13	6	39.9	20.52	-	-	-	-	-	-	8	47.2	21.88
Illinois	21,655	557.6	3.92	6,996	174.9	2.16	3,485	93.7	1.64	2,406	63.4	1.34	1,043	26.0	0.83	661	15.6	0.63
Indiana	5,702	498.0	6.92	1,616	136.4	3.56	982	92.0	3.07	577	51.2	2.23	322	26.0	1.51	223	16.8	1.17
Iowa	758	519.5	21.09	220	154.6	11.43	124	90.2	8.95	75	46.2	6.07	45	29.9	5.02	35	19.9	3.76
Kansas	1,901	553.0	13.51	630	181.0	7.66	317	95.4	5.71	190	58.3	4.50	84	23.2	2.67	65	16.4	2.14
Kentucky	4,072	582.4	9.72	1,158	161.6	5.05	733	112.4	4.42	415	59.5	3.11	236	31.4	2.16	142	17.5	1.57
Louisiana	18,001	604.0	4.74	5,907	195.0	2.67	2,978	104.4	2.02	1,914	64.0	1.54	798	27.0	1.00	676	19.3	0.78
Maine	114	368.6	37.85	36	121.4	21.99	8	31.9	12.03	7	18.8	7.38	-	-	-	7	24.5	10.45
Maryland	18,761	515.6	3.96	7,160	188.7	2.34	2,365	70.1	1.52	1,660	47.4	1.23	805	21.5	0.79	759	18.0	0.69
Massachusetts	4,798	477.1	7.29	1,853	180.3	4.42	526	58.2	2.68	408	40.8	2.14	192	18.5	1.41	204	18.3	1.36
Michigan	17,191	571.9	4.54	5,903	191.3	2.59	2,615	91.9	1.87	1,551	52.3	1.38	927	30.4	1.04	657	19.4	0.79
MI: Metropolitan Detroit	13,084	618.6	5.61	4,594	210.7	3.22	1,934	96.4	2.27	1,191	57.3	1.72	705	32.9	1.28	510	21.3	0.98
Mississippi	12,815	591.9	5.53	4,356	197.3	3.15	2,264	108.8	2.42	1,479	69.3	1.91	601	27.7	1.19	229	9.1	0.64
Missouri	7,608	547.3	6.58	2,237	155.0	3.43	1,342	100.9	2.88	748	56.5	2.16	362	25.1	1.38	369	23.1	1.26

Total Cancer Cases and Average Annual Age-adjusted (1996 Canadian Standard) Cancer Incidence Rates[1] for the Five Most Commonly Diagnosed Cancers
Selected Areas in the NAACCR United States Combined[2], 2010-2014, Black, Males

	All Sites			Prostate			Lung & Bronchus			Colon & Rectum			Kidney & Renal Pelvis			Liver & IBD[3]		
	Count	*Rate*	*S.E.*[3]	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
Montana	50	540.0	85.66	17	186.8	48.62	9	115.5	41.26	-	-	-	-	-	-	-	-	-
Nebraska	909	593.1	21.18	310	199.2	12.08	125	90.7	8.65	107	74.6	7.77	54	34.0	4.99	36	21.1	3.75
New Hampshire	138	396.7	37.04	52	147.9	22.33	12	35.9	11.06	15	43.0	12.10	6	16.8	7.45	7	21.9	8.98
New Jersey	13,468	513.9	4.61	5,172	194.6	2.81	1,622	66.8	1.72	1,253	48.5	1.43	610	22.2	0.93	409	13.8	0.71
New York	38,712	541.0	2.85	15,490	212.1	1.76	4,217	62.0	0.99	3,371	48.5	0.87	1,574	21.3	0.55	1,419	18.3	0.50
North Carolina	24,679	549.8	3.68	8,576	184.7	2.09	4,030	95.0	1.57	2,232	50.4	1.12	1,120	24.5	0.76	737	14.1	0.54
North Dakota	48	394.5	75.94	14	108.9	36.68	7	91.0	41.42	6	55.6	29.05	-	-	-	-	-	-
Ohio	15,808	513.7	4.23	5,272	166.7	2.38	2,660	91.1	1.82	1,459	47.9	1.30	770	24.3	0.91	607	17.3	0.73
Oklahoma	3,182	566.0	10.68	1,082	191.8	6.20	527	98.1	4.53	298	51.4	3.18	144	25.6	2.27	99	15.0	1.60
Oregon	819	483.0	18.60	263	151.7	10.28	119	77.5	7.86	75	48.6	6.09	44	22.3	3.66	50	28.7	4.44
Pennsylvania	16,633	567.8	4.59	5,209	172.5	2.49	2,506	92.3	1.91	1,475	50.8	1.38	905	29.4	1.02	915	27.8	0.96
Rhode Island	578	418.9	18.74	181	122.1	9.73	75	60.9	7.50	43	34.9	5.63	24	16.8	3.72	22	11.5	2.55
South Carolina	15,464	542.2	4.57	5,431	183.9	2.61	2,373	86.1	1.86	1,464	52.2	1.43	641	22.1	0.91	388	11.9	0.63
South Dakota	65	260.1	36.88	10	31.3	10.39	8	37.2	16.32	7	29.6	11.90	6	21.7	9.28	-	-	-
Tennessee	11,230	542.5	5.50	3,743	177.3	3.10	1,900	98.1	2.42	1,094	55.0	1.78	504	23.6	1.12	384	15.5	0.85
Texas	29,896	514.2	3.21	9,029	152.2	1.72	4,573	86.5	1.37	3,236	55.5	1.05	1,514	24.8	0.68	1,516	22.0	0.60
Utah	245	492.5	37.99	89	191.8	23.80	24	54.9	12.94	29	55.1	12.54	14	28.5	9.04	16	30.9	9.01
Vermont	35	270.4	51.60	14	105.8	32.19	-	-	-	-	-	-	-	-	-	-	-	-
Virginia	17,922	517.5	4.05	6,428	177.0	2.31	2,676	83.2	1.68	1,601	47.0	1.23	902	25.3	0.88	545	13.5	0.61
Washington	2,772	501.2	10.61	936	161.9	5.85	315	68.3	4.21	225	41.3	3.06	165	26.7	2.29	140	23.2	2.17
WA: Seattle/Puget Sound	2,520	518.3	11.50	865	168.6	6.32	286	72.0	4.64	199	42.1	3.32	145	26.8	2.44	134	25.8	2.46
West Virginia	859	507.5	18.53	282	158.3	10.09	134	88.3	8.12	92	53.9	6.00	46	24.8	3.83	38	18.6	3.19
Wisconsin	3,914	647.4	11.29	1,243	201.5	6.20	606	111.7	4.91	375	63.1	3.56	193	28.7	2.24	172	24.4	1.98
Wyoming	32	170.2	31.23	9	47.8	16.65	-	-	-	6	37.7	15.51	-	-	-	-	-	-

1. Rates are per 100,000 and are age-adjusted to the 1996 Canadian population standard.
2. See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.
3. Abbreviations: IBD - Intrahepatic Bile Duct; S.E. - Standard Error

\- Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for the U.S. combined.

Total Cancer Cases and Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates[1] for the Five Most Commonly Diagnosed Cancers
Selected Areas in the NAACCR United States Combined[2], 2010-2014, Black, Females

	All Sites			Breast			Lung & Bronchus			Colon & Rectum			Uterus Corpus & Uterus, NOS[3]			Thyroid		
	Count	*Rate*	*S.E.*[3]	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
NAACCR U.S. Combined	**418,018**	**400.2**	**0.63**	**130,554**	**123.0**	**0.35**	**50,098**	**49.1**	**0.22**	**42,301**	**41.0**	**0.20**	**27,125**	**25.2**	**0.16**	**14,716**	**13.6**	**0.11**
Alabama	12,936	377.6	3.39	4,314	124.4	1.93	1,313	38.5	1.09	1,508	44.4	1.17	750	21.6	0.81	339	9.9	0.55
Alaska	157	324.8	27.47	61	126.1	16.94	11	27.1	8.36	12	22.2	6.77	-	-	-	9	15.5	5.53
Arizona	1,989	335.5	7.84	625	101.3	4.19	235	42.5	2.88	193	33.9	2.55	134	21.9	1.96	116	17.1	1.63
Arkansas	4,361	385.7	5.99	1,332	116.3	3.27	543	48.9	2.16	511	45.6	2.07	252	21.7	1.41	114	9.7	0.92
California	25,892	388.3	2.46	8,312	123.0	1.37	3,113	47.8	0.87	2,680	40.9	0.80	1,715	24.7	0.61	797	11.8	0.43
CA: Greater California	9,490	372.0	3.95	3,092	117.0	2.17	1,119	46.8	1.45	959	39.5	1.32	589	22.0	0.93	337	11.8	0.66
CA: Greater Bay Area	5,409	406.8	5.65	1,691	126.2	3.13	702	53.4	2.06	563	42.3	1.83	362	26.3	1.41	107	8.2	0.81
CA: Los Angeles	10,989	395.5	3.82	3,529	126.8	2.17	1,291	46.1	1.30	1,158	41.5	1.24	764	26.4	0.97	353	13.3	0.72
Colorado	1,647	345.9	8.79	555	113.5	4.95	185	42.5	3.22	149	33.3	2.81	88	18.5	2.03	81	14.8	1.68
Connecticut	3,847	386.7	6.37	1,191	116.0	3.43	408	42.9	2.18	360	37.1	2.00	274	26.5	1.63	182	17.2	1.29
Delaware	2,212	422.9	9.24	696	129.4	5.03	278	56.7	3.50	184	35.7	2.71	158	29.5	2.41	95	17.2	1.79
District of Columbia	4,922	463.4	6.80	1,451	140.0	3.79	626	56.9	2.32	508	47.0	2.14	387	34.7	1.81	174	17.9	1.42
Florida	27,785	352.9	2.16	8,474	105.8	1.17	2,677	35.0	0.69	2,776	36.0	0.70	2,052	25.3	0.57	973	11.9	0.39
Georgia	29,098	392.5	2.39	9,650	125.4	1.32	2,877	41.0	0.80	3,061	42.4	0.80	1,863	24.5	0.59	932	11.4	0.38
Hawaii	131	324.2	31.44	54	118.4	17.38	11	41.9	13.13	6	19.1	8.88	6	15.4	6.85	-	-	-
Idaho	42	324.2	55.65	12	96.0	31.41	-	-	-	-	-	-	-	-	-	-	-	-
Illinois	22,521	435.6	2.95	6,781	130.4	1.61	3,228	62.8	1.12	2,489	48.5	0.99	1,469	27.3	0.72	655	12.6	0.50
Indiana	6,152	421.0	5.49	1,928	129.5	3.01	875	61.6	2.13	625	43.7	1.79	342	22.9	1.27	154	9.9	0.81
Iowa	668	440.7	17.92	163	103.1	8.43	102	73.0	7.54	65	46.4	5.99	45	29.5	4.57	17	9.2	2.31
Kansas	1,659	426.2	10.67	488	124.2	5.72	238	63.4	4.18	153	39.8	3.27	96	24.8	2.59	77	18.8	2.19
Kentucky	3,820	447.3	7.42	1,096	126.2	3.90	643	77.9	3.16	423	49.5	2.46	200	23.2	1.68	133	15.1	1.34
Louisiana	15,759	422.1	3.44	4,975	131.6	1.91	1,858	50.3	1.20	1,863	50.4	1.20	841	21.8	0.77	531	14.0	0.62
Maine	77	308.5	37.58	18	69.0	16.81	10	46.3	15.72	-	-	-	7	29.6	11.85	-	-	-
Maryland	19,548	401.5	2.95	6,575	131.3	1.66	2,235	47.3	1.03	1,779	37.2	0.91	1,321	26.2	0.74	775	15.2	0.55
Massachusetts	4,624	363.0	5.45	1,387	106.7	2.92	504	42.2	1.92	425	33.6	1.67	296	22.7	1.35	302	21.8	1.27
Michigan	16,371	428.3	3.42	4,855	126.4	1.85	2,403	63.1	1.32	1,617	42.3	1.07	1,059	26.6	0.84	530	14.0	0.62
MI: Metropolitan Detroit	12,606	447.0	4.07	3,739	132.1	2.21	1,831	64.6	1.55	1,281	45.5	1.30	798	27.1	0.98	433	15.6	0.76
Mississippi	11,187	404.8	3.92	3,399	120.5	2.11	1,271	47.5	1.37	1,436	52.4	1.42	640	22.7	0.92	350	12.1	0.66
Missouri	8,076	444.6	5.04	2,439	132.7	2.74	1,215	67.8	1.99	788	43.8	1.59	499	26.8	1.23	228	12.2	0.82

Total Cancer Cases and Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates[1] for the Five Most Commonly Diagnosed Cancers
Selected Areas in the NAACCR United States Combined[2], 2010-2014, Black, Females

	All Sites			Breast			Lung & Bronchus			Colon & Rectum			Uterus Corpus & Uterus, NOS[3]			Thyroid		
	Count	*Rate*	*S.E.*[3]	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
Montana	19	286.3	69.79	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Nebraska	739	427.6	16.25	216	125.7	8.79	108	64.3	6.36	71	42.7	5.22	35	19.0	3.32	31	15.3	2.82
New Hampshire	68	209.2	26.54	24	75.9	16.05	8	26.1	9.34	6	19.7	8.52	6	18.6	8.05	-	-	-
New Jersey	14,212	401.2	3.41	4,351	120.3	1.85	1,642	47.4	1.19	1,451	41.8	1.11	987	27.2	0.88	490	13.5	0.62
New York	38,608	389.0	2.01	11,487	115.3	1.09	4,031	40.6	0.65	3,680	37.3	0.62	3,148	30.7	0.55	1,602	16.0	0.40
North Carolina	23,841	407.2	2.69	7,874	132.8	1.53	2,732	47.2	0.92	2,224	38.2	0.83	1,525	25.3	0.66	826	14.0	0.49
North Dakota	27	331.4	82.33	6	53.3	21.94	-	-	-	-	-	-	-	-	-	-	-	-
Ohio	15,521	402.0	3.28	4,751	122.5	1.81	2,429	63.3	1.30	1,454	37.8	1.00	920	23.1	0.77	603	16.1	0.66
Oklahoma	2,739	402.0	7.90	845	121.3	4.28	373	57.2	3.05	272	39.8	2.48	165	23.4	1.87	85	11.6	1.29
Oregon	619	389.1	16.20	196	120.2	8.84	78	53.5	6.27	54	36.7	5.18	38	22.9	3.82	30	16.6	3.10
Pennsylvania	17,624	459.7	3.51	4,875	127.2	1.85	2,603	68.7	1.37	1,600	41.6	1.05	1,141	28.9	0.87	841	22.0	0.77
Rhode Island	588	346.0	14.68	179	102.4	7.86	77	47.7	5.58	51	30.9	4.46	33	20.1	3.60	39	20.7	3.38
South Carolina	14,158	389.0	3.35	4,583	124.9	1.89	1,540	42.1	1.10	1,370	37.1	1.03	993	26.4	0.86	461	12.8	0.61
South Dakota	39	203.8	33.61	16	88.0	22.25	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee	10,986	403.8	3.96	3,518	125.7	2.18	1,359	52.1	1.46	1,127	42.6	1.31	620	22.0	0.91	297	10.5	0.62
Texas	29,307	397.5	2.40	9,030	118.1	1.28	3,378	48.9	0.87	3,062	41.8	0.78	1,696	22.4	0.56	1,019	12.6	0.40
Utah	136	310.4	29.22	35	76.5	13.77	13	34.6	10.26	16	43.2	11.46	6	18.8	8.15	8	13.2	5.18
Vermont	23	237.9	55.07	8	70.6	26.78	-	-	-	-	-	-	-	-	-	-	-	-
Virginia	17,037	393.8	3.08	5,844	132.7	1.77	2,015	47.6	1.08	1,710	39.9	0.98	953	21.6	0.71	541	12.3	0.54
Washington	2,034	384.1	9.05	670	122.0	4.97	222	47.7	3.37	156	29.3	2.49	103	19.5	2.03	100	15.5	1.61
WA: Seattle/Puget Sound	1,898	404.4	9.87	632	129.1	5.42	209	51.5	3.75	145	31.1	2.73	96	20.7	2.24	96	16.8	1.79
West Virginia	657	381.4	15.25	201	119.3	8.65	91	51.4	5.49	64	37.1	4.76	36	19.1	3.29	25	15.4	3.16
Wisconsin	3,525	477.4	8.41	1,003	130.7	4.28	522	76.4	3.49	308	42.5	2.53	208	27.8	2.01	130	15.3	1.40
Wyoming	30	171.5	32.17	7	43.8	16.99	-	-	-	-	-	-	-	-	-	-	-	-

1. Rates are per 100,000 population and are age-adjusted by five-year age groups to the 2000 U.S. standard population based on single years of age.
2. See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.
3. Abbreviations: NOS - Not Otherwise Specified; S.E. - Standard Error

\- Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for the U.S. combined.

Total Cancer Cases and Average Annual Age-adjusted (1996 Canadian Standard) Cancer Incidence Rates[1] for the Five Most Commonly Diagnosed Cancers Selected Areas in the NAACCR United States Combined[2], 2010-2014, Black, Females

	All Sites			Breast			Lung & Bronchus			Colon & Rectum			Uterus Corpus & Uterus, NOS[3]			Thyroid		
	Count	*Rate*	*S.E.*[3]	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
NAACCR U.S. Combined	**418,018**	**386.4**	**0.61**	**130,554**	**119.9**	**0.34**	**50,098**	**46.8**	**0.21**	**42,301**	**38.9**	**0.19**	**27,125**	**24.8**	**0.15**	**14,716**	**13.8**	**0.12**
Alabama	12,936	365.3	3.29	4,314	121.8	1.90	1,313	36.8	1.04	1,508	42.0	1.11	750	21.1	0.79	339	10.0	0.56
Alaska	157	318.6	26.73	61	125.9	16.78	11	28.0	8.67	12	21.4	6.49	-	-	-	9	15.4	5.37
Arizona	1,989	323.8	7.47	625	99.2	4.07	235	40.8	2.74	193	32.0	2.37	134	21.5	1.91	116	17.3	1.64
Arkansas	4,361	373.7	5.82	1,332	113.1	3.19	543	46.9	2.07	511	43.7	1.99	252	21.4	1.39	114	9.9	0.95
California	25,892	374.5	2.37	8,312	119.5	1.34	3,113	45.5	0.83	2,680	38.7	0.76	1,715	24.4	0.60	797	11.9	0.43
CA: Greater California	9,490	358.4	3.78	3,092	113.9	2.10	1,119	44.5	1.36	959	37.3	1.24	589	21.8	0.92	337	12.0	0.67
CA: Greater Bay Area	5,409	392.5	5.46	1,691	122.5	3.05	702	50.7	1.96	563	40.2	1.74	362	25.9	1.39	107	8.3	0.82
CA: Los Angeles	10,989	381.6	3.70	3,529	123.2	2.11	1,291	44.1	1.24	1,158	39.4	1.18	764	26.1	0.96	353	13.3	0.73
Colorado	1,647	334.4	8.45	555	110.9	4.82	185	40.8	3.07	149	31.6	2.64	88	17.8	1.94	81	15.2	1.72
Connecticut	3,847	372.9	6.11	1,191	112.9	3.32	408	40.9	2.06	360	35.2	1.89	274	26.2	1.61	182	17.4	1.31
Delaware	2,212	408.0	8.87	696	126.0	4.89	278	53.5	3.28	184	34.0	2.56	158	29.0	2.36	95	17.2	1.81
District of Columbia	4,922	447.9	6.65	1,451	135.7	3.71	626	54.2	2.23	508	44.9	2.08	387	33.9	1.78	174	18.1	1.45
Florida	27,785	341.6	2.08	8,474	103.3	1.14	2,677	33.4	0.66	2,776	34.2	0.66	2,052	25.0	0.56	973	12.0	0.39
Georgia	29,098	379.3	2.29	9,650	122.4	1.28	2,877	39.2	0.76	3,061	40.4	0.75	1,863	24.2	0.58	932	11.5	0.38
Hawaii	131	316.5	30.21	54	119.1	17.35	11	41.5	12.91	6	17.7	8.12	6	15.3	6.59	-	-	-
Idaho	42	314.0	53.07	12	87.1	27.78	-	-	-	-	-	-	-	-	-	-	-	-
Illinois	22,521	420.1	2.85	6,781	127.1	1.57	3,228	59.9	1.07	2,489	45.8	0.93	1,469	26.8	0.71	655	12.8	0.51
Indiana	6,152	404.9	5.27	1,928	126.0	2.93	875	58.8	2.03	625	41.4	1.69	342	22.5	1.24	154	10.2	0.83
Iowa	668	422.1	17.09	163	99.7	8.10	102	68.6	7.07	65	43.9	5.66	45	29.4	4.59	17	9.3	2.33
Kansas	1,659	410.7	10.30	488	121.2	5.59	238	60.2	3.98	153	37.7	3.12	96	24.2	2.52	77	18.8	2.20
Kentucky	3,820	429.7	7.14	1,096	122.6	3.80	643	73.8	2.99	423	47.0	2.35	200	22.9	1.66	133	15.2	1.36
Louisiana	15,759	408.5	3.33	4,975	128.5	1.87	1,858	48.3	1.14	1,863	48.2	1.14	841	21.4	0.76	531	14.2	0.63
Maine	77	306.6	37.31	18	67.4	16.43	10	45.5	15.37	-	-	-	7	30.6	12.29	-	-	-
Maryland	19,548	387.4	2.84	6,575	128.1	1.62	2,235	45.1	0.98	1,779	35.1	0.85	1,321	25.7	0.72	775	15.4	0.56
Massachusetts	4,624	351.4	5.27	1,387	104.2	2.85	504	40.2	1.83	425	32.0	1.58	296	22.4	1.33	302	22.0	1.29
Michigan	16,371	413.2	3.31	4,855	123.3	1.81	2,403	60.0	1.25	1,617	40.1	1.02	1,059	26.1	0.82	530	14.1	0.63
MI: Metropolitan Detroit	12,606	430.9	3.94	3,739	128.9	2.16	1,831	61.4	1.47	1,281	43.1	1.23	798	26.6	0.96	433	15.7	0.77
Mississippi	11,187	391.4	3.79	3,399	117.8	2.07	1,271	45.3	1.30	1,436	50.0	1.35	640	22.4	0.91	350	12.3	0.67
Missouri	8,076	429.0	4.87	2,439	129.0	2.67	1,215	64.6	1.89	788	41.6	1.51	499	26.5	1.21	228	12.3	0.83

Total Cancer Cases and Average Annual Age-adjusted (1996 Canadian Standard) Cancer Incidence Rates[1] for the Five Most Commonly Diagnosed Cancers
Selected Areas in the NAACCR United States Combined[2], 2010-2014, Black, Females

	All Sites			Breast			Lung & Bronchus			Colon & Rectum			Uterus Corpus & Uterus, NOS[3]			Thyroid		
	Count	*Rate*	*S.E.*[3]	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
Montana	19	284.4	68.96	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Nebraska	739	409.4	15.46	216	119.9	8.34	108	62.2	6.13	71	40.7	4.95	35	18.7	3.24	31	15.2	2.79
New Hampshire	68	200.0	25.27	24	73.5	15.61	8	25.4	9.11	6	17.5	7.30	6	17.9	7.61	-	-	-
New Jersey	14,212	387.0	3.29	4,351	117.4	1.80	1,642	45.0	1.12	1,451	39.6	1.05	987	26.7	0.86	490	13.6	0.63
New York	38,608	375.8	1.94	11,487	112.5	1.07	4,031	38.6	0.62	3,680	35.3	0.59	3,148	30.2	0.55	1,602	16.1	0.41
North Carolina	23,841	393.2	2.60	7,874	129.4	1.49	2,732	45.1	0.88	2,224	36.1	0.78	1,525	25.0	0.65	826	14.0	0.50
North Dakota	27	305.9	73.43	6	49.7	20.43	-	-	-	-	-	-	-	-	-	-	-	-
Ohio	15,521	387.8	3.18	4,751	119.1	1.77	2,429	60.4	1.25	1,454	35.8	0.96	920	22.6	0.76	603	16.2	0.68
Oklahoma	2,739	388.6	7.61	845	118.8	4.18	373	54.4	2.89	272	37.6	2.33	165	22.7	1.81	85	11.9	1.32
Oregon	619	373.3	15.45	196	117.8	8.64	78	50.2	5.84	54	33.8	4.74	38	22.6	3.75	30	16.5	3.06
Pennsylvania	17,624	443.9	3.41	4,875	123.8	1.81	2,603	65.7	1.31	1,600	39.5	1.01	1,141	28.4	0.86	841	22.4	0.79
Rhode Island	588	335.6	14.27	179	99.4	7.66	77	46.3	5.44	51	29.3	4.22	33	19.9	3.55	39	20.9	3.43
South Carolina	14,158	375.9	3.24	4,583	121.8	1.85	1,540	40.3	1.05	1,370	35.4	0.98	993	25.8	0.84	461	12.9	0.62
South Dakota	39	201.4	33.19	16	89.7	22.64	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee	10,986	389.8	3.82	3,518	122.6	2.12	1,359	49.6	1.38	1,127	40.4	1.24	620	21.7	0.90	297	10.6	0.63
Texas	29,307	383.6	2.30	9,030	115.2	1.24	3,378	46.7	0.83	3,062	39.8	0.74	1,696	22.0	0.55	1,019	12.7	0.41
Utah	136	296.9	27.64	35	75.7	13.57	13	32.2	9.48	16	40.0	10.59	6	17.1	7.33	8	13.2	5.19
Vermont	23	228.9	51.87	8	70.6	26.13	-	-	-	-	-	-	-	-	-	-	-	-
Virginia	17,037	379.9	2.97	5,844	129.3	1.73	2,015	45.1	1.03	1,710	37.8	0.93	953	21.1	0.70	541	12.5	0.55
Washington	2,034	368.0	8.58	670	118.4	4.79	222	45.5	3.20	156	27.8	2.34	103	18.6	1.93	100	15.6	1.61
WA: Seattle/Puget Sound	1,898	387.5	9.36	632	125.5	5.23	209	48.9	3.54	145	29.5	2.58	96	19.8	2.12	96	17.0	1.79
West Virginia	657	366.4	14.80	201	115.2	8.43	91	49.3	5.30	64	34.8	4.52	36	18.6	3.22	25	15.3	3.19
Wisconsin	3,525	461.0	8.05	1,003	127.3	4.14	522	73.4	3.33	308	40.3	2.38	208	27.5	1.97	130	15.3	1.38
Wyoming	30	170.4	32.20	7	44.2	17.16	-	-	-	-	-	-	-	-	-	-	-	-

1. Rates are per 100,000 and are age-adjusted to the 1996 Canadian population standard.
2. See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.
3. Abbreviations: NOS - Not Otherwise Specified; S.E. - Standard Error

\- Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for the U.S. combined.

NAACCR U.S. Combined Incidence 2010-2014,
The Asian/Pacific Islander Population

Total Cancer Cases and Average Annual Age-adjusted Cancer Incidence Rates*
NAACCR United States Combined, 2010-2014, Asian/Pacific Islander (API)

	MALES (API)					FEMALES (API)				
SITES	*Count*	*Adj. Rate 2000 U.S.*	*S.E.*	*Adj. Rate 1996 Cdn*	*S.E.*	*Count*	*Adj. Rate 2000 U.S.*	*S.E.*	*Adj. Rate 1996 Cdn*	*S.E.*
All Sites (invasive & in situ bladder)	102,876	302.8	0.99	287.4	0.92	128,044	287.6	0.82	277.1	0.78
Oral Cavity & Pharynx	4,207	11.2	0.18	10.9	0.17	2,232	5.0	0.11	4.9	0.10
Lip	39	0.1	0.02	0.1	0.02	30	0.1	0.01	0.1	0.01
Tongue	900	2.5	0.09	2.4	0.08	624	1.4	0.06	1.4	0.06
Major salivary gland	393	1.1	0.06	1.1	0.05	387	0.8	0.04	0.8	0.04
Floor of mouth	89	0.3	0.03	0.2	0.03	57	0.1	0.02	0.1	0.02
Gum & other mouth	652	1.8	0.07	1.7	0.07	376	0.9	0.05	0.8	0.04
Nasopharynx	1,413	3.5	0.10	3.4	0.09	547	1.1	0.05	1.1	0.05
Oropharynx	89	0.3	0.03	0.2	0.03	34	0.1	0.01	0.1	0.01
Hypopharynx	217	0.7	0.05	0.6	0.04	40	0.1	0.02	0.1	0.01
Digestive System	31,206	91.6	0.54	86.5	0.50	24,664	58.5	0.38	54.9	0.35
Esophagus	1,248	3.7	0.11	3.5	0.10	398	1.0	0.05	0.9	0.05
Stomach	4,553	14.1	0.22	13.1	0.20	3,376	8.1	0.14	7.5	0.13
Small intestine	472	1.3	0.06	1.3	0.06	404	0.9	0.05	0.9	0.04
Colon excl. rectum	7,744	23.1	0.28	21.7	0.25	7,904	18.7	0.21	17.5	0.20
Rectum & rectosigmoid	5,059	13.9	0.20	13.3	0.19	3,691	8.3	0.14	7.9	0.13
Anus, anal canal & anorectum	167	0.5	0.04	0.4	0.03	220	0.5	0.03	0.5	0.03
Liver	6,643	18.5	0.24	17.7	0.22	2,699	6.6	0.13	6.1	0.12
Gallbladder	323	1.0	0.06	1.0	0.05	640	1.5	0.06	1.4	0.06
Pancreas	3,190	10.0	0.18	9.3	0.17	3,522	8.7	0.15	8.0	0.14
Respiratory System	15,259	48.3	0.41	45.0	0.37	11,854	28.7	0.27	26.9	0.25
Larynx	740	2.2	0.08	2.1	0.08	123	0.3	0.03	0.3	0.03
Lung & bronchus	14,156	45.2	0.40	42.0	0.36	11,509	27.9	0.27	26.2	0.25
Bones & Joints	330	0.8	0.04	0.8	0.04	228	0.5	0.03	0.5	0.03
Soft Tissue (incl. heart)	1,017	2.7	0.09	2.6	0.08	925	2.1	0.07	2.0	0.07
Skin (excl. basal & sq.)	839	2.4	0.09	2.3	0.08	786	1.8	0.06	1.7	0.06
Melanomas of the skin	534	1.5	0.07	1.5	0.06	530	1.2	0.05	1.1	0.05
Breast (excl. *in situ*)	201	0.6	0.04	0.6	0.04	42,016	90.7	0.45	88.8	0.44
Female Genital System	-	-	-	-	-	16,603	35.5	0.28	34.7	0.27
Cervix uteri	-	-	-	-	-	2,839	6.0	0.11	6.0	0.11
Uterus Corpus	-	-	-	-	-	8,382	17.6	0.20	17.2	0.19
Uterus, NOS[1]	-	-	-	-	-	244	0.6	0.04	0.5	0.03
Ovary	-	-	-	-	-	4,212	9.2	0.14	8.9	0.14
Male Genital System	21,104	61.0	0.44	59.3	0.42	-	-	-	-	-
Prostate	19,972	58.4	0.43	56.6	0.41	-	-	-	-	-
Testis	893	1.9	0.06	2.0	0.07	-	-	-	-	-
Penis	146	0.4	0.04	0.4	0.04	-	-	-	-	-
Urinary System	8,667	26.7	0.30	24.9	0.27	3,819	9.1	0.15	8.6	0.14
Bladder (incl. *in situ*)	4,516	15.0	0.23	13.7	0.21	1,509	3.8	0.10	3.4	0.09
Kidney & renal pelvis	3,892	10.9	0.18	10.5	0.17	2,115	4.9	0.11	4.7	0.10
Ureter	178	0.6	0.05	0.6	0.04	138	0.3	0.03	0.3	0.03
Eye & Orbit	121	0.3	0.03	0.3	0.03	118	0.3	0.03	0.3	0.02
Brain & Other Nervous System	1,626	4.2	0.11	4.1	0.10	1,433	3.2	0.09	3.1	0.08
Brain	1,494	3.9	0.10	3.8	0.10	1,313	3.0	0.08	2.9	0.08
Endocrine System	3,175	7.9	0.14	7.9	0.14	10,619	22.1	0.22	22.4	0.22
Thyroid gland	2,793	6.9	0.13	6.9	0.13	10,254	21.3	0.21	21.6	0.21
Lymphomas	6,017	17.2	0.23	16.4	0.22	5,022	11.5	0.17	11.1	0.16
Hodgkin Lymphoma	628	1.5	0.06	1.5	0.06	461	1.0	0.05	1.0	0.05
Non-Hodgkin Lymphoma	5,389	15.7	0.22	14.9	0.21	4,561	10.6	0.16	10.1	0.15
Multiple Myeloma	1,576	4.8	0.13	4.5	0.12	1,299	3.1	0.09	2.9	0.08
Leukemias	3,408	9.7	0.17	9.2	0.16	2,698	6.2	0.12	6.0	0.12
Acute lymphocytic	618	1.5	0.06	1.4	0.06	519	1.2	0.05	1.2	0.05
Chronic lymphocytic	543	1.7	0.07	1.6	0.07	301	0.7	0.04	0.7	0.04
Acute myeloid	1,351	4.0	0.11	3.7	0.10	1,252	2.9	0.08	2.8	0.08
Chronic myeloid	539	1.5	0.07	1.4	0.06	355	0.8	0.04	0.8	0.04
Other leukemias	357	1.1	0.06	1.0	0.05	271	0.6	0.04	0.6	0.04
Mesothelioma	169	0.6	0.05	0.5	0.04	69	0.2	0.02	0.2	0.02
Kaposi Sarcoma	134	0.3	0.03	0.3	0.03	8	0.0	0.01	0.0	0.01
Ill-defined & Unknown	3,820	12.4	0.21	11.3	0.19	3,651	9.0	0.15	8.2	0.14
Selected tumors (non-invasive)										
Brain & ONS[1] (*benign/borderline*)	2,410	6.6	0.14	6.4	0.13	5,129	11.6	0.17	11.1	0.16
Breast (*in situ*)	27	0.1	0.02	0.1	0.02	13,329	28.5	0.25	27.9	0.24

* Rates are per 100,000 population and are age-adjusted by five-year age groups to the 2000 U.S. standard population based on single years of age and to the 1996 Canadian population standard.

- Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for All Sites combined

[1] Abbreviations: NOS - Not Otherwise Specified; ONS - Other Nervous System.

Total Cancer Cases and Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates[1] for the Five Most Commonly Diagnosed Cancers
Selected Areas in the NAACCR United States Combined[2], 2010-2014, Asian/Pacific Islander, Males

	All Sites			Prostate			Lung & Bronchus			Colon & Rectum			Liver & IBD[3]			Non-Hodgkin Lymphoma		
	Count	*Rate*	*S.E.*[3]	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
NAACCR U.S. Combined	**102,876**	**302.8**	**0.99**	**19,972**	**58.4**	**0.43**	**14,156**	**45.2**	**0.40**	**12,803**	**37.0**	**0.34**	**7,146**	**20.0**	**0.25**	**5,389**	**15.7**	**0.22**
Alabama	272	294.2	21.32	67	72.1	9.96	29	34.2	7.61	21	17.8	4.41	23	27.1	6.30	13	10.5	3.04
Alaska	295	354.8	24.20	57	73.6	10.94	38	55.8	10.38	45	51.7	8.96	21	23.7	6.16	19	27.2	7.55
Arizona	856	228.9	8.51	150	40.4	3.54	122	36.5	3.58	105	28.4	3.03	59	14.4	1.99	41	10.8	1.82
Arkansas	414	651.9	38.07	56	96.7	15.29	45	80.3	13.21	46	61.5	10.28	29	36.7	7.54	18	29.8	8.92
California	39,205	306.7	1.59	7,730	59.0	0.69	5,608	46.2	0.63	5,132	39.5	0.57	2,804	21.0	0.41	2,124	16.6	0.37
CA: Greater California	13,671	286.6	2.54	2,664	54.8	1.10	1,977	44.3	1.03	1,743	35.9	0.89	1,004	19.8	0.65	709	14.8	0.58
CA: Greater Bay Area	13,867	329.7	2.86	2,911	68.4	1.30	2,011	50.3	1.15	1,708	39.8	0.98	1,003	22.7	0.74	786	18.5	0.67
CA: Los Angeles	11,654	305.9	2.89	2,151	53.8	1.18	1,620	43.8	1.11	1,681	43.7	1.08	795	20.5	0.74	628	16.6	0.68
Colorado	732	263.7	10.50	124	47.4	4.50	103	43.3	4.54	80	29.2	3.50	51	16.6	2.49	48	17.4	2.73
Connecticut	642	247.0	11.09	99	37.6	3.99	89	36.3	4.31	68	24.1	3.24	50	19.6	3.13	34	13.8	2.83
Delaware	159	266.3	22.89	35	55.5	9.66	16	33.9	9.44	13	20.1	5.87	10	16.8	5.57	10	18.6	6.61
District of Columbia	143	455.1	41.06	21	72.3	16.47	17	62.6	15.77	19	64.0	15.80	10	30.6	10.28	6	18.5	8.17
Florida	2,541	221.5	4.65	510	43.4	2.01	323	30.4	1.80	316	26.5	1.58	138	11.5	1.02	141	12.0	1.06
Georgia	1,640	271.1	7.77	285	50.2	3.33	181	32.3	2.84	226	36.0	2.77	129	20.3	2.06	82	12.8	1.68
Hawaii	10,385	400.5	3.99	2,251	85.8	1.84	1,511	58.0	1.52	1,452	55.3	1.48	487	18.5	0.85	520	20.4	0.91
Idaho	107	289.2	30.67	19	46.2	11.25	14	52.1	14.53	13	33.5	10.27	7	13.2	5.29	6	15.9	6.99
Illinois	3,631	283.1	5.10	741	56.5	2.24	486	41.2	2.02	466	36.2	1.82	198	15.2	1.17	182	14.7	1.18
Indiana	450	291.4	16.49	65	43.7	6.14	52	41.9	6.77	48	25.2	4.07	28	18.6	4.18	24	10.7	2.38
Iowa	257	343.6	26.07	35	51.4	10.06	29	51.6	11.08	32	45.0	9.46	37	52.2	10.42	15	19.5	5.83
Kansas	364	305.2	18.46	61	59.1	8.37	42	38.7	6.84	37	27.2	4.88	32	23.3	4.38	25	17.3	4.09
Kentucky	186	221.0	19.14	30	48.3	9.83	23	28.9	7.50	24	26.2	6.30	13	15.6	5.38	10	11.7	4.68
Louisiana	468	315.6	16.36	80	60.4	7.38	75	52.6	6.88	60	41.8	6.07	50	28.3	4.35	15	7.6	2.02
Maine	52	216.0	33.62	9	37.9	13.69	-	-	-	6	18.0	7.38	6	24.3	11.16	6	28.6	13.02
Maryland	1,950	262.8	6.34	403	52.1	2.73	235	35.3	2.46	208	27.8	2.05	130	17.0	1.58	100	13.3	1.39
Massachusetts	2,114	308.2	7.15	362	55.7	3.06	342	56.9	3.25	235	33.4	2.31	199	27.7	2.07	81	10.7	1.26
Michigan	1,213	267.2	8.52	313	68.5	4.14	126	31.0	3.08	150	36.0	3.30	62	12.2	1.69	65	13.0	1.78
MI: Metropolitan Detroit	846	291.8	11.09	247	83.9	5.72	86	32.3	3.88	96	34.5	3.95	39	12.5	2.18	40	11.9	2.06
Mississippi	151	311.1	28.73	27	62.5	13.79	27	54.9	11.35	18	37.3	9.66	15	30.6	9.06	6	14.0	6.58
Missouri	498	274.6	13.65	75	42.4	5.27	72	45.6	5.86	61	35.6	5.11	45	22.4	3.57	27	14.9	3.34

Total Cancer Cases and Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates[1] for the Five Most Commonly Diagnosed Cancers
Selected Areas in the NAACCR United States Combined[2], 2010-2014, Asian/Pacific Islander, Males

	All Sites			Prostate			Lung & Bronchus			Colon & Rectum			Liver & IBD[3]			Non-Hodgkin Lymphoma		
	Count	*Rate*	*S.E.*[3]	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
Montana	32	284.2	54.28	7	48.9	19.27	-	-	-	-	-	-	-	-	-	-	-	-
Nebraska	169	325.8	28.44	21	50.9	11.98	23	46.8	10.76	22	38.8	9.26	20	36.7	9.08	9	19.1	7.37
New Hampshire	110	240.6	26.50	28	64.0	13.68	12	31.9	10.00	10	21.4	7.38	6	11.0	5.34	-	-	-
New Jersey	4,326	272.9	4.58	1,004	61.8	2.11	469	33.4	1.69	469	28.3	1.44	223	14.1	1.05	219	13.2	0.98
New York	11,806	343.5	3.35	2,160	61.2	1.39	1,651	52.8	1.38	1,325	38.4	1.12	831	22.7	0.83	551	15.8	0.71
North Carolina	929	256.4	9.79	188	52.4	4.16	127	38.6	3.94	88	24.0	3.00	50	13.3	2.17	53	13.6	2.22
North Dakota	32	359.0	79.36	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Ohio	885	242.3	9.06	202	54.5	4.11	119	38.5	3.92	94	26.9	3.13	58	14.5	2.10	39	10.4	1.87
Oklahoma	431	363.9	19.76	72	61.1	7.90	62	56.3	7.85	68	56.7	8.00	31	26.7	5.31	22	16.8	4.06
Oregon	972	311.6	10.64	157	50.8	4.26	142	51.4	4.54	112	38.8	3.89	106	32.5	3.34	50	16.0	2.43
Pennsylvania	1,894	272.3	6.88	339	51.0	2.99	256	42.1	2.89	200	27.9	2.14	150	19.9	1.76	108	14.8	1.53
Rhode Island	155	272.7	24.43	29	52.3	10.48	16	32.4	9.11	14	26.9	8.00	17	29.1	7.63	11	22.1	7.41
South Carolina	303	244.8	16.46	53	47.3	7.31	52	42.1	6.79	36	28.9	5.53	11	8.7	2.98	25	21.7	5.58
South Dakota	29	314.1	69.54	-	-	-	6	87.7	42.23	6	35.5	15.84	-	-	-	-	-	-
Tennessee	494	295.4	15.60	97	64.0	7.22	54	36.7	5.91	62	37.9	5.61	37	19.6	3.53	33	20.1	4.28
Texas	4,779	246.1	4.04	816	44.1	1.72	644	37.9	1.69	580	29.4	1.40	444	21.3	1.13	271	13.3	0.91
Utah	472	345.1	17.61	118	94.2	9.28	38	34.6	6.04	42	30.9	5.22	32	22.5	4.41	23	15.7	3.74
Vermont	36	311.1	58.10	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Virginia	2,209	255.4	6.08	391	43.4	2.38	308	39.5	2.50	263	30.2	2.12	154	16.9	1.50	110	12.3	1.31
Washington	3,490	324.5	5.85	586	55.3	2.41	489	52.1	2.48	448	39.7	2.00	282	24.2	1.53	196	17.6	1.34
WA: Seattle/Puget Sound	3,274	338.9	6.29	558	59.1	2.63	457	53.6	2.64	423	41.8	2.16	258	24.9	1.64	189	18.7	1.44
West Virginia	54	203.9	32.82	10	30.9	10.06	13	43.2	13.36	11	44.3	15.30	-	-	-	-	-	-
Wisconsin	527	314.5	15.95	77	58.9	7.33	54	42.2	6.36	62	32.8	4.80	50	28.8	4.66	37	19.0	3.82
Wyoming	17	154.5	38.94	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-

1. Rates are per 100,000 population and are age-adjusted by five-year age groups to the 2000 U.S. standard population based on single years of age.
2. See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.
3. Abbreviations: IBD - Intrahepatic Bile Duct; S.E. - Standard Error

\- Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for the U.S. combined.

Total Cancer Cases and Average Annual Age-adjusted (1996 Canadian Standard) Cancer Incidence Rates[1] for the Five Most Commonly Diagnosed Cancers
Selected Areas in the NAACCR United States Combined[2], 2010-2014, Asian/Pacific Islander, Males

	All Sites			Prostate			Lung & Bronchus			Colon & Rectum			Liver & IBD[3]			Non-Hodgkin Lymphoma		
	Count	*Rate*	*S.E.*[3]	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
NAACCR U.S. Combined	**102,876**	**287.4**	**0.92**	**19,972**	**56.6**	**0.41**	**14,156**	**42.0**	**0.36**	**12,803**	**35.0**	**0.32**	**7,146**	**19.1**	**0.23**	**5,389**	**14.9**	**0.21**
Alabama	272	277.6	19.07	67	70.0	9.29	29	32.0	6.67	21	17.8	4.28	23	25.7	5.82	13	10.4	3.05
Alaska	295	337.8	22.03	57	72.0	10.35	38	52.1	9.20	45	49.7	8.30	21	22.1	5.50	19	24.4	6.44
Arizona	856	216.4	7.79	150	38.9	3.31	122	33.8	3.20	105	26.8	2.76	59	13.9	1.89	41	10.3	1.68
Arkansas	414	622.4	34.66	56	92.1	13.83	45	77.6	12.41	46	59.5	9.69	29	35.8	7.24	18	27.6	7.70
California	39,205	291.4	1.49	7,730	57.4	0.66	5,608	42.9	0.58	5,132	37.5	0.53	2,804	20.1	0.39	2,124	15.8	0.35
CA: Greater California	13,671	272.4	2.37	2,664	53.3	1.05	1,977	41.3	0.94	1,743	34.1	0.83	1,004	19.0	0.61	709	14.1	0.54
CA: Greater Bay Area	13,867	313.2	2.70	2,911	66.4	1.25	2,011	46.7	1.06	1,708	37.7	0.93	1,003	21.8	0.70	786	17.7	0.64
CA: Los Angeles	11,654	290.7	2.73	2,151	52.6	1.15	1,620	40.7	1.02	1,681	41.3	1.02	795	19.5	0.70	628	15.8	0.64
Colorado	732	249.5	9.70	124	45.7	4.24	103	39.6	4.04	80	27.4	3.20	51	15.7	2.31	48	16.4	2.51
Connecticut	642	236.3	10.15	99	37.1	3.89	89	34.5	3.94	68	23.0	3.02	50	18.9	2.89	34	13.1	2.50
Delaware	159	250.9	20.83	35	54.2	9.33	16	29.6	7.93	13	18.6	5.38	10	16.0	5.23	10	16.7	5.65
District of Columbia	143	431.3	38.39	21	69.5	15.68	17	60.0	14.95	19	61.4	15.02	10	30.2	10.09	6	17.5	7.67
Florida	2,541	211.7	4.33	510	42.6	1.93	323	28.3	1.63	316	25.3	1.47	138	11.1	0.97	141	11.6	1.00
Georgia	1,640	258.2	7.03	285	48.5	3.08	181	30.4	2.51	226	34.4	2.52	129	19.3	1.87	82	12.0	1.49
Hawaii	10,385	381.9	3.82	2,251	82.8	1.77	1,511	54.7	1.43	1,452	53.0	1.42	487	17.6	0.81	520	19.4	0.87
Idaho	107	268.0	27.74	19	44.7	10.76	14	46.8	12.86	13	30.0	9.07	7	12.8	4.92	6	14.3	6.19
Illinois	3,631	268.5	4.66	741	54.7	2.09	486	38.3	1.82	466	34.1	1.65	198	14.3	1.07	182	13.7	1.07
Indiana	450	270.7	14.47	65	41.4	5.58	52	37.6	5.76	48	24.8	3.91	28	17.1	3.63	24	10.6	2.35
Iowa	257	319.9	22.99	35	49.7	9.11	29	46.4	9.51	32	41.4	8.32	37	48.2	9.07	15	17.9	5.25
Kansas	364	293.2	16.87	61	56.7	7.71	42	37.2	6.24	37	27.0	4.72	32	23.1	4.28	25	17.0	3.85
Kentucky	186	210.9	17.33	30	45.0	8.83	23	26.6	6.38	24	25.4	5.78	13	15.0	4.77	10	11.1	4.23
Louisiana	468	300.4	15.05	80	58.0	6.90	75	49.2	6.18	60	39.5	5.54	50	27.7	4.21	15	7.7	2.07
Maine	52	205.5	30.84	9	37.6	13.33	-	-	-	6	18.2	7.46	6	21.1	9.22	6	26.6	11.73
Maryland	1,950	249.3	5.83	403	50.7	2.59	235	32.5	2.19	208	26.3	1.89	130	16.0	1.45	100	12.9	1.32
Massachusetts	2,114	293.4	6.65	362	53.8	2.91	342	52.7	2.94	235	31.9	2.17	199	26.5	1.95	81	10.4	1.21
Michigan	1,213	254.0	7.76	313	66.5	3.91	126	28.7	2.73	150	33.4	2.93	62	11.9	1.59	65	12.5	1.66
MI: Metropolitan Detroit	846	278.6	10.18	247	81.6	5.42	86	30.4	3.50	96	32.3	3.54	39	12.1	2.04	40	11.5	1.93
Mississippi	151	298.1	26.47	27	59.0	12.32	27	54.4	11.08	18	36.0	9.13	15	28.8	8.27	6	13.6	6.09
Missouri	498	260.1	12.46	75	41.5	4.99	72	43.1	5.32	61	32.3	4.45	45	21.3	3.34	27	13.9	2.98

Total Cancer Cases and Average Annual Age-adjusted (1996 Canadian Standard) Cancer Incidence Rates[1] for the Five Most Commonly Diagnosed Cancers
Selected Areas in the NAACCR United States Combined[2], 2010-2014, Asian/Pacific Islander, Males

	All Sites			Prostate			Lung & Bronchus			Colon & Rectum			Liver & IBD[3]			Non-Hodgkin Lymphoma		
	Count	*Rate*	*S.E.*[3]	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
Montana	32	269.4	50.56	7	50.4	19.86	-	-	-	-	-	-	-	-	-	-	-	-
Nebraska	169	307.1	25.87	21	47.5	10.94	23	44.9	10.02	22	36.9	8.48	20	35.5	8.56	9	16.9	6.18
New Hampshire	110	229.2	24.32	28	62.8	12.81	12	29.8	9.18	10	21.1	7.20	6	10.4	4.84	-	-	-
New Jersey	4,326	259.9	4.19	1,004	59.8	1.98	469	31.3	1.52	469	27.1	1.33	223	13.3	0.95	219	12.7	0.91
New York	11,806	327.3	3.12	2,160	59.6	1.32	1,651	49.1	1.25	1,325	36.4	1.04	831	21.8	0.78	551	15.1	0.67
North Carolina	929	242.9	8.83	188	51.3	3.95	127	35.8	3.50	88	22.7	2.70	50	12.6	1.97	53	13.0	2.01
North Dakota	32	341.2	72.49	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Ohio	885	228.8	8.21	202	52.9	3.87	119	35.4	3.45	94	24.9	2.75	58	13.8	1.92	39	9.9	1.71
Oklahoma	431	348.2	18.17	72	60.1	7.49	62	53.7	7.28	68	53.0	7.07	31	25.4	4.92	22	16.5	3.81
Oregon	972	296.9	9.92	157	49.9	4.10	142	48.1	4.16	112	36.0	3.53	106	31.3	3.17	50	15.0	2.23
Pennsylvania	1,894	258.7	6.31	339	49.1	2.79	256	38.8	2.57	200	26.5	1.98	150	19.2	1.66	108	14.3	1.45
Rhode Island	155	261.3	22.69	29	51.3	10.09	16	30.5	8.26	14	26.3	7.53	17	28.8	7.40	11	20.2	6.63
South Carolina	303	233.6	14.84	53	45.0	6.68	52	40.8	6.19	36	27.6	5.02	11	8.4	2.80	25	19.8	4.63
South Dakota	29	304.8	65.18	-	-	-	6	81.0	36.53	6	36.9	16.75	-	-	-	-	-	-
Tennessee	494	280.5	14.06	97	61.4	6.68	54	34.1	5.18	62	36.0	5.09	37	19.0	3.37	33	18.8	3.73
Texas	4,779	231.6	3.64	816	42.0	1.57	644	34.9	1.49	580	27.5	1.25	444	20.3	1.04	271	12.6	0.83
Utah	472	329.3	16.37	118	91.7	8.86	38	31.7	5.42	42	29.3	4.87	32	21.4	4.06	23	15.0	3.45
Vermont	36	303.0	54.51	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Virginia	2,209	240.4	5.47	391	42.1	2.23	308	36.5	2.22	263	28.1	1.88	154	16.0	1.37	110	11.7	1.19
Washington	3,490	306.4	5.38	586	53.2	2.26	489	47.8	2.23	448	37.4	1.83	282	23.1	1.43	196	16.7	1.24
WA: Seattle/Puget Sound	3,274	320.0	5.80	558	56.9	2.48	457	49.3	2.38	423	39.3	1.98	258	23.7	1.53	189	17.8	1.34
West Virginia	54	190.8	28.72	10	30.2	9.65	13	42.5	12.64	11	41.2	13.78	-	-	-	-	-	-
Wisconsin	527	294.4	14.37	77	54.7	6.56	54	38.3	5.58	62	31.3	4.43	50	26.9	4.17	37	18.1	3.46
Wyoming	17	150.4	37.31	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-

1. Rates are per 100,000 and are age-adjusted to the 1996 Canadian population standard.
2. See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.
3. Abbreviations: IBD - Intrahepatic Bile Duct; S.E. - Standard Error

\- Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for the U.S. combined.

Total Cancer Cases and Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates[1] for the Five Most Commonly Diagnosed Cancers
Selected Areas in the NAACCR United States Combined[2], 2010-2014, Asian/Pacific Islander, Females

	All Sites			Breast			Colon & Rectum			Lung & Bronchus			Thyroid			Uterus Corpus & Uterus, NOS[3]		
	Count	*Rate*	*S.E.*[3]	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
NAACCR U.S. Combined	**128,044**	**287.6**	**0.82**	**42,016**	**90.7**	**0.45**	**11,595**	**27.0**	**0.26**	**11,509**	**27.9**	**0.27**	**10,254**	**21.3**	**0.21**	**8,626**	**18.2**	**0.20**
Alabama	400	280.7	15.33	136	87.4	8.03	33	24.6	4.70	42	35.5	5.88	24	12.5	2.57	15	9.3	2.55
Alaska	374	300.2	16.31	120	92.3	8.77	28	22.6	4.48	43	38.4	6.19	33	25.0	4.46	34	23.3	4.14
Arizona	1,402	254.0	7.12	450	75.6	3.68	99	19.9	2.10	150	30.8	2.65	134	21.0	1.86	85	13.5	1.52
Arkansas	503	534.3	26.12	119	126.9	12.67	48	53.5	8.54	51	62.1	9.47	26	24.5	5.36	25	24.4	5.30
California	48,846	293.9	1.35	16,207	95.7	0.76	4,796	29.0	0.42	4,325	26.6	0.41	3,169	19.1	0.34	3,467	19.9	0.34
CA: Greater California	17,690	282.2	2.15	5,743	88.5	1.18	1,709	27.9	0.68	1,552	26.0	0.67	1,271	19.7	0.56	1,221	18.3	0.53
CA: Greater Bay Area	16,477	304.3	2.40	5,585	101.4	1.37	1,516	28.0	0.73	1,503	28.5	0.74	936	17.2	0.57	1,190	21.0	0.62
CA: Los Angeles	14,672	297.8	2.50	4,876	98.8	1.44	1,571	31.2	0.80	1,269	25.5	0.72	962	20.6	0.68	1,057	20.8	0.65
Colorado	1,044	251.3	8.06	339	76.1	4.25	99	27.5	2.85	106	28.5	2.86	112	22.6	2.18	66	14.1	1.77
Connecticut	845	260.3	9.81	314	89.2	5.41	69	25.6	3.38	64	26.8	3.60	97	23.3	2.54	40	11.4	1.90
Delaware	217	278.7	20.09	74	85.9	10.19	10	15.7	5.32	17	26.6	6.93	29	31.4	5.96	10	12.5	4.16
District of Columbia	184	338.7	26.50	52	90.4	13.15	13	24.0	6.93	17	35.2	9.05	28	34.5	7.18	9	16.2	5.48
Florida	3,506	224.6	3.93	1,165	70.1	2.10	284	18.6	1.15	323	22.3	1.29	260	15.5	0.98	237	14.0	0.94
Georgia	1,990	248.7	6.14	662	74.4	3.09	171	23.9	2.04	137	20.3	1.89	165	16.4	1.32	107	12.2	1.25
Hawaii	11,920	382.6	3.62	4,015	132.1	2.14	1,177	36.3	1.09	1,207	35.9	1.06	621	23.9	0.98	942	32.1	1.07
Idaho	169	283.2	22.81	50	76.0	11.13	13	22.3	6.47	18	36.2	8.90	9	13.3	4.58	13	19.0	5.39
Illinois	4,505	271.6	4.20	1,641	92.7	2.35	394	25.7	1.35	325	21.8	1.26	406	21.7	1.10	324	18.2	1.04
Indiana	570	247.1	11.38	170	66.1	5.44	54	29.7	4.43	48	25.2	3.95	43	13.9	2.24	42	16.8	2.76
Iowa	272	256.1	17.31	80	68.2	8.07	19	20.0	5.11	23	25.8	6.02	22	13.1	3.06	19	15.7	3.87
Kansas	449	274.9	14.09	115	63.2	6.22	40	26.8	4.61	61	46.6	6.37	57	26.6	3.65	29	17.5	3.51
Kentucky	310	260.6	16.23	89	64.4	7.21	34	35.1	6.56	36	39.8	7.16	27	17.8	3.52	22	16.1	3.56
Louisiana	466	256.9	12.72	117	62.6	6.10	30	17.8	3.52	47	30.5	4.74	40	17.2	2.76	32	14.0	2.49
Maine	105	300.6	32.87	28	76.3	15.97	8	24.5	9.06	13	50.0	14.94	19	38.8	8.96	6	16.6	7.19
Maryland	2,510	264.3	5.47	861	84.3	2.95	212	24.1	1.73	185	22.1	1.69	264	25.4	1.59	160	15.6	1.27
Massachusetts	2,563	294.8	6.08	816	87.7	3.17	221	27.7	1.95	244	32.6	2.17	280	27.0	1.67	118	12.9	1.22
Michigan	1,512	264.0	7.23	534	85.3	3.86	104	19.6	2.06	152	31.5	2.71	110	15.9	1.58	106	17.0	1.72
MI: Metropolitan Detroit	958	275.5	9.45	358	94.8	5.24	68	21.4	2.78	80	27.9	3.30	78	19.6	2.31	67	17.9	2.30
Mississippi	194	282.2	22.22	47	60.6	9.41	17	26.7	7.21	28	53.4	10.83	17	17.3	4.23	9	9.3	3.12
Missouri	746	297.8	11.69	238	88.2	6.03	60	28.3	3.93	72	33.8	4.20	61	19.0	2.53	40	14.2	2.33

Total Cancer Cases and Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates[1] for the Five Most Commonly Diagnosed Cancers
Selected Areas in the NAACCR United States Combined[2], 2010-2014, Asian/Pacific Islander, Females

	All Sites			Breast			Colon & Rectum			Lung & Bronchus			Thyroid			Uterus Corpus & Uterus, NOS[3]		
	Count	*Rate*	*S.E.*[3]	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
Montana	78	344.2	40.81	27	112.5	22.63	6	29.0	12.30	9	41.4	14.42	-	-	-	9	35.5	12.20
Nebraska	200	262.4	20.22	51	65.6	9.82	18	24.6	6.24	18	31.2	7.95	30	31.7	6.34	11	15.0	4.95
New Hampshire	168	250.6	21.62	55	72.1	10.50	16	24.6	6.88	12	21.0	6.78	17	18.5	4.54	15	24.4	6.89
New Jersey	5,260	268.5	3.90	1,938	92.2	2.17	401	22.3	1.18	377	22.5	1.22	509	22.3	1.01	352	16.4	0.91
New York	13,341	314.7	2.80	4,076	92.2	1.47	1,106	27.5	0.85	1,173	30.3	0.91	1,567	33.5	0.86	858	19.2	0.67
North Carolina	1,304	257.8	7.86	434	77.7	4.03	94	19.8	2.25	126	30.3	2.91	109	15.9	1.62	81	14.9	1.78
North Dakota	39	291.9	55.10	-	-	-	-	-	-	-	-	-	9	51.7	21.75	-	-	-
Ohio	1,173	239.1	7.40	419	79.8	4.07	85	19.0	2.20	109	26.5	2.68	119	19.6	1.85	58	11.0	1.50
Oklahoma	564	324.8	14.71	159	80.5	6.63	62	41.9	5.74	65	44.9	5.95	29	16.0	3.30	44	22.8	3.61
Oregon	1,284	281.7	8.16	380	76.9	4.04	123	28.9	2.71	120	31.5	2.97	109	21.2	2.08	75	14.2	1.67
Pennsylvania	2,350	271.1	5.95	699	73.8	2.91	184	23.2	1.82	211	28.5	2.09	262	24.1	1.54	113	11.8	1.15
Rhode Island	169	239.4	19.71	52	68.9	10.01	15	24.0	6.57	13	21.1	6.24	10	11.2	3.57	14	17.5	4.82
South Carolina	464	266.8	13.35	145	76.3	6.66	39	24.8	4.30	45	30.9	4.95	32	15.3	2.72	25	12.7	2.61
South Dakota	57	297.9	43.28	13	72.8	22.26	7	44.7	18.15	-	-	-	-	-	-	-	-	-
Tennessee	596	250.1	11.31	182	70.2	5.62	59	26.4	3.79	64	33.4	4.57	43	15.2	2.50	29	11.2	2.20
Texas	5,794	225.8	3.20	1,854	65.0	1.59	531	22.1	1.04	515	23.7	1.12	503	16.2	0.75	337	11.7	0.67
Utah	597	326.2	14.19	175	91.2	7.28	59	36.0	4.93	45	31.0	4.83	45	19.6	3.03	72	34.3	4.19
Vermont	43	249.2	42.79	14	89.6	26.76	-	-	-	8	59.4	22.56	-	-	-	-	-	-
Virginia	3,134	252.2	4.77	1,057	79.0	2.53	260	22.5	1.48	272	25.9	1.67	368	25.2	1.35	170	12.3	0.98
Washington	5,004	317.8	4.60	1,618	98.3	2.49	424	27.5	1.37	513	35.6	1.61	352	20.3	1.10	331	19.4	1.09
WA: Seattle/Puget Sound	4,642	328.2	4.92	1,483	100.3	2.65	408	29.4	1.49	474	36.7	1.72	334	21.5	1.19	308	20.2	1.17
West Virginia	87	221.9	25.03	33	77.8	14.01	-	-	-	6	18.6	8.13	9	21.8	7.29	7	17.6	7.12
Wisconsin	710	302.2	12.19	184	74.2	5.74	62	31.5	4.23	62	32.8	4.36	66	20.7	2.73	60	22.4	3.04
Wyoming	26	196.2	39.91	8	58.4	21.46	-	-	-	-	-	-	-	-	-	-	-	-

1. Rates are per 100,000 population and are age-adjusted by five-year age groups to the 2000 U.S. standard population based on single years of age.
2. See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.
3. Abbreviations: NOS - Not Otherwise Specified; S.E. - Standard Error

\- Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for the U.S. combined.

Total Cancer Cases and Average Annual Age-adjusted (1996 Canadian Standard) Cancer Incidence Rates[1] for the Five Most Commonly Diagnosed Cancers
Selected Areas in the NAACCR United States Combined[2], 2010-2014, Asian/Pacific Islander, Females

	All Sites			Breast			Colon & Rectum			Lung & Bronchus			Thyroid			Uterus Corpus & Uterus, NOS[3]		
	Count	*Rate*	*S.E.*[3]	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
NAACCR U.S. Combined	**128,044**	**277.1**	**0.78**	**42,016**	**88.8**	**0.44**	**11,595**	**25.4**	**0.24**	**11,509**	**26.2**	**0.25**	**10,254**	**21.6**	**0.21**	**8,626**	**17.8**	**0.19**
Alabama	400	266.8	14.10	136	84.7	7.59	33	23.1	4.26	42	33.0	5.32	24	12.9	2.65	15	8.8	2.38
Alaska	374	293.2	15.74	120	90.5	8.56	28	22.1	4.32	43	36.2	5.71	33	25.9	4.61	34	23.4	4.14
Arizona	1,402	242.4	6.65	450	73.6	3.55	99	18.4	1.90	150	28.3	2.38	134	21.2	1.85	85	13.3	1.47
Arkansas	503	512.4	24.41	119	121.7	11.85	48	49.6	7.72	51	57.9	8.62	26	23.8	5.01	25	23.4	4.94
California	48,846	283.3	1.30	16,207	93.9	0.75	4,796	27.3	0.40	4,325	24.9	0.38	3,169	19.4	0.35	3,467	19.4	0.33
CA: Greater California	17,690	272.3	2.07	5,743	86.9	1.16	1,709	26.3	0.64	1,552	24.3	0.62	1,271	19.9	0.57	1,221	17.9	0.52
CA: Greater Bay Area	16,477	292.8	2.31	5,585	99.3	1.35	1,516	26.4	0.69	1,503	26.5	0.69	936	17.5	0.58	1,190	20.4	0.60
CA: Los Angeles	14,672	287.6	2.43	4,876	97.0	1.42	1,571	29.5	0.76	1,269	24.0	0.69	962	20.8	0.69	1,057	20.4	0.64
Colorado	1,044	240.3	7.63	339	74.1	4.11	99	25.7	2.64	106	26.7	2.66	112	22.6	2.17	66	13.7	1.72
Connecticut	845	249.9	9.14	314	86.4	5.14	69	23.5	3.01	64	24.5	3.21	97	23.6	2.50	40	11.2	1.84
Delaware	217	268.3	18.77	74	85.2	10.03	10	14.7	4.82	17	23.8	6.04	29	32.6	6.12	10	12.4	4.05
District of Columbia	184	325.4	25.11	52	88.3	12.71	13	23.6	6.78	17	33.1	8.40	28	35.4	7.19	9	15.2	5.08
Florida	3,506	218.2	3.76	1,165	69.3	2.06	284	17.8	1.08	323	21.2	1.21	260	15.7	0.98	237	13.8	0.92
Georgia	1,990	238.7	5.69	662	72.8	2.96	171	22.5	1.84	137	19.0	1.72	165	16.5	1.32	107	12.1	1.22
Hawaii	11,920	371.0	3.56	4,015	129.4	2.12	1,177	34.6	1.06	1,207	34.1	1.02	621	24.4	1.01	942	31.7	1.07
Idaho	169	269.1	21.48	50	74.9	10.91	13	20.0	5.71	18	34.6	8.46	9	13.0	4.51	13	18.5	5.28
Illinois	4,505	262.8	4.00	1,641	91.4	2.29	394	24.2	1.25	325	20.5	1.16	406	22.1	1.11	324	18.0	1.02
Indiana	570	236.4	10.55	170	64.6	5.21	54	26.6	3.82	48	23.5	3.57	43	14.0	2.21	42	16.4	2.66
Iowa	272	241.9	15.80	80	67.0	7.86	19	18.6	4.60	23	23.1	5.16	22	13.5	3.06	19	14.9	3.60
Kansas	449	265.4	13.22	115	62.9	6.08	40	25.5	4.25	61	43.2	5.80	57	27.7	3.78	29	16.3	3.21
Kentucky	310	250.3	15.06	89	63.8	7.03	34	32.6	5.89	36	36.4	6.35	27	17.6	3.47	22	15.9	3.48
Louisiana	466	246.9	11.94	117	60.7	5.81	30	16.4	3.17	47	28.4	4.33	40	17.8	2.84	32	14.1	2.52
Maine	105	285.2	29.89	28	72.1	14.58	8	24.6	9.03	13	45.5	13.13	19	39.2	9.07	6	15.7	6.71
Maryland	2,510	254.5	5.19	861	83.0	2.88	212	22.5	1.58	185	20.7	1.55	264	25.4	1.59	160	15.3	1.23
Massachusetts	2,563	284.8	5.79	816	86.5	3.11	221	25.9	1.79	244	30.6	2.01	280	27.2	1.66	118	12.6	1.19
Michigan	1,512	252.6	6.75	534	83.5	3.72	104	18.1	1.85	152	29.4	2.47	110	15.9	1.55	106	16.5	1.65
MI: Metropolitan Detroit	958	263.4	8.82	358	92.1	5.01	68	19.8	2.50	80	25.9	2.99	78	19.6	2.27	67	17.5	2.20
Mississippi	194	264.9	20.13	47	58.7	8.92	17	24.0	6.24	28	47.5	9.38	17	18.2	4.44	9	9.1	3.06
Missouri	746	284.5	10.85	238	85.8	5.75	60	26.1	3.50	72	31.6	3.85	61	19.6	2.57	40	13.9	2.27

Total Cancer Cases and Average Annual Age-adjusted (1996 Canadian Standard) Cancer Incidence Rates[1] for the Five Most Commonly Diagnosed Cancers
Selected Areas in the NAACCR United States Combined[2], 2010-2014, Asian/Pacific Islander, Females

	All Sites			Breast			Colon & Rectum			Lung & Bronchus			Thyroid			Uterus Corpus & Uterus, NOS[3]		
	Count	*Rate*	*S.E.*[3]	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
Montana	78	335.0	39.54	27	108.2	21.75	6	28.4	12.05	9	39.7	13.64	-	-	-	9	33.0	11.28
Nebraska	200	253.2	19.09	51	63.4	9.37	18	22.8	5.66	18	29.0	7.20	30	32.2	6.31	11	14.1	4.55
New Hampshire	168	239.6	20.07	55	71.0	10.20	16	23.1	6.18	12	20.0	6.25	17	18.8	4.60	15	23.3	6.53
New Jersey	5,260	258.6	3.67	1,938	90.2	2.10	401	20.9	1.08	377	20.9	1.11	509	22.8	1.02	352	16.0	0.87
New York	13,341	304.9	2.69	4,076	90.5	1.44	1,106	26.0	0.80	1,173	28.5	0.85	1,567	34.0	0.87	858	18.7	0.65
North Carolina	1,304	247.6	7.32	434	75.4	3.82	94	18.8	2.07	126	28.8	2.70	109	16.1	1.60	81	14.6	1.71
North Dakota	39	276.6	50.00	-	-	-	-	-	-	-	-	-	9	51.4	20.13	-	-	-
Ohio	1,173	229.2	6.93	419	77.6	3.89	85	18.0	2.02	109	24.8	2.45	119	19.9	1.85	58	10.6	1.43
Oklahoma	564	311.4	13.71	159	79.4	6.49	62	38.5	5.11	65	41.7	5.37	29	15.4	3.03	44	22.4	3.50
Oregon	1,284	270.6	7.74	380	75.9	3.97	123	27.1	2.51	120	28.8	2.69	109	21.4	2.08	75	14.0	1.64
Pennsylvania	2,350	259.5	5.56	699	72.3	2.81	184	21.7	1.66	211	26.2	1.87	262	24.6	1.55	113	11.4	1.10
Rhode Island	169	228.4	18.42	52	65.5	9.42	15	22.4	6.06	13	19.5	5.64	10	11.7	3.72	14	17.8	4.91
South Carolina	464	255.7	12.46	145	74.7	6.42	39	22.9	3.87	45	28.6	4.47	32	15.8	2.81	25	12.5	2.59
South Dakota	57	294.8	42.22	13	68.9	20.48	7	40.4	16.22	-	-	-	-	-	-	-	-	-
Tennessee	596	242.1	10.54	182	68.8	5.35	59	25.3	3.51	64	31.2	4.11	43	15.2	2.42	29	11.0	2.13
Texas	5,794	216.5	2.98	1,854	63.7	1.53	531	20.8	0.95	515	22.3	1.02	503	16.2	0.74	337	11.5	0.65
Utah	597	314.4	13.50	175	88.3	6.95	59	34.4	4.68	45	28.6	4.44	45	20.1	3.11	72	34.2	4.17
Vermont	43	237.6	39.45	14	85.6	24.74	-	-	-	8	56.0	20.83	-	-	-	-	-	-
Virginia	3,134	243.5	4.50	1,057	77.3	2.44	260	21.4	1.38	272	24.3	1.52	368	25.6	1.35	170	12.2	0.96
Washington	5,004	305.2	4.38	1,618	96.3	2.43	424	26.1	1.29	513	33.3	1.49	352	20.4	1.10	331	19.1	1.07
WA: Seattle/Puget Sound	4,642	314.8	4.69	1,483	98.2	2.58	408	27.9	1.40	474	34.3	1.60	334	21.6	1.19	308	19.7	1.14
West Virginia	87	216.4	23.85	33	77.7	13.73	-	-	-	6	17.4	7.41	9	21.1	7.10	7	17.2	6.77
Wisconsin	710	290.2	11.49	184	73.1	5.60	62	28.8	3.81	62	31.3	4.10	66	21.3	2.80	60	21.9	2.93
Wyoming	26	187.5	37.78	8	55.4	20.20	-	-	-	-	-	-	-	-	-	-	-	-

1. Rates are per 100,000 and are age-adjusted to the 1996 Canadian population standard.
2. See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.
3. Abbreviations: NOS - Not Otherwise Specified; S.E. - Standard Error

\- Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for the U.S. combined.

NAACCR U.S. Combined Incidence 2010-2014,
The American Indian/Alaskan Native CHSDA Counties Population

UNITED STATES

III-43 Combined Incidence – American Indian/Alaskan Native

Total Cancer Cases and Average Annual Age-adjusted Cancer Incidence Rates*
NAACCR United States Combined, 2010-2014, American Indian/Alaskan Native (AI/AN) for CHSDA[1] Counties

	MALES (AI/AN)					FEMALES (AI/AN)				
SITES	*Count*	*Adj. Rate 2000 U.S.*	*S.E.*	*Adj. Rate 1996 Cdn*	*S.E.*	*Count*	*Adj. Rate 2000 U.S.*	*S.E.*	*Adj. Rate 1996 Cdn*	*S.E.*
All Sites (invasive & in situ bladder)	12,813	425.3	4.17	406.0	3.85	14,277	388.7	3.41	374.0	3.23
Oral Cavity & Pharynx	554	16.3	0.76	15.6	0.71	228	6.1	0.43	5.8	0.40
Lip	26	1.0	0.22	0.9	0.19	-	-	-	-	-
Tongue	135	3.7	0.34	3.6	0.33	59	1.5	0.20	1.5	0.20
Major salivary gland	37	1.0	0.19	1.0	0.18	37	1.0	0.17	0.9	0.16
Floor of mouth	38	1.1	0.20	1.0	0.18	13	0.4	0.11	0.3	0.10
Gum & other mouth	53	1.6	0.24	1.5	0.22	35	1.0	0.18	0.9	0.16
Nasopharynx	42	1.2	0.21	1.2	0.19	24	0.7	0.15	0.6	0.13
Oropharynx	30	0.9	0.17	0.9	0.17	7	0.2	0.07	0.2	0.06
Hypopharynx	53	1.8	0.27	1.7	0.25	11	0.3	0.10	0.3	0.09
Digestive System	3,383	110.4	2.10	105.1	1.94	2,776	79.2	1.58	74.9	1.47
Esophagus	236	7.8	0.56	7.4	0.52	80	2.3	0.27	2.2	0.25
Stomach	329	11.6	0.71	10.9	0.64	222	6.5	0.46	6.2	0.43
Small intestine	53	1.7	0.26	1.6	0.24	47	1.3	0.20	1.3	0.19
Colon excl. rectum	964	33.4	1.20	31.6	1.09	1,030	30.0	0.98	28.4	0.91
Rectum & rectosigmoid	549	16.7	0.78	16.0	0.73	433	11.3	0.57	10.8	0.54
Anus, anal canal & anorectum	38	1.0	0.17	1.0	0.17	61	1.5	0.20	1.4	0.19
Liver	657	19.0	0.80	18.3	0.76	293	8.1	0.50	7.6	0.46
Gallbladder	57	2.4	0.35	2.2	0.31	95	3.0	0.32	2.7	0.29
Pancreas	358	12.0	0.70	11.5	0.65	359	10.8	0.60	10.0	0.55
Respiratory System	2,107	78.1	1.87	73.8	1.71	1,930	57.2	1.36	54.5	1.28
Larynx	159	5.1	0.44	4.9	0.41	35	0.8	0.15	0.8	0.15
Lung & bronchus	1,911	71.9	1.80	67.8	1.64	1,875	55.9	1.35	53.1	1.26
Bones & Joints	64	1.6	0.22	1.5	0.20	32	0.7	0.12	0.7	0.12
Soft Tissue (incl. heart)	110	3.1	0.33	3.1	0.31	98	2.4	0.26	2.3	0.25
Skin (excl. basal & sq.)	282	9.7	0.65	9.1	0.58	256	6.8	0.44	6.7	0.43
Melanomas of the skin	251	8.6	0.61	8.0	0.55	210	5.6	0.40	5.4	0.39
Breast (excl. *in situ*)	15	0.5	0.14	0.5	0.13	3,839	100.7	1.69	98.0	1.63
Female Genital System	-	-	-	-	-	1,808	46.0	1.13	45.2	1.09
Cervix uteri	-	-	-	-	-	374	9.1	0.49	9.2	0.49
Uterus Corpus	-	-	-	-	-	872	21.7	0.76	21.4	0.74
Uterus, NOS[1]	-	-	-	-	-	24	0.6	0.14	0.6	0.13
Ovary	-	-	-	-	-	399	10.6	0.56	10.3	0.53
Male Genital System	2,570	83.8	1.82	81.6	1.71	-	-	-	-	-
Prostate	2,311	78.3	1.78	75.6	1.67	-	-	-	-	-
Testis	243	4.9	0.32	5.3	0.35	-	-	-	-	-
Penis	14	0.5	0.15	0.5	0.14	-	-	-	-	-
Urinary System	1,509	50.5	1.44	48.0	1.32	878	23.8	0.84	22.8	0.79
Bladder (incl. *in situ*)	489	19.4	0.98	17.9	0.86	194	5.9	0.45	5.5	0.41
Kidney & renal pelvis	994	30.0	1.03	29.2	0.98	670	17.4	0.70	16.9	0.67
Ureter	16	0.6	0.16	0.5	0.15	8	0.3	0.10	0.2	0.09
Eye & Orbit	17	0.5	0.14	0.5	0.13	19	0.4	0.10	0.4	0.09
Brain & Other Nervous System	211	5.4	0.42	5.3	0.40	154	3.6	0.31	3.5	0.30
Brain	192	4.9	0.40	4.8	0.38	142	3.4	0.30	3.3	0.29
Endocrine System	188	4.9	0.38	4.9	0.37	642	15.1	0.62	15.3	0.62
Thyroid gland	164	4.4	0.37	4.4	0.36	618	14.5	0.60	14.7	0.61
Lymphomas	565	18.2	0.86	17.3	0.79	516	14.5	0.67	13.9	0.63
Hodgkin Lymphoma	63	1.5	0.20	1.5	0.19	43	1.0	0.17	1.0	0.16
Non-Hodgkin Lymphoma	502	16.7	0.84	15.9	0.77	473	13.5	0.65	12.9	0.61
Multiple Myeloma	214	7.8	0.59	7.3	0.53	173	5.0	0.40	4.8	0.38
Leukemias	417	12.7	0.71	12.1	0.65	353	9.0	0.51	8.7	0.48
Acute lymphocytic	74	1.4	0.18	1.4	0.17	69	1.4	0.17	1.4	0.17
Chronic lymphocytic	95	3.5	0.39	3.2	0.36	58	1.8	0.25	1.7	0.22
Acute myeloid	136	4.3	0.41	4.1	0.39	128	3.2	0.30	3.2	0.29
Chronic myeloid	55	1.6	0.24	1.6	0.23	60	1.5	0.20	1.5	0.20
Other leukemias	57	1.9	0.29	1.8	0.26	38	1.1	0.19	1.0	0.17
Mesothelioma	27	1.3	0.28	1.2	0.24	9	0.3	0.10	0.3	0.09
Kaposi Sarcoma	18	0.6	0.15	0.5	0.13	-	-	-	-	-
Ill-defined & Unknown	562	20.0	0.95	18.9	0.86	-	17.6	0.78	16.2	0.70
Selected tumors (non-invasive)										
Brain & ONS[1] (*benign/borderline*)	249	7.4	0.53	7.0	0.48	539	14.1	0.63	13.9	0.62
Breast (*in situ*)	6	0.2	0.07	0.2	0.08	737	18.7	0.71	18.3	0.69

* Rates are per 100,000 population and are age-adjusted by five-year age groups to the 2000 U.S. standard population based on single years of age and to the 1996 Canadian population standard. Statistics exclude data for KS for all years.

\- Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for All Sites combined.

[1] CHSDA - Indian Health Services Contract Health Service Delivery Areas (See Appendix G)

[2] Abbreviations: NOS - Not Otherwise Specified; ONS - Other Nervous System

Total Cancer Cases and Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates[1] for the Five Most Commonly Diagnosed Cancers Selected Areas in the NAACCR United States Combined[2], 2010-2014, American Indian/Alaskan Native for CHSDA[3] Counties, Males

	All Sites			Prostate			Lung & Bronchus			Colon & Rectum			Kidney & Renal Pelvis			Liver & IBD[4]		
	Count	*Rate*	*S.E.*[4]	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
NAACCR U.S. Combined	**12,813**	**425.3**	**4.17**	**2,311**	**78.3**	**1.78**	**1,911**	**71.9**	**1.80**	**1,513**	**50.1**	**1.43**	**994**	**30.0**	**1.03**	**687**	**20.1**	**0.84**
Alabama	35	200.2	39.19	8	61.7	27.48	7	33.3	13.73	-	-	-	-	-	-	-	-	-
Alaska	985	507.6	18.08	99	54.1	6.03	178	108.8	8.95	178	91.7	7.60	54	24.5	3.60	34	15.7	3.04
Arizona	1,730	319.9	8.40	312	66.3	4.01	104	22.9	2.40	189	32.6	2.55	216	35.4	2.58	121	20.1	1.96
California	1,194	248.0	7.95	243	50.5	3.51	151	37.0	3.30	102	22.2	2.43	81	15.3	1.88	88	15.6	1.82
CA: Greater California	1,194	248.0	7.95	243	50.5	3.51	151	37.0	3.30	102	22.2	2.43	81	15.3	1.88	88	15.6	1.82
Colorado	31	298.2	61.41	6	75.5	34.46	6	62.7	27.67	-	-	-	-	-	-	-	-	-
Connecticut	55	185.2	28.30	8	28.2	10.79	14	57.5	16.93	9	24.0	8.74	-	-	-	-	-	-
Florida	81	209.9	26.06	17	39.8	10.64	8	19.5	7.48	9	24.5	9.13	6	16.7	7.71	-	-	-
Idaho	91	359.6	42.26	22	71.5	15.89	10	45.6	15.43	12	57.1	19.11	-	-	-	9	30.8	10.66
Indiana	9	151.2	55.42	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Iowa	38	704.9	139.67	7	159.1	72.82	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana	22	272.4	65.29	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Maine	64	550.0	77.91	17	151.2	39.58	9	64.4	22.24	-	-	-	-	-	-	-	-	-
Massachusetts	35	122.7	21.86	12	43.2	13.06	-	-	-	8	26.6	10.04	-	-	-	-	-	-
Michigan	416	512.2	28.01	70	87.0	11.37	88	110.1	12.71	51	65.5	10.26	32	37.0	7.17	16	18.1	5.04
Mississippi	51	401.6	62.62	-	-	-	-	-	-	6	46.1	21.90	-	-	-	9	87.4	31.18
Montana	496	564.5	28.82	100	106.9	11.88	83	115.9	14.19	63	74.4	10.74	43	47.6	8.60	22	20.6	4.63
Nebraska	85	440.5	62.65	12	49.6	15.16	15	116.5	38.02	8	38.7	15.08	8	23.2	8.47	9	33.7	12.08
New York	264	501.7	33.54	64	111.1	14.60	47	92.9	14.76	29	59.0	12.03	9	18.8	6.68	20	29.8	6.93
North Carolina	136	372.4	36.09	27	80.5	18.25	22	60.4	14.29	16	43.2	12.01	9	18.9	6.42	12	32.2	10.03
North Dakota	252	579.3	40.91	48	101.8	15.40	47	126.4	20.33	33	65.2	12.13	24	51.3	11.12	7	15.6	6.84
Oklahoma	3,961	604.2	10.49	727	112.3	4.53	689	113.0	4.66	454	69.7	3.58	293	41.2	2.57	158	20.9	1.78
Oregon	512	384.3	19.10	58	48.9	7.13	101	85.7	9.86	62	47.2	6.53	33	25.0	4.89	31	20.1	3.90
Pennsylvania	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Rhode Island	8	252.5	89.79	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina	74	162.7	22.53	19	34.7	8.64	10	26.8	9.97	6	14.0	6.29	-	-	-	-	-	-
South Dakota	475	543.7	27.96	97	99.6	10.71	78	117.0	14.72	67	70.6	9.54	42	45.4	7.49	25	25.4	5.86

Total Cancer Cases and Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates[1] for the Five Most Commonly Diagnosed Cancers
Selected Areas in the NAACCR United States Combined[2], 2010-2014, American Indian/Alaskan Native for CHSDA[3] Counties, Males

	All Sites			Prostate			Lung & Bronchus			Colon & Rectum			Kidney & Renal Pelvis			Liver & IBD[4]		
	Count	*Rate*	*S.E.*[4]	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
Texas	38	230.1	39.75	10	55.2	18.45	-	-	-	-	-	-	-	-	-	-	-	-
Utah	75	249.7	32.43	18	71.0	18.24	6	18.4	7.80	13	47.6	15.24	-	-	-	-	-	-
Washington	1,077	485.9	17.43	195	85.6	6.90	138	72.4	7.07	116	53.6	5.91	73	28.7	3.70	89	33.6	4.06
WA: Seattle/Puget Sound	733	527.5	23.20	135	93.2	8.91	98	80.4	9.37	76	54.4	7.48	44	30.9	5.27	62	36.3	5.37
Wisconsin	453	688.0	38.39	90	126.4	15.73	76	143.8	19.01	50	78.5	13.41	30	41.2	8.38	16	30.1	9.42
Wyoming	69	519.3	70.65	11	60.7	19.16	10	92.2	31.33	8	47.6	18.98	6	38.2	17.72	-	-	-

1. Rates are per 100,000 population and are age-adjusted by five-year age groups to the 2000 U.S. standard population based on single years of age.
2. See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.
3. CHSDA - Indian Health Services Contract Health Service Delivery Areas (See Appendix G)
4. Abbreviations: IBD - Intrahepatic Bile Duct; S.E. - Standard Error

\- Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for the U.S. combined. Statistics exclude data for KS for all years.

Total Cancer Cases and Average Annual Age-adjusted (1996 Canadian Standard) Cancer Incidence Rates[1] for the Five Most Commonly Diagnosed Cancers Selected Areas in the NAACCR United States Combined[2], 2010-2014, American Indian/Alaskan Native for CHSDA[3] Counties, Males

	All Sites			Prostate			Lung & Bronchus			Colon & Rectum			Kidney & Renal Pelvis			Liver & IBD[4]		
	Count	*Rate*	*S.E.*[4]	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
NAACCR U.S. Combined	**12,813**	**406.0**	**3.85**	**2,311**	**75.6**	**1.67**	**1,911**	**67.8**	**1.64**	**1,513**	**47.6**	**1.31**	**994**	**29.2**	**0.98**	**687**	**19.4**	**0.79**
Alabama	35	196.9	37.10	8	58.2	24.61	7	33.5	13.26	-	-	-	-	-	-	-	-	-
Alaska	985	481.5	16.60	99	51.0	5.50	178	100.8	8.04	178	87.8	7.06	54	24.2	3.50	34	14.8	2.78
Arizona	1,730	305.3	7.78	312	62.7	3.70	104	21.5	2.19	189	31.7	2.42	216	34.9	2.49	121	19.1	1.84
California	1,194	236.5	7.35	243	49.7	3.36	151	34.5	2.97	102	21.0	2.23	81	14.7	1.75	88	15.2	1.73
CA: Greater California	1,194	236.5	7.35	243	49.7	3.36	151	34.5	2.97	102	21.0	2.23	81	14.7	1.75	88	15.2	1.73
Colorado	31	285.2	56.99	6	70.2	31.24	6	62.4	26.68	-	-	-	-	-	-	-	-	-
Connecticut	55	175.9	26.21	8	28.1	10.57	14	54.5	15.66	9	22.8	8.07	-	-	-	-	-	-
Florida	81	198.0	23.91	17	38.3	9.99	8	18.9	7.16	9	23.1	8.17	6	15.3	6.90	-	-	-
Idaho	91	342.6	38.63	22	72.6	15.93	10	43.4	14.35	12	52.6	16.73	-	-	-	9	30.7	10.52
Indiana	9	145.4	51.57	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Iowa	38	667.6	124.67	7	146.4	62.92	-	-	-	-	-	-	-	-	-	-	-	-
Louisiana	22	257.6	59.12	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Maine	64	534.2	72.87	17	149.4	38.42	9	62.4	21.90	-	-	-	-	-	-	-	-	-
Massachusetts	35	116.3	20.37	12	41.8	12.46	-	-	-	8	24.2	8.87	-	-	-	-	-	-
Michigan	416	491.7	25.98	70	84.2	10.76	88	108.1	12.21	51	61.3	9.29	32	35.7	6.76	16	17.3	4.69
Mississippi	51	390.8	59.88	-	-	-	-	-	-	6	42.5	19.57	-	-	-	9	86.6	30.62
Montana	496	542.6	26.63	100	106.9	11.44	83	108.5	12.82	63	70.6	9.85	43	44.0	7.50	22	20.6	4.63
Nebraska	85	417.3	55.07	12	51.4	15.80	15	104.6	31.44	8	37.5	14.22	8	24.0	8.84	9	34.0	12.34
New York	264	479.7	31.18	64	110.9	14.39	47	88.6	13.63	29	55.1	10.93	9	18.0	6.28	20	28.6	6.62
North Carolina	136	357.9	33.50	27	78.1	16.99	22	59.2	13.30	16	40.8	10.92	9	18.2	6.23	12	30.5	9.36
North Dakota	252	557.3	38.33	48	102.1	15.40	47	122.0	19.05	33	63.1	11.62	24	50.3	10.75	7	14.8	6.29
Oklahoma	3,961	575.1	9.65	727	107.3	4.19	689	106.4	4.26	454	66.0	3.27	293	39.9	2.44	158	20.3	1.70
Oregon	512	372.1	17.85	58	46.2	6.54	101	81.1	8.85	62	45.8	6.21	33	23.8	4.49	31	20.1	3.82
Pennsylvania	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Rhode Island	8	261.8	93.30	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina	74	151.6	19.94	19	33.9	8.30	10	24.1	8.76	6	12.6	5.61	-	-	-	-	-	-
South Dakota	475	525.7	26.15	97	100.5	10.72	78	107.9	13.07	67	69.2	9.06	42	45.2	7.41	25	24.3	5.35

Total Cancer Cases and Average Annual Age-adjusted (1996 Canadian Standard) Cancer Incidence Rates[1] for the Five Most Commonly Diagnosed Cancers Selected Areas in the NAACCR United States Combined[2], 2010-2014, American Indian/Alaskan Native for CHSDA[3] Counties, Males

	All Sites			Prostate			Lung & Bronchus			Colon & Rectum			Kidney & Renal Pelvis			Liver & IBD[4]		
	Count	*Rate*	*S.E.*[4]	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
Texas	38	223.1	37.66	10	54.2	17.87	-	-	-	-	-	-	-	-	-	-	-	-
Utah	75	238.7	29.83	18	67.9	17.04	6	19.0	8.06	13	43.1	13.02	-	-	-	-	-	-
Washington	1,077	461.3	15.76	195	83.8	6.50	138	68.1	6.40	116	50.4	5.28	73	28.2	3.56	89	32.3	3.82
WA: Seattle/Puget Sound	733	504.4	21.10	135	92.6	8.61	98	77.1	8.63	76	51.8	6.76	44	30.1	4.98	62	34.9	5.03
Wisconsin	453	657.1	34.92	90	122.8	14.42	76	133.8	16.96	50	73.4	11.93	30	40.7	8.07	16	27.8	8.17
Wyoming	69	490.9	64.82	11	62.3	19.75	10	86.5	28.99	8	45.6	17.56	6	35.8	16.13	-	-	-

1. Rates are per 100,000 and are age-adjusted to the 1996 Canadian population standard.
2. See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.
3. CHSDA - Indian Health Services Contract Health Service Delivery Areas (See Appendix G)
4. Abbreviations: IBD - Intrahepatic Bile Duct; S.E. - Standard Error

\- Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for the U.S. combined. Statistics exclude data for KS for all years.

Total Cancer Cases and Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates[1] for the Five Most Commonly Diagnosed Cancers Selected Areas in the NAACCR United States Combined[2], 2010-2014, American Indian/Alaskan Native for CHSDA[3] Counties, Females

	All Sites			Breast			Lung & Bronchus			Colon & Rectum			Uterus Corpus & Uterus, NOS[4]			Kidney & Renal Pelvis		
	Count	*Rate*	*S.E.*[4]	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
NAACCR U.S. Combined	**14,277**	**388.7**	**3.41**	**3,839**	**100.7**	**1.69**	**1,875**	**55.9**	**1.35**	**1,463**	**41.3**	**1.13**	**896**	**22.4**	**0.77**	**670**	**17.4**	**0.70**
Alabama	28	153.4	30.48	-	-	-	6	27.0	11.42	-	-	-	-	-	-	-	-	-
Alaska	1,182	518.8	15.83	343	144.8	8.14	144	68.8	5.99	184	86.4	6.66	43	17.7	2.82	48	19.3	2.91
Arizona	1,986	280.8	6.54	429	58.7	2.92	119	18.8	1.78	164	24.1	1.95	197	25.0	1.81	129	17.5	1.58
California	1,270	221.5	6.55	350	57.6	3.21	129	25.2	2.33	123	22.4	2.13	97	15.7	1.67	53	8.4	1.22
CA: Greater California	1,269	221.3	6.55	350	57.6	3.21	129	25.2	2.33	123	22.4	2.13	97	15.7	1.67	53	8.4	1.22
Colorado	50	381.4	59.30	12	94.1	29.42	-	-	-	6	40.4	18.23	8	52.7	20.66	-	-	-
Connecticut	55	143.3	20.53	20	48.5	11.46	7	19.7	7.96	-	-	-	-	-	-	-	-	-
Florida	74	147.0	17.60	30	58.0	10.93	11	22.4	6.92	8	14.4	5.10	-	-	-	-	-	-
Idaho	125	446.4	43.44	34	119.5	21.68	10	41.2	14.02	18	80.8	21.04	7	19.2	7.31	6	18.9	7.77
Indiana	8	150.1	58.09	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Iowa	39	498.2	91.48	9	104.1	37.96	-	-	-	6	111.0	52.00	-	-	-	-	-	-
Louisiana	16	163.2	43.40	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Maine	55	487.5	71.14	20	162.4	37.74	10	85.9	29.91	-	-	-	-	-	-	-	-	-
Massachusetts	52	168.5	24.26	13	37.9	10.81	12	45.2	13.41	-	-	-	-	-	-	-	-	-
Michigan	439	467.3	23.71	122	120.3	11.43	101	115.5	12.24	36	35.4	6.26	14	13.1	3.67	23	27.5	6.06
Mississippi	76	534.9	71.45	12	73.3	21.35	-	-	-	9	51.4	17.24	6	53.8	28.37	-	-	-
Montana	558	521.9	23.68	147	131.6	11.58	83	92.6	10.84	66	60.6	7.87	32	30.9	5.80	28	22.9	4.53
Nebraska	119	460.3	48.43	37	121.1	22.31	17	94.3	24.66	13	56.6	18.12	-	-	-	-	-	-
New York	291	468.2	28.07	72	116.4	14.02	43	72.2	11.28	32	50.2	9.05	17	26.5	6.57	11	17.3	5.34
North Carolina	148	366.5	32.24	44	105.0	16.85	19	46.7	11.34	11	29.0	9.42	11	23.9	7.43	7	18.1	7.43
North Dakota	276	516.7	33.36	72	122.8	15.25	52	101.2	14.76	34	65.2	12.02	16	28.1	7.28	15	31.8	9.36
Oklahoma	4,295	537.7	8.48	1,187	145.8	4.36	661	86.5	3.48	433	55.8	2.77	238	28.1	1.87	219	26.4	1.84
Oregon	580	399.6	18.11	147	102.2	9.19	89	71.5	8.24	53	37.4	5.53	46	25.5	3.97	16	10.9	2.91
Pennsylvania	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Rhode Island	14	435.8	124.05	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina	71	128.9	16.48	21	38.5	9.01	9	13.1	4.47	-	-	-	8	14.9	5.83	-	-	-
South Dakota	503	445.7	20.88	152	126.4	10.58	75	77.2	9.35	55	49.6	6.97	25	20.1	4.14	16	13.9	3.71

Total Cancer Cases and Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates[1] for the Five Most Commonly Diagnosed Cancers
Selected Areas in the NAACCR United States Combined[2], 2010-2014, American Indian/Alaskan Native for CHSDA[3] Counties, Females

	All Sites			Breast			Lung & Bronchus			Colon & Rectum			Uterus Corpus & Uterus, NOS[4]			Kidney & Renal Pelvis		
	Count	*Rate*	*S.E.*[4]	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
Texas	28	144.5	28.70	7	36.4	14.45	-	-	-	-	-	-	-	-	-	-	-	-
Utah	100	262.0	27.72	18	40.1	9.64	-	-	-	14	40.0	11.35	10	24.6	8.05	-	-	-
Washington	1,247	465.0	14.32	377	136.3	7.56	160	68.0	5.84	113	46.5	4.75	69	22.1	2.84	39	13.0	2.21
WA: Seattle/Puget Sound	859	505.2	18.91	263	150.2	10.06	100	68.1	7.48	75	48.4	6.05	48	23.4	3.64	23	13.0	2.90
Wisconsin	507	607.2	28.45	124	138.8	12.97	89	119.5	13.31	46	53.5	8.33	30	30.9	5.83	23	26.1	5.65
Wyoming	84	448.4	50.78	28	144.8	28.10	11	60.7	19.03	10	50.4	16.63	-	-	-	-	-	-

1. Rates are per 100,000 population and are age-adjusted by five-year age groups to the 2000 U.S. standard population based on single years of age.
2. See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.
3. CHSDA - Indian Health Services Contract Health Service Delivery Areas (See Appendix G)
4. Abbreviations: NOS - Not Otherwise Specified; S.E. - Standard Error

\- Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for the U.S. combined. Statistics exclude data for KS for all years.

Total Cancer Cases and Average Annual Age-adjusted (1996 Canadian Standard) Cancer Incidence Rates[1] for the Five Most Commonly Diagnosed Cancers Selected Areas in the NAACCR United States Combined[2], 2010-2014, American Indian/Alaskan Native for CHSDA[3] Counties, Females

	All Sites			Breast			Lung & Bronchus			Colon & Rectum			Uterus Corpus & Uterus, NOS[4]			Kidney & Renal Pelvis		
	Count	*Rate*	*S.E.*[4]	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
NAACCR U.S. Combined	**14,277**	**374.0**	**3.23**	**3,839**	**98.0**	**1.63**	**1,875**	**53.1**	**1.26**	**1,463**	**39.2**	**1.06**	**896**	**22.0**	**0.76**	**670**	**16.9**	**0.67**
Alabama	28	152.3	29.97	-	-	-	6	24.8	10.26	-	-	-	-	-	-	-	-	-
Alaska	1,182	501.0	15.11	343	141.6	7.91	144	65.6	5.64	184	82.2	6.26	43	17.4	2.76	48	19.1	2.86
Arizona	1,986	271.2	6.24	429	57.0	2.82	119	17.8	1.67	164	22.9	1.83	197	25.0	1.81	129	17.1	1.54
California	1,270	213.1	6.21	350	57.1	3.16	129	24.2	2.20	123	21.0	1.97	97	15.3	1.61	53	8.2	1.18
CA: Greater California	1,269	212.9	6.21	350	57.1	3.16	129	24.2	2.20	123	21.0	1.97	97	15.3	1.61	53	8.2	1.18
Colorado	50	361.6	54.60	12	91.4	27.71	-	-	-	6	35.9	15.53	8	52.5	19.86	-	-	-
Connecticut	55	138.5	19.68	20	47.9	11.20	7	19.1	7.68	-	-	-	-	-	-	-	-	-
Florida	74	143.9	17.09	30	55.7	10.42	11	22.3	6.81	8	15.5	5.53	-	-	-	-	-	-
Idaho	125	421.9	39.57	34	117.8	20.94	10	37.2	12.38	18	68.0	17.01	7	19.5	7.50	6	18.2	7.55
Indiana	8	142.7	53.84	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Iowa	39	479.6	84.47	9	96.3	33.76	-	-	-	6	103.2	46.80	-	-	-	-	-	-
Louisiana	16	155.9	40.68	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Maine	55	467.2	66.44	20	160.2	36.95	10	79.4	27.11	-	-	-	-	-	-	-	-	-
Massachusetts	52	159.1	22.80	13	37.7	10.84	12	41.7	12.30	-	-	-	-	-	-	-	-	-
Michigan	439	444.8	22.13	122	116.8	10.97	101	107.6	11.17	36	34.6	6.02	14	12.9	3.60	23	25.2	5.44
Mississippi	76	513.7	64.51	12	74.9	21.95	-	-	-	9	52.1	17.46	6	48.7	23.30	-	-	-
Montana	558	500.0	22.15	147	126.8	10.92	83	85.5	9.75	66	59.3	7.59	32	29.5	5.43	28	22.5	4.44
Nebraska	119	440.0	44.93	37	119.1	21.69	17	86.9	22.21	13	51.8	16.05	-	-	-	-	-	-
New York	291	449.4	26.80	72	113.1	13.56	43	67.3	10.43	32	48.7	8.73	17	25.4	6.25	11	16.8	5.17
North Carolina	148	344.3	29.59	44	98.8	15.60	19	45.6	10.83	11	26.5	8.38	11	23.4	7.17	7	16.7	6.73
North Dakota	276	501.2	31.61	72	121.1	14.84	52	98.9	14.23	34	62.8	11.29	16	27.2	7.08	15	29.3	8.17
Oklahoma	4,295	519.2	8.10	1,187	142.2	4.21	661	83.0	3.29	433	53.0	2.60	238	27.7	1.83	219	25.5	1.77
Oregon	580	381.9	16.90	147	96.7	8.52	89	67.6	7.59	53	35.7	5.18	46	25.0	3.84	16	10.4	2.74
Pennsylvania	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Rhode Island	14	394.3	111.60	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
South Carolina	71	120.9	15.08	21	35.9	8.19	9	12.5	4.31	-	-	-	8	13.2	4.99	-	-	-
South Dakota	503	425.6	19.64	152	124.3	10.33	75	72.0	8.59	55	47.3	6.57	25	19.5	3.99	16	13.2	3.43

Total Cancer Cases and Average Annual Age-adjusted (1996 Canadian Standard) Cancer Incidence Rates[1] for the Five Most Commonly Diagnosed Cancers
Selected Areas in the NAACCR United States Combined[2], 2010-2014, American Indian/Alaskan Native for CHSDA[3] Counties, Females

	All Sites			Breast			Lung & Bronchus			Colon & Rectum			Uterus Corpus & Uterus, NOS[4]			Kidney & Renal Pelvis		
	Count	*Rate*	*S.E.*[4]	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
Texas	28	140.6	27.42	7	36.1	14.06	-	-	-	-	-	-	-	-	-	-	-	-
Utah	100	253.8	26.44	18	41.3	10.01	-	-	-	14	38.1	10.68	10	24.3	7.93	-	-	-
Washington	1,247	443.5	13.32	377	131.0	7.10	160	64.2	5.38	113	43.5	4.35	69	21.7	2.75	39	12.7	2.14
WA: Seattle/Puget Sound	859	481.9	17.57	263	143.9	9.39	100	63.5	6.80	75	45.5	5.58	48	23.0	3.52	23	12.6	2.79
Wisconsin	507	584.0	26.92	124	134.4	12.47	89	113.7	12.42	46	50.7	7.75	30	30.3	5.71	23	26.2	5.63
Wyoming	84	445.3	50.29	28	144.0	27.96	11	60.7	19.05	10	46.6	15.22	-	-	-	-	-	-

1. Rates are per 100,000 and are age-adjusted to the 1996 Canadian population standard.
2. See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.
3. CHSDA - Indian Health Services Contract Health Service Delivery Areas (See Appendix G)
4. Abbreviations: NOS - Not Otherwise Specified; S.E. - Standard Error

- Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for the U.S. combined. Statistics exclude data for KS for all years.

Total Cancer Cases and Average Annual Age-Adjusted Cancer Incidence Rates*
Most Frequent Cancer Sites by IHS[1] Region, NAACCR U.S. Combined Selected CHSDA[1] Counties
2010-2014, American Indian/Alaskan Native, Males

	U.S. (CHSDA Counties)					Northern Plains				
Cancer site or type	*Rank*	*Count*	*Rate*	*95% CI*	*S.E.*	*Rank*	*Count*	*Rate*	*95% CI*	*S.E.*
All Sites		12,813	425.3	417.1 - 433.5	4.17		2,293	555.6	530.0 - 582.0	13.21
Prostate	1	2,311	78.3	74.8 - 81.8	1.78	1	438	99.1	89.1 - 109.9	5.23
Lung & Bronchus	2	1,911	71.9	68.4 - 75.5	1.80	2	402	117.0	104.6 - 130.3	6.52
Colon & Rectum	3	1,513	50.1	47.4 - 53.0	1.43	3	284	66.8	58.3 - 76.2	4.50
Kidney & Renal Pelvis	4	994	30.0	28.0 - 32.1	1.03	4	190	43.0	36.4 - 50.4	3.51
Liver and Intrahepatic Bile Duct	5	687	20.1	18.5 - 21.8	0.84	5	102	22.9	18.1 - 28.5	2.60
Oral Cavity & Pharynx	6	554	16.3	14.8 - 17.8	0.76	6	100	20.1	15.9 - 25.0	2.25
Non-Hodgkin Lymphoma	7	502	16.7	15.1 - 18.4	0.84	7	75	18.9	14.3 - 24.4	2.50
Urinary Bladder	8	489	19.4	17.6 - 21.4	0.98	8	72	21.5	16.1 - 27.9	2.95
Leukemia	9	417	12.7	11.4 - 14.2	0.71	9	65	14.5	10.7 - 19.1	2.08
Pancreas	10	358	12.0	10.7 - 13.5	0.70	10	56	13.6	9.9 - 18.2	2.04
Stomach	11	329	11.6	10.2 - 13.0	0.71	11	48	12.0	8.4 - 16.4	1.98
Melanoma of the Skin	12	251	8.6	7.4 - 9.8	0.61	15	34	7.7	4.8 - 11.4	1.61
Testis	13	243	4.9	4.3 - 5.6	0.32	14	42	6.2	4.4 - 8.5	0.98
Esophagus	14	236	7.8	6.7 - 9.0	0.56	13	43	11.5	7.9 - 15.9	1.98
Brain & Other Nervous System	15	214	7.8	6.6 - 9.0	0.59	16	32	8.7	5.7 - 12.6	1.69
Myeloma	16	211	5.4	4.6 - 6.3	0.42	17	32	5.9	3.7 - 8.8	1.24
Thyroid	17	164	4.4	3.7 - 5.1	0.37	18	30	5.2	3.5 - 7.6	0.97
Larynx	18	159	5.1	4.2 - 6.0	0.44	12	44	11.2	7.7 - 15.5	1.94

	Alaska					Southern Plains				
Cancer site or type	*Rank*	*Count*	*Rate*	*95% CI*	*S.E.*	*Rank*	*Count*	*Rate*	*95% CI*	*S.E.*
All Sites		985	507.6	472.8 - 544.1	18.08		3,999	594.2	574.3 - 614.6	10.25
Prostate	3	99	54.1	42.9 - 67.0	6.03	1	737	110.7	102.1 - 119.6	4.43
Lung & Bronchus	1	178	108.8	91.9 - 127.5	8.95	2	692	110.6	101.9 - 119.8	4.54
Colon & Rectum	2	178	91.7	77.4 - 107.7	7.60	3	459	68.7	62.0 - 75.9	3.50
Kidney & Renal Pelvis	5	54	24.5	18.0 - 32.6	3.60	4	295	40.3	35.5 - 45.5	2.51
Liver and Intrahepatic Bile Duct	8	34	15.7	10.3 - 22.8	3.04	8	160	20.8	17.5 - 24.6	1.77
Oral Cavity & Pharynx	6	49	21.9	15.6 - 29.8	3.52	6	171	22.7	19.2 - 26.6	1.86
Non-Hodgkin Lymphoma	10	31	14.6	9.4 - 21.4	2.93	7	167	25.4	21.3 - 30.0	2.19
Urinary Bladder	9	34	22.3	14.8 - 31.8	4.21	5	185	31.7	26.9 - 37.0	2.54
Leukemia	7	39	15.8	10.4 - 22.7	3.01	9	127	17.8	14.6 - 21.6	1.75
Pancreas	11	29	16.1	10.4 - 23.6	3.25	11	105	15.8	12.7 - 19.3	1.67
Stomach	4	59	27.8	20.5 - 36.7	4.00	14	64	10.3	7.7 - 13.3	1.39
Melanoma of the Skin	20	6	2.7	0.9 - 6.2	1.20	10	117	19.1	15.5 - 23.2	1.93
Testis	12	25	8.2	5.2 - 12.4	1.70	16	47	4.8	3.5 - 6.5	0.72
Esophagus	13	20	9.8	5.6 - 15.8	2.48	13	66	9.3	7.1 - 12.0	1.22
Brain & Other Nervous System	17	9	5.9	2.3 - 11.7	2.27	15	58	9.6	7.1 - 12.7	1.39
Myeloma	15	14	5.9	2.9 - 10.5	1.79	12	69	8.4	6.4 - 10.9	1.12
Thyroid	14	17	7.8	4.3 - 12.9	2.07	18	42	5.2	3.7 - 7.2	0.85
Larynx	18	9	4.3	1.8 - 8.5	1.57	17	47	6.2	4.4 - 8.4	0.96

* Rates are per 100,000 population and are age-adjusted by five-year age groups to the 2000 U.S. standard population based on single years of age. The rates are based on cases and populations included in the combined incidence for the United States excluding KS.

[1] Abbreviations: IHS - Indian Health Services; CHSDA - Contract Health Service Delivery Area

~ Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for All Sites combined.

Total Cancer Cases and Average Annual Age-Adjusted Cancer Incidence Rates*
Most Frequent Cancer Sites by IHS[1] Region, NAACCR U.S. Combined Selected CHSDA[1] Counties
2010-2014, American Indian/Alaskan Native, Males

	Pacific Coast					East				
Cancer site or type	*Rank*	*Count*	*Rate*	*95% CI*	*S.E.*	*Rank*	*Count*	*Rate*	*95% CI*	*S.E.*
All Sites		2,874	331.4	317.9 - 345.3	6.95		826	284.2	263.0 - 306.5	11.01
Prostate	1	518	60.1	54.5 - 66.0	2.90	1	183	62.2	52.7 - 72.8	5.04
Lung & Bronchus	2	400	53.5	47.8 - 59.6	2.99	2	123	45.0	36.6 - 54.5	4.49
Colon & Rectum	3	292	34.7	30.4 - 39.4	2.28	3	96	35.4	27.8 - 44.1	4.06
Kidney & Renal Pelvis	5	191	20.2	17.2 - 23.7	1.62	6	38	12.1	8.2 - 17.0	2.15
Liver and Intrahepatic Bile Duct	4	217	21.4	18.4 - 24.7	1.59	4	50	15.5	11.2 - 20.8	2.36
Oral Cavity & Pharynx	8	134	13.5	11.1 - 16.2	1.29	5	40	13.5	9.3 - 18.9	2.37
Non-Hodgkin Lymphoma	6	137	16.0	13.1 - 19.3	1.56	11	25	8.6	5.4 - 13.0	1.87
Urinary Bladder	7	135	19.4	15.9 - 23.3	1.87	8	30	12.1	7.8 - 17.8	2.47
Leukemia	9	90	9.5	7.4 - 12.0	1.15	7	32	10.9	7.0 - 16.0	2.20
Pancreas	10	82	9.7	7.5 - 12.3	1.20	9	28	8.4	5.5 - 12.4	1.68
Stomach	15	51	6.3	4.5 - 8.4	0.99	10	27	10.0	6.3 - 15.1	2.15
Melanoma of the Skin	14	55	6.2	4.5 - 8.3	0.96	13	16	5.4	2.8 - 9.3	1.58
Testis	11	62	4.3	3.3 - 5.6	0.56	12	17	3.6	2.1 - 6.0	0.88
Esophagus	12	57	6.0	4.3 - 8.0	0.90	14	15	5.1	2.7 - 8.7	1.44
Brain & Other Nervous System	16	44	5.1	3.6 - 7.1	0.86	16	11	3.9	1.7 - 7.5	1.37
Myeloma	13	56	4.6	3.4 - 6.2	0.68	~	~	~	~	~
Thyroid	18	34	3.1	2.1 - 4.6	0.61	17	9	2.8	1.2 - 5.6	1.01
Larynx	17	38	4.6	3.1 - 6.4	0.82	15	12	4.1	1.9 - 7.4	1.32

	Southwest				
Cancer site or type	*Rank*	*Count*	*Rate*	*95% CI*	*S.E.*
All Sites		1,836	315.9	300.3 - 332.1	8.07
Prostate	1	336	66.6	59.2 - 74.6	3.88
Lung & Bronchus	5	116	23.4	19.1 - 28.3	2.33
Colon & Rectum	3	204	33.0	28.3 - 38.2	2.50
Kidney & Renal Pelvis	2	226	34.6	29.9 - 39.7	2.46
Liver and Intrahepatic Bile Duct	4	124	19.2	15.7 - 23.2	1.86
Oral Cavity & Pharynx	9	60	9.9	7.3 - 12.9	1.39
Non-Hodgkin Lymphoma	7	67	10.8	8.2 - 14.0	1.45
Urinary Bladder	15	33	6.5	4.3 - 9.3	1.24
Leukemia	8	64	9.8	7.2 - 12.9	1.41
Pancreas	11	58	10.1	7.5 - 13.3	1.45
Stomach	6	80	15.7	12.2 - 19.8	1.91
Melanoma of the Skin	19	23	4.1	2.5 - 6.2	0.92
Testis	12	50	4.7	3.5 - 6.3	0.69
Esophagus	14	35	6.7	4.5 - 9.4	1.22
Brain & Other Nervous System	10	60	11.1	8.2 - 14.5	1.56
Myeloma	13	37	4.8	3.1 - 6.9	0.91
Thyroid	16	32	4.2	2.8 - 6.0	0.78
Larynx	21	9	1.4	0.6 - 2.8	0.50

* Rates are per 100,000 population and are age-adjusted by five-year age groups to the 2000 U.S. standard population based on single years of age. The rates are based on cases and populations included in the combined incidence for the United States excluding KS.

[1] Abbreviations: IHS - Indian Health Services; CHSDA - Contract Health Service Delivery Area

~ Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for All Sites combined.

Total Cancer Cases and Average Annual Age-Adjusted Cancer Incidence Rates*
Most Frequent Cancer Sites by IHS[1] Region, NAACCR U.S. Combined Selected CHSDA[1] Counties
2010-2014, American Indian/Alaskan Native, Females

	U.S. (CHSDA Counties)					Northern Plains				
Cancer site or type	*Rank*	*Count*	*Rate*	*95% CI*	*S.E.*	*Rank*	*Count*	*Rate*	*95% CI*	*S.E.*
All Sites		14,277	388.7	382.0 - 395.4	3.41		2,533	494.4	474.2 - 515.3	10.45
Breast	1	3,839	100.7	97.4 - 104.0	1.69	1	691	125.5	115.9 - 135.7	5.00
Lung & Bronchus	2	1,875	55.9	53.2 - 58.6	1.35	2	435	96.6	87.2 - 106.6	4.91
Colon & Rectum	3	1,463	41.3	39.1 - 43.6	1.13	3	268	52.2	45.8 - 59.2	3.38
Corpus & Uterus, NOS	4	896	22.4	20.9 - 23.9	0.77	4	125	22.2	18.3 - 26.6	2.08
Kidney & Renal Pelvis	5	670	17.4	16.1 - 18.8	0.70	5	117	22.8	18.6 - 27.6	2.25
Thyroid	6	618	14.5	13.4 - 15.7	0.60	6	91	13.9	11.2 - 17.2	1.49
Non-Hodgkin Lymphoma	7	473	13.5	12.3 - 14.8	0.65	8	71	14.4	11.0 - 18.3	1.82
Ovary	8	399	10.6	9.6 - 11.8	0.56	10	59	12.0	8.9 - 15.6	1.66
Cervix Uteri	9	374	9.1	8.2 - 10.1	0.49	7	72	12.8	9.9 - 16.2	1.56
Pancreas	10	359	10.8	9.6 - 12.0	0.60	11	53	11.4	8.4 - 15.1	1.67
Leukemia	11	353	9.0	8.1 - 10.1	0.51	9	61	11.7	8.7 - 15.3	1.64
Liver and Intrahepatic Bile Duct	12	320	8.8	7.8 - 9.9	0.52	12	49	9.4	6.8 - 12.7	1.45
Oral Cavity & Pharynx	13	228	6.1	5.3 - 7.0	0.43	13	46	9.1	6.6 - 12.3	1.42
Stomach	14	222	6.5	5.7 - 7.5	0.46	16	29	6.2	4.0 - 8.9	1.20
Melanoma of the Skin	15	210	5.6	4.8 - 6.4	0.40	21	17	2.9	1.7 - 4.7	0.74
Urinary Bladder	16	194	5.9	5.1 - 6.8	0.45	15	36	7.7	5.2 - 10.8	1.38
Myeloma	17	173	5.0	4.3 - 5.9	0.40	18	22	4.9	2.9 - 7.5	1.11
Brain & Other Nervous System	18	154	3.6	3.0 - 4.3	0.31	14	37	6.3	4.3 - 8.9	1.12

	Alaska					Southern Plains				
Cancer site or type	*Rank*	*Count*	*Rate*	*95% CI*	*S.E.*	*Rank*	*Count*	*Rate*	*95% CI*	*S.E.*
All Sites		1,182	518.8	488.3 - 550.7	15.83		4,323	528.1	512.0 - 544.6	8.30
Breast	1	343	144.8	129.3 - 161.6	8.14	1	1,194	143.1	134.9 - 151.7	4.26
Lung & Bronchus	3	144	68.8	57.5 - 81.4	5.99	2	661	84.5	78.0 - 91.4	3.40
Colon & Rectum	2	184	86.4	73.9 - 100.4	6.66	3	437	54.9	49.7 - 60.5	2.72
Corpus & Uterus, NOS	6	43	17.7	12.6 - 24.1	2.82	4	240	27.7	24.2 - 31.5	1.84
Kidney & Renal Pelvis	5	48	19.3	14.0 - 25.8	2.91	5	221	25.9	22.5 - 29.7	1.80
Thyroid	4	53	21.6	15.9 - 28.5	3.11	6	173	19.2	16.3 - 22.3	1.49
Non-Hodgkin Lymphoma	8	31	13.4	8.9 - 19.2	2.52	7	148	19.1	16.0 - 22.5	1.62
Ovary	11	23	9.4	5.8 - 14.3	2.05	9	111	13.2	10.8 - 16.0	1.30
Cervix Uteri	13	18	7.9	4.5 - 12.6	1.95	10	109	12.8	10.4 - 15.4	1.25
Pancreas	9	30	15.3	10.1 - 22.0	2.93	8	116	15.0	12.3 - 18.0	1.44
Leukemia	12	23	8.2	5.0 - 12.6	1.81	11	107	12.3	9.9 - 14.9	1.24
Liver and Intrahepatic Bile Duct	17	12	5.4	2.7 - 9.5	1.63	12	75	8.9	6.9 - 11.2	1.07
Oral Cavity & Pharynx	10	29	11.4	7.5 - 16.6	2.21	15	49	6.0	4.4 - 8.0	0.89
Stomach	7	38	17.5	12.2 - 24.2	2.96	16	46	5.9	4.2 - 7.9	0.89
Melanoma of the Skin	15	13	5.5	2.9 - 9.6	1.61	14	65	8.3	6.3 - 10.6	1.05
Urinary Bladder	14	14	6.9	3.6 - 11.7	1.94	13	67	8.8	6.8 - 11.3	1.12
Myeloma	21	6	3.0	1.1 - 6.5	1.27	18	42	5.0	3.6 - 6.9	0.80
Brain & Other Nervous System	16	13	4.6	2.4 - 8.0	1.32	17	44	4.6	3.3 - 6.3	0.73

* Rates are per 100,000 population and are age-adjusted by five-year age groups to the 2000 U.S. standard population based on single years of age. The rates are based on cases and populations included in the combined incidence for the United States excluding KS.

[1] Abbreviations: IHS - Indian Health Services; CHSDA - Contract Health Service Delivery Area

~ Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for All Sites combined.

Total Cancer Cases and Average Annual Age-Adjusted Cancer Incidence Rates*
Most Frequent Cancer Sites by IHS[1] Region, NAACCR U.S. Combined Selected CHSDA[1] Counties
2010-2014, American Indian/Alaskan Native, Females

	Pacific Coast					East				
Cancer site or type	*Rank*	*Count*	*Rate*	*95% CI*	*S.E.*	*Rank*	*Count*	*Rate*	*95% CI*	*S.E.*
All Sites		3,222	315.7	304.2 - 327.5	5.93		881	264.1	246.2 - 282.9	9.29
Breast	1	908	85.8	80.0 - 91.9	3.01	1	244	70.7	61.8 - 80.5	4.71
Lung & Bronchus	2	388	42.9	38.5 - 47.7	2.32	2	123	38.6	31.8 - 46.3	3.63
Colon & Rectum	3	307	32.1	28.4 - 36.1	1.95	3	83	25.0	19.7 - 31.2	2.86
Corpus & Uterus, NOS	4	219	19.0	16.5 - 21.8	1.35	4	54	15.9	11.7 - 20.9	2.26
Kidney & Renal Pelvis	7	114	10.3	8.4 - 12.5	1.02	5	35	10.3	7.1 - 14.5	1.83
Thyroid	5	137	11.6	9.7 - 13.9	1.04	8	23	6.4	3.9 - 9.7	1.40
Non-Hodgkin Lymphoma	6	119	12.5	10.2 - 15.1	1.22	6	25	7.7	4.9 - 11.4	1.60
Ovary	10	81	7.8	6.1 - 9.8	0.93	14	19	5.3	3.1 - 8.4	1.27
Cervix Uteri	8	107	8.6	7.0 - 10.4	0.85	7	24	6.2	4.0 - 9.3	1.29
Pancreas	12	77	8.2	6.4 - 10.4	1.00	13	20	6.3	3.7 - 9.8	1.48
Leukemia	9	83	8.0	6.2 - 10.0	0.93	11	21	5.7	3.4 - 8.8	1.28
Liver and Intrahepatic Bile Duct	11	80	7.6	5.9 - 9.6	0.92	10	22	7.6	4.6 - 11.5	1.68
Oral Cavity & Pharynx	14	59	5.6	4.2 - 7.4	0.78	15	17	5.6	3.2 - 9.1	1.41
Stomach	17	36	3.5	2.4 - 4.9	0.62	9	23	7.7	4.8 - 11.6	1.67
Melanoma of the Skin	13	68	6.1	4.7 - 7.8	0.78	12	21	6.2	3.7 - 9.5	1.40
Urinary Bladder	15	47	5.6	4.1 - 7.6	0.88	17	12	3.7	1.9 - 6.6	1.12
Myeloma	16	43	4.6	3.2 - 6.3	0.76	16	14	5.0	2.7 - 8.3	1.36
Brain & Other Nervous System	18	30	2.8	1.8 - 4.0	0.54	18	9	2.3	1.0 - 4.5	0.80

	Southwest				
Cancer site or type	*Rank*	*Count*	*Rate*	*95% CI*	*S.E.*
All Sites		2,136	281.4	269.2 - 294.1	6.33
Breast	1	459	58.3	53.0 - 64.1	2.81
Lung & Bronchus	6	124	18.1	14.9 - 21.7	1.68
Colon & Rectum	3	184	25.2	21.5 - 29.2	1.93
Corpus & Uterus, NOS	2	215	25.4	22.0 - 29.1	1.77
Kidney & Renal Pelvis	5	135	17.0	14.2 - 20.2	1.51
Thyroid	4	141	15.9	13.3 - 18.8	1.36
Non-Hodgkin Lymphoma	9	79	10.8	8.5 - 13.6	1.27
Ovary	7	106	13.5	11.0 - 16.4	1.36
Cervix Uteri	14	44	5.2	3.7 - 7.0	0.80
Pancreas	10	63	9.6	7.3 - 12.4	1.27
Leukemia	11	58	7.1	5.3 - 9.2	0.98
Liver and Intrahepatic Bile Duct	8	82	11.7	9.2 - 14.6	1.34
Oral Cavity & Pharynx	15	28	3.5	2.3 - 5.2	0.69
Stomach	12	50	7.8	5.7 - 10.3	1.13
Melanoma of the Skin	17	26	3.5	2.2 - 5.2	0.72
Urinary Bladder	20	18	2.6	1.5 - 4.1	0.63
Myeloma	13	46	6.5	4.7 - 8.7	0.99
Brain & Other Nervous System	19	21	2.3	1.4 - 3.6	0.53

* Rates are per 100,000 population and are age-adjusted by five-year age groups to the 2000 U.S. standard population based on single years of age. The rates are based on cases and populations included in the combined incidence for the United States excluding KS.

[1] Abbreviations: IHS - Indian Health Services; CHSDA - Contract Health Service Delivery Area

~ Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for All Sites combined.

NAACCR United States Combined Incidence 2010-2014, Hispanic/Latino, All Races Combined

Total Cancer Cases and Average Annual Age-adjusted Cancer Incidence Rates*
NAACCR United States Combined, 2010-2014, Hispanic/Latino, All Races

	MALES (HISPANIC/LATINO)					FEMALES (HISPANIC/LATINA)				
SITES	*Count*	*Adj. Rate 2000 U.S.*	*S.E.*	*Adj. Rate 1996 Cdn*	*S.E.*	*Count*	*Adj. Rate 2000 US.*	*S.E.*	*Adj. Rate 1996 Cdn*	*S.E.*
All Sites (invasive & in situ bladder)	267,627	386.3	0.82	368.6	0.76	292,872	329.6	0.63	317.7	0.60
Oral Cavity & Pharynx	7,836	10.5	0.13	10.1	0.12	3,649	4.2	0.07	4.0	0.07
Lip	291	0.4	0.03	0.4	0.03	127	0.2	0.01	0.2	0.01
Tongue	2,129	2.8	0.07	2.7	0.06	1,068	1.2	0.04	1.2	0.04
Major salivary gland	788	1.1	0.04	1.0	0.04	751	0.8	0.03	0.8	0.03
Floor of mouth	368	0.5	0.03	0.5	0.03	118	0.1	0.01	0.1	0.01
Gum & other mouth	961	1.4	0.05	1.3	0.05	738	0.9	0.03	0.9	0.03
Nasopharynx	456	0.5	0.03	0.5	0.03	210	0.2	0.02	0.2	0.01
Oropharynx	385	0.5	0.03	0.5	0.03	107	0.1	0.01	0.1	0.01
Hypopharynx	577	0.8	0.04	0.8	0.04	91	0.1	0.01	0.1	0.01
Digestive System	67,716	99.1	0.41	94.0	0.38	53,219	65.5	0.29	61.8	0.27
Esophagus	3,152	4.8	0.09	4.6	0.09	842	1.1	0.04	1.0	0.04
Stomach	8,479	12.9	0.15	12.1	0.14	6,462	7.7	0.10	7.4	0.09
Small intestine	1,355	1.9	0.06	1.8	0.05	1,338	1.6	0.04	1.5	0.04
Colon excl. rectum	18,575	28.2	0.23	26.6	0.21	17,390	21.5	0.17	20.1	0.16
Rectum & rectosigmoid	10,037	13.7	0.15	13.1	0.14	6,815	7.8	0.10	7.5	0.09
Anus, anal canal & anorectum	775	1.0	0.04	0.9	0.04	1,351	1.6	0.04	1.5	0.04
Liver	13,779	18.5	0.17	17.8	0.16	5,158	6.5	0.09	6.1	0.09
Gallbladder	733	1.2	0.05	1.1	0.04	2,033	2.6	0.06	2.4	0.06
Pancreas	7,607	12.0	0.15	11.3	0.14	7,924	10.3	0.12	9.6	0.11
Respiratory System	27,555	46.5	0.30	43.6	0.28	20,420	26.4	0.19	24.9	0.18
Larynx	3,211	4.8	0.09	4.6	0.09	543	0.6	0.03	0.6	0.03
Lung & bronchus	23,424	40.6	0.28	37.9	0.26	19,375	25.2	0.19	23.8	0.17
Bones & Joints	1,130	1.0	0.03	1.0	0.03	944	0.8	0.03	0.8	0.03
Soft Tissue (incl. heart)	3,136	3.6	0.07	3.5	0.07	2,574	2.5	0.05	2.4	0.05
Skin (excl. basal & sq.)	4,157	6.0	0.10	5.6	0.09	4,724	5.1	0.08	4.9	0.07
Melanomas of the skin	3,394	4.9	0.09	4.6	0.08	3,985	4.2	0.07	4.1	0.07
Breast (excl. *in situ*)	526	0.8	0.04	0.7	0.03	83,873	91.9	0.33	89.5	0.32
Female Genital System	-	-	-	-	-	42,811	45.5	0.23	44.7	0.22
Cervix uteri	-	-	-	-	-	10,009	9.7	0.10	9.8	0.10
Uterus Corpus	-	-	-	-	-	19,691	21.2	0.16	20.9	0.15
Uterus, NOS[1]	-	-	-	-	-	888	1.0	0.04	1.0	0.03
Ovary	-	-	-	-	-	9,452	10.3	0.11	9.9	0.11
Male Genital System	71,169	103.2	0.42	100.6	0.40	-	-	-	-	-
Prostate	63,511	97.1	0.41	94.3	0.39	-	-	-	-	-
Testis	6,559	4.5	0.06	4.8	0.06	-	-	-	-	-
Penis	966	1.4	0.05	1.3	0.04	-	-	-	-	-
Urinary System	26,975	41.0	0.27	38.7	0.25	14,672	17.3	0.15	16.6	0.14
Bladder (incl. *in situ*)	10,984	19.5	0.20	17.9	0.18	3,791	5.0	0.08	4.6	0.08
Kidney & renal pelvis	15,570	20.8	0.18	20.1	0.17	10,632	12.0	0.12	11.6	0.12
Ureter	260	0.5	0.03	0.4	0.03	149	0.2	0.02	0.2	0.02
Eye & Orbit	547	0.6	0.03	0.6	0.03	532	0.5	0.02	0.5	0.02
Brain & Other Nervous System	5,571	5.9	0.09	5.8	0.09	4,677	4.5	0.07	4.4	0.07
Brain	5,206	5.6	0.09	5.4	0.08	4,247	4.1	0.07	4.0	0.06
Endocrine System	5,685	6.0	0.09	6.0	0.09	22,133	20.7	0.14	20.9	0.14
Thyroid gland	4,987	5.4	0.08	5.3	0.08	21,544	20.2	0.14	20.4	0.14
Lymphomas	17,533	22.6	0.19	21.6	0.18	15,119	17.1	0.15	16.5	0.14
Hodgkin Lymphoma	2,981	2.8	0.06	2.8	0.06	2,343	2.1	0.05	2.1	0.04
Non-Hodgkin Lymphoma	14,552	19.8	0.18	18.8	0.17	12,776	15.1	0.14	14.4	0.13
Multiple Myeloma	5,092	7.9	0.12	7.4	0.11	4,262	5.3	0.08	5.1	0.08
Leukemias	11,319	13.1	0.14	12.4	0.13	9,004	9.1	0.10	8.7	0.10
Acute lymphocytic	3,575	2.5	0.05	2.5	0.04	2,762	2.0	0.04	2.0	0.04
Chronic lymphocytic	1,649	2.8	0.07	2.6	0.07	1,242	1.6	0.05	1.5	0.04
Acute myeloid	3,273	4.2	0.08	4.0	0.08	2,917	3.1	0.06	3.0	0.06
Chronic myeloid	1,577	1.9	0.06	1.9	0.05	1,120	1.2	0.04	1.2	0.04
Other leukemias	1,245	1.6	0.05	1.5	0.05	963	1.1	0.04	1.0	0.03
Mesothelioma	714	1.2	0.05	1.2	0.05	280	0.3	0.02	0.3	0.02
Kaposi Sarcoma	1,043	1.1	0.04	1.1	0.04	133	0.2	0.02	0.2	0.01
Ill-defined & Unknown	9,923	16.1	0.18	14.8	0.16	9,846	12.6	0.13	11.6	0.12
Selected tumors (non-invasive)										
Brain & ONS[1] (*benign/borderline*)	5,831	7.1	0.11	6.8	0.10	12,308	13.6	0.13	13.1	0.12
Breast (*in situ*)	58	0.1	0.01	0.1	0.01	20,446	21.9	0.16	21.5	0.15

* Rates are per 100,000 population and are age-adjusted by five-year age groups to the 2000 U.S. standard population based on single years of age and to the 1996 Canadian population standard.

- Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for All Sites combined.

[1] Abbreviations: NOS - Not Otherwise Specified; ONS - Other Nervous System

N.B. In areas with small Latino populations, methods to indirectly identify Latinos (like NHIAv2) can overestimate the ethnicity-specific counts of cancer cases. Also, even small errors in Latino population estimates can affect the magnitude of the cancer rates. Data Item 190 used in place of NHIA for MA and RI, see technical notes.

Total Cancer Cases and Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates[1] for Commonly Diagnosed Cancers
Selected Areas in the NAACCR United States Combined[2], 2010-2014, Hispanic/Latino, All Races, Males
{Page 1 of 4}

	All Sites			Prostate			Colon & Rectum			Lung & Bronchus			Kidney & Renal Pelvis			Liver & IBD[3]		
	Count	*Rate*	*S.E.*[3]	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
NAACCR U.S. Combined	**267,627**	**386.3**	**0.82**	**63,511**	**97.1**	**0.41**	**28,612**	**41.9**	**0.27**	**23,424**	**40.6**	**0.28**	**15,570**	**20.8**	**0.18**	**14,558**	**19.7**	**0.18**
Alabama	492	222.9	12.32	101	55.8	6.18	45	22.1	3.93	38	24.2	4.45	31	13.1	2.85	18	7.9	2.21
Alaska	148	296.3	30.65	25	57.4	14.56	19	27.2	6.67	18	50.4	14.35	-	-	-	9	18.4	7.40
Arizona	8,866	341.4	4.00	1,654	66.2	1.75	1,086	43.5	1.44	780	38.1	1.45	648	23.5	1.00	579	20.0	0.90
Arkansas	1,238	735.5	25.53	192	122.6	10.11	133	78.1	8.31	171	115.9	10.41	77	45.3	6.16	37	15.3	2.79
California	72,388	369.0	1.50	17,520	95.9	0.77	7,706	39.2	0.49	4,989	31.8	0.48	4,358	20.7	0.34	4,164	20.0	0.33
CA: Greater California	38,029	371.7	2.10	8,802	93.4	1.06	4,102	39.5	0.67	2,751	33.7	0.68	2,452	22.4	0.49	2,273	20.7	0.47
CA: Greater Bay Area	9,879	404.4	4.47	2,644	115.3	2.39	932	38.9	1.39	678	35.5	1.44	536	19.3	0.91	570	21.2	0.96
CA: Los Angeles	24,461	352.2	2.46	6,066	92.5	1.26	2,671	39.0	0.82	1,559	27.6	0.74	1,370	18.5	0.54	1,319	18.5	0.55
Colorado	5,588	383.6	5.67	1,206	86.5	2.67	645	43.5	1.88	486	41.2	2.00	385	24.9	1.41	378	23.7	1.31
Connecticut	3,222	516.5	10.44	722	120.5	4.95	317	51.6	3.35	329	62.7	3.82	135	19.5	1.94	197	27.7	2.20
Delaware	356	424.5	25.80	103	139.1	14.98	28	31.1	6.60	28	37.6	7.85	12	12.4	4.15	20	26.5	6.45
District of Columbia	269	318.0	23.10	58	78.4	11.59	32	31.7	6.81	18	32.1	8.33	12	16.7	5.87	9	12.5	4.64
Florida	38,208	425.3	2.22	9,859	110.0	1.12	4,052	45.7	0.73	4,584	53.8	0.81	1,656	17.8	0.45	1,217	13.1	0.38
Georgia	2,671	336.9	8.44	578	89.8	4.33	233	30.8	2.60	225	39.3	3.15	124	14.8	1.65	84	10.3	1.35
Hawaii	649	444.1	19.59	108	75.1	7.91	67	44.5	6.10	93	79.8	8.98	35	20.0	3.73	36	20.8	3.80
Idaho	690	363.1	16.29	143	81.7	7.65	69	37.7	5.11	65	43.8	6.13	47	22.3	3.81	41	19.7	3.52
Illinois	9,593	375.1	4.42	2,403	99.8	2.25	944	37.7	1.41	713	37.2	1.53	611	21.5	1.00	473	18.7	0.97
Indiana	1,429	322.0	9.94	291	75.3	4.86	157	36.6	3.35	137	38.0	3.58	96	19.4	2.32	61	13.6	2.00
Iowa	459	288.2	16.51	81	62.1	7.86	46	29.6	5.12	34	33.8	6.37	21	17.0	4.28	28	18.6	4.23
Kansas	1,267	391.3	12.67	290	100.4	6.41	136	45.9	4.42	100	40.3	4.41	91	27.3	3.31	49	13.7	2.20
Kentucky	386	248.5	15.09	76	55.8	7.08	42	26.9	4.86	54	42.0	6.46	25	13.3	3.17	20	14.1	3.59
Louisiana	1,168	370.9	11.78	309	107.5	6.46	102	31.5	3.37	125	44.1	4.20	66	21.3	2.83	48	14.2	2.23
Maine	78	296.0	36.28	16	63.9	16.97	-	-	-	7	34.1	13.38	8	32.4	12.01	-	-	-
Maryland	1,672	310.6	9.28	460	96.5	5.11	160	29.2	2.83	100	26.7	2.99	78	16.2	2.22	61	10.9	1.66
Massachusetts	3,225	388.7	7.86	953	120.0	4.29	262	34.9	2.51	263	38.6	2.65	166	17.4	1.55	195	22.3	1.79
Michigan	2,084	377.8	9.19	481	89.8	4.42	223	42.5	3.13	207	45.3	3.40	136	23.8	2.26	112	19.2	1.99
MI: Metropolitan Detroit	1,035	512.7	17.37	250	127.8	8.61	114	57.6	5.86	109	62.3	6.37	65	32.5	4.36	43	21.1	3.50
Mississippi	220	215.6	17.19	55	64.5	9.77	25	24.9	5.77	22	27.8	6.50	12	11.0	3.85	10	10.2	3.74

Total Cancer Cases and Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates[1] for Commonly Diagnosed Cancers
Selected Areas in the NAACCR United States Combined[2], 2010-2014, Hispanic/Latino, All Races, Males
{Page 2 of 4}

	All Sites			Prostate			Colon & Rectum			Lung & Bronchus			Kidney & Renal Pelvis			Liver & IBD[3]		
	Count	*Rate*	*S.E.*[3]	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
Missouri	758	282.6	11.53	140	59.6	5.39	78	29.9	3.77	87	41.8	4.79	47	14.7	2.35	39	13.2	2.27
Montana	184	467.6	38.21	41	96.4	16.11	23	53.6	12.42	21	73.8	17.10	10	29.0	9.79	6	10.9	4.90
Nebraska	647	357.4	17.23	138	96.3	9.23	55	30.3	4.99	49	35.8	5.72	52	27.4	4.58	27	14.1	3.27
New Hampshire	178	376.5	32.19	50	109.3	16.88	14	27.9	8.80	16	48.5	12.96	7	17.8	7.45	-	-	-
New Jersey	10,677	444.2	4.74	2,989	129.4	2.55	1,056	44.3	1.50	932	45.0	1.59	501	17.8	0.87	404	16.1	0.88
New York	26,324	468.9	3.11	7,906	141.3	1.68	2,496	46.0	0.99	2,319	46.8	1.03	1,073	17.3	0.56	1,368	22.8	0.66
North Carolina	2,246	300.1	8.26	486	81.2	4.27	186	25.8	2.48	168	31.2	2.90	131	15.7	1.82	107	14.1	1.70
North Dakota	39	283.5	55.40	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Ohio	1,334	295.3	9.01	322	76.0	4.57	143	30.3	2.82	133	37.1	3.45	66	12.0	1.65	81	15.2	1.85
Oklahoma	1,152	320.6	11.47	202	70.5	5.61	129	34.3	3.59	112	43.7	4.63	81	19.9	2.71	55	13.4	2.05
Oregon	1,697	368.0	10.91	326	80.1	5.04	175	40.5	3.72	129	36.7	3.74	113	23.9	2.77	107	20.6	2.33
Pennsylvania	3,895	421.8	7.76	1,006	116.3	4.05	388	43.3	2.52	376	49.5	2.84	170	16.5	1.45	250	23.2	1.64
Rhode Island	573	372.9	18.33	150	103.1	9.50	62	39.5	6.02	56	43.5	6.51	22	10.9	2.50	25	14.1	3.30
South Carolina	739	296.0	13.51	147	67.6	6.22	72	25.6	3.79	71	34.3	4.80	39	18.2	3.45	29	11.9	2.65
South Dakota	74	255.3	33.44	13	53.6	15.61	8	25.5	9.60	6	33.9	13.96	-	-	-	-	-	-
Tennessee	742	238.6	11.00	142	59.8	5.68	57	19.6	3.16	73	30.0	4.06	37	12.1	2.46	33	10.1	2.16
Texas	52,057	377.5	1.82	10,000	78.2	0.84	6,400	46.5	0.63	4,663	41.0	0.64	3,914	26.1	0.45	3,769	25.6	0.45
Utah	1,449	391.8	12.24	354	108.0	6.41	141	36.4	3.59	92	33.6	3.92	91	25.0	3.05	71	19.4	2.64
Vermont	31	192.9	37.12	6	44.9	19.22	-	-	-	-	-	-	-	-	-	-	-	-
Virginia	1,952	284.7	8.13	452	73.6	4.06	182	29.1	2.70	168	33.5	3.04	107	14.1	1.71	72	10.5	1.54
Washington	2,861	367.8	8.41	594	92.7	4.33	253	31.0	2.29	227	40.8	3.07	156	18.8	1.82	157	19.0	1.80
WA: Seattle/Puget Sound	1,734	431.4	12.99	358	105.6	6.44	147	36.4	3.67	127	44.3	4.62	102	21.3	2.64	99	22.2	2.72
West Virginia	81	195.5	22.95	18	49.4	12.04	11	25.5	8.19	9	24.3	8.46	-	-	-	6	12.4	5.05
Wisconsin	1,339	397.7	12.97	292	92.2	6.07	110	30.8	3.47	104	39.8	4.40	85	22.7	2.89	88	25.6	3.19
Wyoming	232	270.0	18.90	48	57.3	8.72	33	38.3	7.07	22	32.9	7.36	18	18.9	4.72	10	10.8	3.60

1. Rates are per 100,000 population and are age-adjusted by five-year age groups to the 2000 U.S. standard population based on single years of age.
2. See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.
3. Abbreviations: IBD - Intrahepatic Bile Duct; S.E. - Standard Error

\- Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for the U.S. combined.

N.B. In areas with small Latino populations, methods to indirectly identify Latinos (like NHIAv2) can overestimate the ethnicity-specific counts of cancer cases. Also, even small errors in Latino population estimates can affect the magnitude of the cancer rates. Data Item 190 used in place of NHIA for MA and RI, see technical notes.

Total Cancer Cases and Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates[1] for Commonly Diagnosed Cancers
Selected Areas in the NAACCR United States Combined[2], 2010-2014, Hispanic/Latino, All Races, Males
{Page 3 of 4}

	Non-Hodgkin Lymphoma			Leukemia			Bladder			Stomach			Oral Cavity & Pharynx			Pancreas		
	Count	*Rate*	*S.E.*[3]	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
NAACCR U.S. Combined	**14,552**	**19.8**	**0.18**	**11,319**	**13.1**	**0.14**	**10,984**	**19.5**	**0.20**	**8,479**	**12.9**	**0.15**	**7,836**	**10.5**	**0.13**	**7,607**	**12.0**	**0.15**
Alabama	25	11.6	2.92	35	12.9	3.29	13	7.0	2.30	19	6.2	1.62	16	6.4	1.78	17	9.3	2.68
Alaska	15	27.4	8.32	-	-	-	6	15.9	9.16	6	6.3	2.66	-	-	-	-	-	-
Arizona	421	15.7	0.85	413	11.8	0.68	332	16.0	0.94	283	11.4	0.74	209	8.0	0.60	288	11.9	0.76
Arkansas	58	35.0	5.93	56	23.9	4.35	72	59.8	8.02	20	15.7	4.12	57	29.7	4.80	40	26.3	5.10
California	4,219	20.6	0.35	3,206	12.5	0.26	2,838	18.0	0.36	2,470	13.1	0.29	1,968	9.3	0.23	2,067	11.6	0.27
CA: Greater California	2,183	20.6	0.49	1,723	12.6	0.36	1,485	18.1	0.50	1,191	12.4	0.39	1,045	9.5	0.32	1,088	11.8	0.39
CA: Greater Bay Area	594	23.4	1.06	428	14.6	0.81	431	21.7	1.11	279	11.9	0.78	263	9.8	0.66	278	12.5	0.81
CA: Los Angeles	1,442	19.7	0.57	1,055	11.7	0.41	922	16.4	0.58	1,000	14.7	0.50	659	8.8	0.37	701	11.0	0.45
Colorado	253	17.1	1.20	211	11.3	0.91	221	18.0	1.31	175	12.7	1.05	176	11.5	0.95	153	11.4	1.00
Connecticut	180	26.1	2.32	148	21.3	2.15	186	39.0	3.17	116	20.0	2.15	88	13.5	1.64	77	14.4	1.83
Delaware	13	13.4	4.09	23	15.4	4.29	15	23.5	6.74	6	9.4	4.20	13	12.5	3.98	12	15.2	4.95
District of Columbia	15	17.8	5.56	11	8.0	2.93	6	9.5	4.48	11	16.2	5.76	12	10.2	3.07	8	9.8	3.77
Florida	1,903	20.4	0.48	1,336	14.3	0.40	2,100	25.4	0.56	947	10.7	0.36	1,381	14.6	0.40	1,118	12.9	0.39
Georgia	195	18.9	1.87	174	15.0	1.70	89	18.2	2.22	77	10.5	1.60	92	10.9	1.47	58	8.1	1.29
Hawaii	33	23.8	4.69	30	20.4	4.36	36	29.1	5.45	17	13.1	3.44	26	15.4	3.22	28	19.0	3.90
Idaho	37	16.4	3.24	33	10.1	2.28	26	18.0	4.01	11	6.9	2.33	16	8.6	2.42	18	11.9	3.11
Illinois	504	18.5	0.97	467	13.5	0.78	363	18.9	1.10	327	13.6	0.86	259	9.3	0.66	298	13.5	0.88
Indiana	71	15.7	2.20	77	9.1	1.34	62	17.3	2.47	58	14.1	2.17	33	6.3	1.26	48	12.9	2.08
Iowa	28	15.2	3.67	25	8.9	2.43	20	17.4	4.32	12	6.9	2.45	15	8.8	2.71	15	11.3	3.27
Kansas	63	17.2	2.56	69	11.8	1.86	57	22.7	3.31	31	10.5	2.11	38	11.1	2.10	22	8.0	1.86
Kentucky	26	13.7	3.59	20	8.6	2.37	16	13.7	3.88	-	-	-	11	7.5	2.67	9	7.9	3.03
Louisiana	65	17.3	2.37	45	11.8	2.03	55	20.6	2.92	31	10.9	2.09	50	13.7	2.11	18	6.2	1.55
Maine	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	6	21.1	8.98
Maryland	96	15.3	1.96	97	11.6	1.67	66	18.6	2.59	68	11.1	1.72	50	7.9	1.33	31	6.5	1.38
Massachusetts	175	19.1	1.69	109	11.1	1.32	112	16.4	1.78	121	16.0	1.67	129	15.6	1.56	73	9.7	1.32
Michigan	121	22.1	2.25	89	12.4	1.58	97	21.0	2.32	56	12.1	1.77	60	9.5	1.33	57	11.1	1.60
MI: Metropolitan Detroit	59	29.0	4.12	42	14.4	2.52	58	33.2	4.66	28	16.8	3.37	40	16.1	2.71	33	17.6	3.29
Mississippi	11	9.6	3.41	7	5.2	2.70	-	-	-	-	-	-	-	-	-	-	-	-

Total Cancer Cases and Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates[1] for Commonly Diagnosed Cancers
Selected Areas in the NAACCR United States Combined[2], 2010-2014, Hispanic/Latino, All Races, Males
{Page 4 of 4}

	Non-Hodgkin Lymphoma			Leukemia			Bladder			Stomach			Oral Cavity & Pharynx			Pancreas		
	Count	*Rate*	*S.E.*[3]	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
Missouri	52	18.6	3.01	41	12.9	2.49	28	12.5	2.57	30	11.4	2.31	27	9.0	1.90	16	7.0	1.88
Montana	-	-	-	8	15.3	6.48	9	25.2	9.04	-	-	-	6	10.3	4.23	6	13.8	6.12
Nebraska	42	25.1	4.79	46	17.0	3.71	25	18.3	4.15	16	7.6	2.23	23	9.6	2.25	15	8.3	2.65
New Hampshire	10	16.4	6.24	10	21.0	7.68	9	23.7	8.62	-	-	-	-	-	-	-	-	-
New Jersey	631	24.5	1.09	435	15.9	0.87	499	25.9	1.26	336	14.8	0.89	288	11.4	0.74	285	13.1	0.84
New York	1,416	23.5	0.68	875	14.6	0.54	1,106	23.9	0.77	892	16.8	0.61	806	13.3	0.50	699	13.1	0.53
North Carolina	143	15.6	1.80	144	10.7	1.37	70	14.2	2.01	91	12.8	1.73	65	7.8	1.23	56	10.2	1.64
North Dakota	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Ohio	66	14.6	2.02	58	8.8	1.41	66	19.3	2.52	33	8.2	1.59	39	8.3	1.44	33	7.9	1.51
Oklahoma	59	15.5	2.49	67	11.7	2.04	48	19.5	3.16	21	6.0	1.51	43	10.2	1.85	45	16.3	2.78
Oregon	107	20.1	2.47	90	12.6	1.78	68	20.0	2.80	49	11.2	1.93	53	9.8	1.52	60	14.4	2.22
Pennsylvania	186	18.1	1.55	158	12.8	1.30	174	24.9	2.10	104	12.6	1.41	124	12.0	1.22	101	12.5	1.41
Rhode Island	28	18.2	4.05	26	13.2	3.41	21	21.1	5.13	23	16.8	4.05	22	12.9	3.28	12	9.3	2.86
South Carolina	37	13.8	2.95	44	8.5	1.65	23	15.6	3.65	36	14.6	3.11	21	7.1	1.97	21	10.7	2.71
South Dakota	11	36.9	13.37	-	-	-	-	-	-	-	-	-	6	23.7	10.37	-	-	-
Tennessee	39	10.6	2.28	60	9.6	1.70	27	12.5	2.82	23	10.0	2.51	29	8.1	1.77	17	6.2	1.87
Texas	2,682	19.0	0.41	2,168	12.8	0.32	1,695	15.0	0.39	1,666	13.0	0.35	1,368	9.2	0.27	1,541	12.1	0.33
Utah	89	24.0	3.09	57	8.9	1.63	55	17.7	2.72	44	13.4	2.41	32	9.0	1.85	45	16.0	2.68
Vermont	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Virginia	128	16.4	1.88	102	10.7	1.51	72	12.9	1.82	96	14.6	1.90	42	6.1	1.18	52	8.6	1.44
Washington	182	22.2	2.06	151	12.2	1.41	114	18.6	2.01	89	10.7	1.39	86	9.9	1.29	67	10.6	1.49
WA: Seattle/Puget Sound	96	21.9	2.85	83	13.0	2.01	84	28.1	3.66	38	8.8	1.72	61	14.2	2.19	44	13.5	2.38
West Virginia	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Wisconsin	88	24.8	3.28	79	17.3	2.65	51	20.1	3.21	37	16.2	3.02	34	9.3	1.85	50	16.1	2.62
Wyoming	9	11.0	3.88	9	11.0	3.94	16	21.7	5.70	-	-	-	10	9.2	2.91	6	5.8	2.37

1. Rates are per 100,000 population and are age-adjusted by five-year age groups to the 2000 U.S. standard population based on single years of age.
2. See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.
3. Abbreviations: S.E. - Standard Error

\- Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for the U.S. combined.

N.B. In areas with small Latino populations, methods to indirectly identify Latinos (like NHIAv2) can overestimate the ethnicity-specific counts of cancer cases. Also, even small errors in Latino population estimates can affect the magnitude of the cancer rates. Data Item 190 used in place of NHIA for MA and RI, see technical notes.

Total Cancer Cases and Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates[1] for Commonly Diagnosed Cancers
Selected Areas in the NAACCR United States Combined[2], 2010-2014, Hispanic/Latina, All Races, Females
{Page 1 of 4}

	All Sites			Breast			Colon & Rectum			Thyroid			Uterus Corpus & Uterus NOS[3]			Lung & Bronchus		
	Count	*Rate*	*S.E.*[3]	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
NAACCR U.S. Combined	**292,872**	**329.6**	**0.63**	**83,873**	**91.9**	**0.33**	**24,205**	**29.3**	**0.19**	**21,544**	**20.2**	**0.14**	**20,579**	**22.2**	**0.16**	**19,375**	**25.2**	**0.19**
Alabama	456	205.9	10.80	143	64.3	5.83	44	25.1	4.08	38	12.1	2.25	23	10.8	2.49	33	20.8	3.79
Alaska	167	261.0	22.91	54	80.4	12.10	22	32.3	7.21	13	13.6	3.88	8	9.8	3.51	9	15.4	5.58
Arizona	10,581	324.1	3.31	2,976	88.9	1.69	847	28.2	1.01	901	22.3	0.77	712	20.3	0.79	736	26.9	1.02
Arkansas	1,329	692.5	22.43	297	148.1	10.10	116	70.7	7.64	82	24.4	3.13	80	39.7	5.10	130	90.6	8.76
California	80,046	315.2	1.16	23,487	89.2	0.60	6,633	27.9	0.36	5,433	17.7	0.25	5,684	21.4	0.29	4,552	21.4	0.33
CA: Greater California	41,428	319.6	1.65	12,442	92.2	0.86	3,400	28.3	0.51	2,921	18.2	0.35	2,687	19.9	0.40	2,508	23.5	0.49
CA: Greater Bay Area	10,520	334.9	3.39	3,065	94.7	1.77	848	28.4	1.01	648	16.4	0.67	746	23.2	0.88	627	23.6	0.96
CA: Los Angeles	28,084	302.5	1.86	7,976	83.0	0.95	2,384	27.1	0.57	1,864	17.3	0.41	2,251	22.9	0.50	1,416	17.7	0.48
Colorado	6,376	359.6	4.73	1,910	104.2	2.47	516	32.1	1.47	512	23.0	1.05	373	19.5	1.04	528	35.6	1.60
Connecticut	3,813	447.3	7.83	1,142	127.5	4.03	294	36.7	2.31	344	31.6	1.79	247	28.3	1.93	279	41.3	2.61
Delaware	379	376.5	21.64	102	100.0	11.00	21	26.9	6.29	40	28.2	4.73	28	26.6	5.54	30	38.4	7.39
District of Columbia	304	278.9	16.99	80	70.8	8.22	20	22.8	5.43	29	21.4	4.09	18	15.6	3.76	19	23.7	5.60
Florida	39,523	342.5	1.73	11,343	98.1	0.92	3,823	33.2	0.54	2,716	23.5	0.45	2,778	23.7	0.45	3,137	27.3	0.49
Georgia	3,180	314.0	6.50	1,012	93.2	3.35	209	26.2	2.06	259	17.4	1.24	197	19.2	1.53	173	24.0	1.98
Hawaii	822	447.5	16.57	282	151.6	9.51	71	42.3	5.29	40	14.9	2.44	45	22.2	3.41	82	51.8	6.00
Idaho	745	328.3	13.57	200	89.1	7.05	55	26.4	3.97	68	21.5	2.86	57	22.5	3.20	43	23.9	3.93
Illinois	10,557	341.8	3.61	2,928	89.5	1.76	795	29.0	1.11	873	21.1	0.76	843	24.9	0.91	663	26.2	1.08
Indiana	1,602	315.5	8.69	438	83.7	4.33	137	30.2	2.79	114	16.6	1.71	117	21.0	2.08	123	30.9	2.95
Iowa	520	268.2	13.54	142	67.1	6.13	36	25.1	4.73	38	13.4	2.41	40	16.9	2.90	35	28.1	5.03
Kansas	1,323	329.8	9.92	347	84.2	4.85	80	22.9	2.74	138	27.5	2.54	87	20.9	2.37	91	28.9	3.19
Kentucky	308	202.7	12.82	82	54.2	6.51	25	18.1	3.98	21	8.6	2.05	15	10.4	2.93	33	29.2	5.27
Louisiana	1,156	326.5	9.79	323	91.5	5.15	108	31.5	3.09	71	17.2	2.09	74	20.2	2.38	105	32.3	3.19
Maine	68	231.0	29.00	19	56.6	13.26	-	-	-	-	-	-	6	17.6	7.24	11	44.1	13.62
Maryland	2,253	301.7	6.98	703	91.8	3.75	157	22.7	1.95	200	19.8	1.50	150	19.1	1.68	137	24.6	2.22
Massachusetts	3,604	320.3	5.78	1,022	87.2	2.90	270	27.0	1.77	390	26.9	1.43	237	20.5	1.40	258	28.1	1.86
Michigan	2,026	302.6	7.11	547	80.0	3.58	150	24.2	2.07	143	16.7	1.49	173	24.5	1.94	178	31.2	2.43
MI: Metropolitan Detroit	1,223	484.5	14.54	315	125.3	7.37	96	39.6	4.19	75	23.0	2.78	106	39.1	3.95	123	56.2	5.25
Mississippi	197	188.5	14.49	46	41.4	6.44	14	18.5	5.14	15	11.7	3.23	10	8.8	2.93	19	22.1	5.24

Total Cancer Cases and Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates[1] for Commonly Diagnosed Cancers
Selected Areas in the NAACCR United States Combined[2], 2010-2014, Hispanic/Latina, All Races, Females
{Page 2 of 4}

	All Sites			Breast			Colon & Rectum			Thyroid			Uterus Corpus & Uterus NOS[3]			Lung & Bronchus		
	Count	*Rate*	*S.E.*[3]	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
Missouri	888	287.7	10.34	250	78.9	5.28	59	22.3	3.04	66	17.4	2.28	64	18.0	2.35	86	34.7	3.88
Montana	214	437.9	31.34	66	134.3	17.13	16	33.0	8.57	12	20.5	6.41	9	20.4	7.03	25	56.8	11.85
Nebraska	704	357.7	15.35	188	93.4	7.67	55	32.5	4.90	51	16.2	2.56	42	20.4	3.51	61	39.6	5.48
New Hampshire	225	350.5	25.54	60	92.2	12.88	20	31.7	7.70	28	33.4	6.62	12	18.5	5.51	18	36.5	9.16
New Jersey	11,643	365.4	3.52	3,299	98.7	1.77	1,010	34.6	1.13	953	24.9	0.83	828	25.1	0.90	802	28.4	1.04
New York	28,042	362.1	2.20	7,996	101.3	1.15	2,262	30.6	0.65	2,259	26.4	0.56	2,088	26.1	0.58	2,126	29.3	0.65
North Carolina	2,756	310.0	7.04	755	82.5	3.49	172	22.5	2.00	251	17.1	1.24	184	20.8	1.76	177	29.3	2.44
North Dakota	37	200.5	36.43	12	58.5	18.17	-	-	-	-	-	-	-	-	-	-	-	-
Ohio	1,284	234.6	6.91	357	64.3	3.55	115	22.4	2.19	91	12.7	1.41	91	16.4	1.78	115	24.1	2.34
Oklahoma	1,433	358.1	10.78	395	96.9	5.43	115	32.9	3.43	82	14.8	1.90	77	17.2	2.14	104	34.6	3.67
Oregon	1,976	366.6	9.44	532	97.0	4.72	140	28.2	2.67	156	18.2	1.64	147	24.6	2.26	170	44.3	3.66
Pennsylvania	3,948	336.4	5.80	1,040	86.5	2.87	298	27.4	1.71	351	22.7	1.28	297	25.5	1.57	291	29.9	1.86
Rhode Island	601	286.5	12.78	191	85.2	6.68	43	22.5	3.73	43	15.6	2.52	42	20.6	3.47	36	21.2	3.77
South Carolina	819	285.8	11.35	266	89.6	6.10	76	28.3	3.57	62	15.1	2.07	38	13.4	2.41	60	29.8	4.17
South Dakota	60	182.5	25.19	16	46.8	12.24	11	36.1	11.52	-	-	-	-	-	-	-	-	-
Tennessee	844	241.0	9.60	228	66.5	4.91	55	18.9	2.81	64	11.8	1.68	55	15.1	2.27	61	25.8	3.54
Texas	56,426	321.8	1.41	15,884	88.0	0.72	4,624	27.8	0.42	3,623	17.5	0.30	3,982	21.3	0.35	3,197	21.4	0.39
Utah	1,744	371.0	10.03	479	101.8	5.14	124	30.8	3.05	207	31.0	2.37	125	25.7	2.52	110	33.0	3.37
Vermont	40	235.6	39.66	15	88.1	23.96	-	-	-	-	-	-	7	45.7	18.62	-	-	-
Virginia	2,805	293.2	6.30	823	80.6	3.13	209	25.1	1.95	314	21.9	1.34	161	16.2	1.41	166	23.0	1.96
Washington	3,306	350.3	7.01	930	93.6	3.41	227	27.9	2.07	290	20.4	1.37	195	20.4	1.62	220	32.0	2.36
WA: Seattle/Puget Sound	2,255	464.9	11.30	697	137.3	5.84	143	34.4	3.24	195	25.6	2.10	133	27.9	2.70	172	47.9	4.00
West Virginia	61	142.5	18.61	14	34.5	9.40	-	-	-	-	-	-	-	-	-	9	21.7	7.30
Wisconsin	1,407	334.0	10.05	376	90.2	5.14	101	28.4	3.11	131	20.1	2.00	111	25.1	2.57	104	32.4	3.41
Wyoming	274	307.8	19.12	76	85.5	10.00	16	19.7	5.03	15	13.7	3.68	14	14.8	4.03	22	28.4	6.18

1. Rates are per 100,000 population and are age-adjusted by five-year age groups to the 2000 U.S. standard population based on single years of age.
2. See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.
3. Abbreviations: NOS - Not Otherwise Specified; S.E. - Standard Error

\- Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for the U.S. combined.

N.B. In areas with small Latino populations, methods to indirectly identify Latinos (like NHIAv2) can overestimate the ethnicity-specific counts of cancer cases. Also, even small errors in Latino population estimates can affect the magnitude of the cancer rates. Data Item 190 used in place of NHIA for MA and RI, see technical notes.

Total Cancer Cases and Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates[1] for Commonly Diagnosed Cancers
Selected Areas in the NAACCR United States Combined[2], 2010-2014, Hispanic/Latina, All Races, Females
{Page 3 of 4}

	Non-Hodgkin Lymphoma			Kidney & Renal Pelvis			Cervix			Ovary			Leukemia			Pancreas		
	Count	*Rate*	*S.E.*[3]	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
NAACCR U.S. Combined	**12,776**	**15.1**	**0.14**	**10,632**	**12.0**	**0.12**	**10,009**	**9.7**	**0.10**	**9,452**	**10.3**	**0.11**	**9,004**	**9.1**	**0.10**	**7,924**	**10.3**	**0.12**
Alabama	20	9.7	2.42	12	5.6	1.77	14	3.8	1.15	21	8.2	2.00	15	5.1	1.73	6	3.4	1.48
Alaska	7	13.6	5.37	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Arizona	421	13.7	0.70	504	15.3	0.72	334	8.6	0.49	341	9.8	0.56	340	9.3	0.55	309	11.4	0.67
Arkansas	43	19.9	3.50	46	24.5	4.29	69	23.8	3.26	77	40.1	5.26	57	25.7	4.40	26	18.9	4.13
California	3,649	15.2	0.26	2,906	11.7	0.23	2,644	9.0	0.18	2,865	10.7	0.21	2,496	8.6	0.18	2,199	10.2	0.22
CA: Greater California	1,857	15.2	0.37	1,563	12.4	0.33	1,350	8.8	0.25	1,397	10.3	0.29	1,284	8.3	0.25	1,126	10.4	0.32
CA: Greater Bay Area	496	16.6	0.77	385	12.4	0.65	320	8.8	0.51	345	10.4	0.58	340	9.7	0.56	321	11.9	0.68
CA: Los Angeles	1,295	14.7	0.42	957	10.5	0.35	974	9.3	0.31	1,123	11.5	0.35	871	8.5	0.30	752	9.2	0.35
Colorado	226	13.4	0.93	281	16.0	1.00	165	7.8	0.63	216	12.0	0.86	172	8.1	0.68	163	10.7	0.87
Connecticut	187	23.2	1.83	102	11.4	1.21	123	11.8	1.13	93	10.2	1.15	113	12.5	1.30	101	14.0	1.50
Delaware	15	17.2	4.76	10	8.3	2.80	13	8.4	2.44	10	8.5	2.90	13	9.3	3.39	15	16.8	4.78
District of Columbia	15	13.7	3.71	9	9.7	3.34	15	10.1	2.69	-	-	-	7	4.4	1.77	8	9.9	3.65
Florida	1,717	15.0	0.36	1,078	9.4	0.29	1,058	9.2	0.28	1,243	10.7	0.31	1,131	9.9	0.29	1,159	10.1	0.30
Georgia	120	13.2	1.40	88	9.0	1.09	142	9.6	0.90	94	9.2	1.12	123	9.7	1.12	64	8.5	1.17
Hawaii	31	17.8	3.41	19	11.2	2.73	17	7.7	1.94	16	9.1	2.39	17	7.6	2.11	31	19.6	3.73
Idaho	27	14.0	3.06	32	13.6	2.63	31	9.8	1.82	22	9.2	2.25	27	9.5	2.23	24	14.0	3.08
Illinois	464	16.4	0.82	412	13.2	0.70	389	10.3	0.56	304	9.3	0.59	343	9.3	0.58	265	11.2	0.72
Indiana	67	14.9	1.97	64	12.8	1.74	54	8.3	1.24	56	10.2	1.49	63	10.6	1.61	37	9.2	1.61
Iowa	17	12.8	3.44	23	11.4	2.71	27	9.4	1.89	20	10.1	2.66	19	8.1	2.44	-	-	-
Kansas	47	13.4	2.11	59	13.9	1.95	61	12.8	1.78	39	10.0	1.75	45	9.6	1.67	28	8.2	1.64
Kentucky	15	8.6	2.47	8	3.8	1.42	12	5.5	1.70	9	5.2	2.04	12	4.9	1.60	7	5.3	2.21
Louisiana	54	15.6	2.18	47	12.9	1.91	41	9.9	1.58	27	7.7	1.52	29	7.5	1.46	41	12.4	1.95
Maine	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Maryland	87	12.2	1.46	68	9.3	1.24	92	8.7	0.96	77	9.5	1.20	71	8.5	1.18	57	9.2	1.31
Massachusetts	176	16.5	1.35	103	8.8	0.94	99	8.3	0.92	78	6.5	0.80	102	8.2	0.91	79	8.0	0.96
Michigan	90	13.3	1.50	83	12.4	1.44	79	9.7	1.13	65	10.0	1.32	55	7.9	1.15	67	11.8	1.51
MI: Metropolitan Detroit	62	23.0	3.07	52	20.0	2.94	38	12.2	2.06	37	15.5	2.67	35	14.0	2.48	36	16.9	2.92
Mississippi	6	6.5	2.92	8	6.8	2.59	11	8.4	2.76	-	-	-	12	7.7	2.50	-	-	-

Total Cancer Cases and Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates[1] for Commonly Diagnosed Cancers
Selected Areas in the NAACCR United States Combined[2], 2010-2014, Hispanic/Latina, All Races, Females
{Page 4 of 4}

	Non-Hodgkin Lymphoma			Kidney & Renal Pelvis			Cervix			Ovary			Leukemia			Pancreas		
	Count	*Rate*	*S.E.*[3]	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
Missouri	36	12.7	2.28	34	11.6	2.13	35	9.9	1.79	23	6.5	1.45	28	6.6	1.43	19	7.4	1.77
Montana	6	15.9	6.67	6	13.6	5.75	-	-	-	-	-	-	-	-	-	10	24.5	8.00
Nebraska	38	21.8	3.88	24	12.1	2.85	26	8.7	1.90	20	8.3	2.16	21	8.0	2.06	14	10.8	3.04
New Hampshire	8	13.8	5.48	-	-	-	6	7.1	2.91	6	9.3	3.89	-	-	-	6	10.8	4.80
New Jersey	524	17.1	0.78	319	9.9	0.57	378	10.6	0.57	322	10.0	0.58	348	10.7	0.60	315	11.5	0.67
New York	1,268	16.6	0.47	734	9.5	0.36	822	10.0	0.35	738	9.3	0.35	798	10.1	0.37	769	10.8	0.40
North Carolina	107	13.7	1.54	98	10.9	1.27	135	10.6	1.10	87	8.4	1.08	113	9.5	1.15	64	8.5	1.23
North Dakota	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Ohio	57	11.1	1.54	46	8.1	1.25	32	4.9	0.89	35	6.3	1.12	39	5.8	1.04	41	9.3	1.49
Oklahoma	60	17.9	2.61	45	9.9	1.61	72	11.7	1.48	62	14.8	2.13	53	8.9	1.56	42	16.4	2.68
Oregon	80	15.8	2.04	73	14.6	1.91	54	7.4	1.14	54	10.3	1.62	63	8.7	1.41	54	13.9	2.04
Pennsylvania	155	13.9	1.21	129	10.3	0.98	170	11.6	0.95	110	9.9	1.02	105	7.0	0.78	78	8.2	0.99
Rhode Island	30	14.4	2.89	20	10.9	2.64	19	7.5	1.80	9	4.6	1.68	14	6.0	1.76	16	8.1	2.26
South Carolina	33	13.4	2.73	30	12.7	2.60	28	6.4	1.29	23	7.5	1.79	30	7.5	1.69	14	6.4	1.87
South Dakota	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee	33	9.0	1.89	25	6.6	1.57	46	9.9	1.62	22	6.1	1.49	43	10.0	1.96	8	3.0	1.17
Texas	2,427	14.8	0.31	2,770	15.9	0.31	2,314	11.3	0.24	1,928	10.7	0.25	1,735	8.9	0.23	1,566	10.3	0.27
Utah	70	14.9	2.01	63	12.6	1.75	52	7.4	1.08	52	10.7	1.66	58	9.0	1.53	46	13.3	2.12
Vermont	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Virginia	116	13.7	1.44	72	8.0	1.06	123	10.7	1.10	105	11.1	1.24	84	7.0	0.91	59	8.0	1.14
Washington	143	16.9	1.61	107	11.1	1.21	124	9.6	0.98	98	10.1	1.17	130	9.8	1.07	67	10.0	1.34
WA: Seattle/Puget Sound	90	18.5	2.24	59	10.3	1.51	66	9.5	1.37	63	12.5	1.85	75	12.1	1.72	40	10.7	1.87
West Virginia	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Wisconsin	66	16.0	2.22	51	12.1	1.88	58	9.9	1.42	52	11.2	1.75	52	11.0	1.90	23	6.5	1.50
Wyoming	8	9.3	3.35	12	13.0	3.84	10	10.8	3.55	8	7.8	2.76	10	9.1	3.05	10	12.0	3.91

1. Rates are per 100,000 population and are age-adjusted by five-year age groups to the 2000 U.S. standard population based on single years of age.
2. See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.
3. Abbreviations: S.E. - Standard Error

\- Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for the U.S. combined.

N.B. In areas with small Latino populations, methods to indirectly identify Latinos (like NHIAv2) can overestimate the ethnicity-specific counts of cancer cases. Also, even small errors in Latino population estimates can affect the magnitude of the cancer rates. Data Item 190 used in place of NHIA for MA and RI, see technical notes.

NAACCR United States Combined Incidence 2010-2014, The Non-Hispanic White Population

UNITED STATES

III-67 Combined Incidence – Non-Hispanic White

Total Cancer Cases and Average Annual Age-adjusted Cancer Incidence Rates*
NAACCR United States Combined, 2010-2014, Non-Hispanic White

	MALES (NON-HISPANIC WHITE)					FEMALES (NON-HISPANIC WHITE)				
SITES	*Count*	*Adj. Rate 2000 U.S.*	*S.E.*	*Adj. Rate 1996 Cdn*	*S.E.*	*Count*	*Adj. Rate 2000 U.S.*	*S.E.*	*Adj. Rate 1996 Cdn*	*S.E.*
All Sites (invasive & in situ bladder)	3,019,830	510.8	0.30	488.2	0.29	2,897,505	436.0	0.27	421.5	0.26
Oral Cavity & Pharynx	115,165	18.8	0.06	18.2	0.05	45,991	6.8	0.03	6.5	0.03
Lip	6,300	1.1	0.01	1.0	0.01	2,329	0.3	0.01	0.3	0.01
Tongue	37,168	6.0	0.03	5.8	0.03	14,105	2.1	0.02	2.0	0.02
Major salivary gland	10,056	1.8	0.02	1.7	0.02	6,300	1.0	0.01	1.0	0.01
Floor of mouth	5,456	0.9	0.01	0.8	0.01	2,516	0.4	0.01	0.3	0.01
Gum & other mouth	11,550	1.9	0.02	1.8	0.02	9,405	1.3	0.01	1.3	0.01
Nasopharynx	3,388	0.6	0.01	0.6	0.01	1,338	0.2	0.01	0.2	0.01
Oropharynx	5,497	0.9	0.01	0.8	0.01	1,725	0.2	0.01	0.2	0.01
Hypopharynx	6,648	1.1	0.01	1.0	0.01	1,691	0.2	0.01	0.2	0.01
Digestive System	555,641	93.9	0.13	88.9	0.12	442,724	62.8	0.10	59.1	0.09
Esophagus	52,340	8.6	0.04	8.2	0.04	13,113	1.8	0.02	1.7	0.02
Stomach	46,073	7.8	0.04	7.4	0.04	24,856	3.5	0.02	3.3	0.02
Small intestine	15,572	2.6	0.02	2.5	0.02	13,518	2.0	0.02	1.9	0.02
Colon excl. rectum	179,193	31.0	0.07	29.1	0.07	182,534	25.7	0.06	24.0	0.06
Rectum & rectosigmoid	83,812	14.2	0.05	13.6	0.05	59,135	8.8	0.04	8.4	0.04
Anus, anal canal & anorectum	8,898	1.5	0.02	1.4	0.02	16,427	2.4	0.02	2.3	0.02
Liver	56,422	8.9	0.04	8.5	0.04	18,163	2.6	0.02	2.4	0.02
Gallbladder	4,031	0.7	0.01	0.6	0.01	8,174	1.1	0.01	1.0	0.01
Pancreas	85,017	14.3	0.05	13.5	0.05	78,507	10.8	0.04	10.1	0.04
Respiratory System	495,104	83.2	0.12	78.7	0.11	423,041	59.6	0.09	56.9	0.09
Larynx	37,392	6.1	0.03	5.8	0.03	10,060	1.5	0.01	1.4	0.01
Lung & bronchus	450,926	75.9	0.12	71.7	0.11	408,808	57.5	0.09	54.9	0.09
Bones & Joints	5,810	1.2	0.02	1.1	0.02	4,551	0.9	0.01	0.9	0.01
Soft Tissue (incl. heart)	22,051	4.0	0.03	3.9	0.03	16,756	2.8	0.02	2.7	0.02
Skin (excl. basal & sq.)	201,710	35.3	0.08	33.7	0.08	136,648	22.5	0.06	22.3	0.06
Melanomas of the skin	187,736	32.8	0.08	31.4	0.07	128,264	21.3	0.06	21.1	0.06
Breast (excl. *in situ*)	7,711	1.3	0.02	1.2	0.01	841,438	128.7	0.15	125.4	0.14
Female Genital System	-	-	-	-	-	330,996	50.3	0.09	49.3	0.09
Cervix uteri	-	-	-	-	-	36,848	7.0	0.04	7.2	0.04
Uterus Corpus	-	-	-	-	-	177,674	25.9	0.06	25.3	0.06
Uterus, NOS[1]	-	-	-	-	-	4,962	0.7	0.01	0.7	0.01
Ovary	-	-	-	-	-	79,140	12.0	0.04	11.6	0.04
Male Genital System	716,009	114.9	0.14	112.7	0.14	-	-	-	-	-
Prostate	679,454	107.1	0.13	104.4	0.13	-	-	-	-	-
Testis	30,642	6.8	0.04	7.3	0.04	-	-	-	-	-
Penis	4,625	0.8	0.01	0.8	0.01	-	-	-	-	-
Urinary System	368,623	63.3	0.11	59.5	0.10	150,165	21.6	0.06	20.7	0.06
Bladder (incl. *in situ*)	229,998	39.8	0.08	36.9	0.08	70,592	9.7	0.04	9.1	0.04
Kidney & renal pelvis	130,539	22.1	0.06	21.4	0.06	75,325	11.3	0.04	11.0	0.04
Ureter	5,265	0.9	0.01	0.8	0.01	3,079	0.4	0.01	0.4	0.01
Eye & Orbit	6,292	1.1	0.01	1.1	0.01	5,229	0.9	0.01	0.8	0.01
Brain & Other Nervous System	47,193	8.7	0.04	8.5	0.04	37,151	6.3	0.03	6.2	0.03
Brain	44,831	8.2	0.04	8.1	0.04	34,684	5.8	0.03	5.7	0.03
Endocrine System	47,906	8.8	0.04	8.8	0.04	124,479	23.4	0.07	23.8	0.07
Thyroid gland	43,776	8.0	0.04	8.0	0.04	120,802	22.7	0.07	23.2	0.07
Lymphomas	153,036	27.2	0.07	26.1	0.07	125,163	19.0	0.06	18.3	0.05
Hodgkin Lymphoma	16,069	3.3	0.03	3.3	0.03	12,933	2.6	0.02	2.7	0.02
Non-Hodgkin Lymphoma	136,967	23.9	0.07	22.8	0.06	112,230	16.4	0.05	15.6	0.05
Multiple Myeloma	42,996	7.3	0.04	6.9	0.03	32,044	4.5	0.03	4.2	0.02
Leukemias	103,021	18.4	0.06	17.4	0.06	73,124	11.1	0.04	10.6	0.04
Acute lymphocytic	7,890	1.8	0.02	1.7	0.02	6,184	1.4	0.02	1.3	0.02
Chronic lymphocytic	39,767	6.7	0.03	6.3	0.03	25,740	3.5	0.02	3.3	0.02
Acute myeloid	30,053	5.4	0.03	5.1	0.03	23,550	3.6	0.02	3.5	0.02
Chronic myeloid	12,682	2.3	0.02	2.2	0.02	9,196	1.4	0.02	1.4	0.02
Other leukemias	12,629	2.3	0.02	2.1	0.02	8,454	1.2	0.01	1.1	0.01
Mesothelioma	10,542	1.9	0.02	1.7	0.02	3,210	0.5	0.01	0.4	0.01
Kaposi Sarcoma	2,174	0.4	0.01	0.4	0.01	307	0.0	0.00	0.0	0.00
Ill-defined & Unknown	118,846	20.9	0.06	19.2	0.06	104,488	14.4	0.05	13.3	0.04
Selected tumors (non-invasive)										
Brain & ONS[1] (*benign/borderline*)	47,713	8.7	0.04	8.4	0.04	94,799	14.9	0.05	14.5	0.05
Breast (*in situ*)	803	0.1	0.00	0.1	0.00	200,320	31.6	0.07	30.9	0.07

* Rates are per 100,000 population and are age-adjusted by five-year age groups to the 2000 U.S. standard population based on single years of age and to the 1996 Canadian population standard.

- Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for All Sites combined.

[1] Abbreviations: NOS - Not Otherwise Specified; ONS - Other Nervous System

N.B. In areas with small Latino populations, methods to indirectly identify Latinos (like NHIAv2) can overestimate the ethnicity-specific counts of cancer cases. Also, even small errors in Latino population estimates can affect the magnitude of the cancer rates. Data Item 190 used in place of NHIA for MA and RI, see technical notes.

Total Cancer Cases and Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates[1] for Commonly Diagnosed Cancers
Selected Areas in the NAACCR United States Combined[2], 2010-2014, Non-Hispanic White, Males
{Page 1 of 4}

	All Sites			Prostate			Colon & Rectum			Lung & Bronchus			Kidney & Renal Pelvis			Liver & IBD[3]		
	Count	*Rate*	*S.E.*[3]	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
NAACCR U.S. Combined	**3,019,830**	**510.8**	**0.30**	**679,454**	**107.1**	**0.13**	**263,005**	**45.2**	**0.09**	**450,926**	**75.9**	**0.12**	**130,539**	**22.1**	**0.06**	**62,694**	**10.0**	**0.04**
Alabama	51,296	523.6	2.38	11,375	108.3	1.04	4,837	50.2	0.74	9,250	93.1	0.99	2,222	22.8	0.50	1,048	10.2	0.32
Alaska	5,283	444.0	6.90	1,267	95.4	3.03	442	39.0	2.11	678	61.5	2.66	271	22.2	1.51	155	10.5	0.95
Arizona	58,197	428.3	1.83	11,759	80.6	0.76	5,142	38.4	0.55	8,520	60.1	0.66	2,620	19.7	0.40	1,229	8.7	0.25
Arkansas	34,928	512.8	2.81	7,865	108.3	1.24	3,246	48.4	0.87	6,749	97.0	1.20	1,570	23.4	0.61	660	9.3	0.37
California	255,755	504.1	1.02	59,883	110.1	0.46	21,396	42.5	0.30	28,557	56.3	0.34	10,354	20.4	0.21	5,880	10.7	0.14
CA: Greater California	160,953	504.3	1.29	37,199	108.4	0.57	13,518	42.8	0.38	19,129	59.6	0.44	6,624	20.8	0.26	3,688	10.7	0.18
CA: Greater Bay Area	48,827	513.4	2.39	12,314	120.0	1.11	3,752	39.8	0.67	4,470	48.1	0.74	1,856	19.2	0.46	1,183	11.3	0.34
CA: Los Angeles	45,837	493.2	2.35	10,328	105.2	1.05	4,118	44.4	0.70	4,942	53.0	0.76	1,874	20.3	0.48	1,004	10.3	0.33
Colorado	44,561	440.7	2.17	10,993	100.1	0.99	3,592	36.6	0.63	4,717	48.7	0.73	1,770	17.2	0.42	896	8.0	0.28
Connecticut	42,191	525.3	2.63	9,638	111.6	1.16	3,425	43.3	0.76	5,668	70.2	0.95	1,740	21.9	0.54	933	10.9	0.36
Delaware	11,435	578.7	5.57	2,755	128.3	2.49	839	43.4	1.54	1,671	83.4	2.08	477	25.2	1.20	264	13.1	0.83
District of Columbia	1,914	392.2	9.36	504	101.6	4.66	127	25.9	2.41	164	37.0	2.98	76	15.5	1.83	39	7.7	1.27
Florida	214,342	492.1	1.11	43,335	93.1	0.45	17,716	41.2	0.32	34,837	76.2	0.42	8,192	19.4	0.22	4,828	10.9	0.16
Georgia	83,284	536.8	1.93	18,487	110.2	0.84	7,240	47.2	0.57	13,374	87.1	0.78	3,420	21.8	0.38	1,708	10.3	0.26
Hawaii	5,719	520.9	7.38	1,072	87.7	2.85	459	41.2	2.05	586	52.9	2.33	210	19.2	1.42	147	11.4	1.00
Idaho	18,124	476.8	3.63	4,691	115.9	1.73	1,482	39.4	1.05	2,101	56.0	1.25	765	20.0	0.74	326	7.9	0.45
Illinois	127,145	529.9	1.52	28,724	112.1	0.68	12,361	52.1	0.48	19,540	81.8	0.60	5,763	23.8	0.32	2,296	9.1	0.19
Indiana	72,877	494.0	1.88	14,283	90.7	0.78	7,090	48.7	0.59	13,240	90.1	0.80	3,499	23.5	0.41	1,415	9.0	0.24
Iowa	42,909	531.0	2.62	9,538	110.8	1.15	4,132	52.0	0.82	6,403	78.8	1.00	1,975	24.6	0.57	708	8.3	0.32
Kansas	33,655	516.2	2.87	8,015	116.6	1.33	3,010	46.8	0.87	4,776	73.1	1.08	1,497	22.8	0.60	597	8.6	0.36
Kentucky	61,976	590.6	2.45	11,920	106.1	1.00	6,166	59.5	0.78	12,388	117.7	1.09	2,778	26.2	0.51	1,252	11.3	0.33
Louisiana	45,436	571.0	2.75	10,920	128.2	1.26	4,203	53.2	0.84	6,880	86.9	1.07	2,377	29.6	0.62	1,009	11.7	0.38
Maine	20,841	517.9	3.70	4,332	98.3	1.53	1,689	42.9	1.07	3,435	84.9	1.48	851	21.1	0.75	353	8.3	0.45
Maryland	49,572	511.6	2.36	12,000	114.3	1.07	3,957	41.6	0.68	6,557	68.4	0.86	2,009	20.6	0.47	1,011	9.8	0.32
Massachusetts	75,273	501.5	1.87	17,028	105.0	0.82	6,301	42.7	0.55	10,666	71.8	0.71	3,303	22.0	0.39	1,716	10.8	0.27
Michigan	114,049	500.5	1.52	27,507	112.2	0.69	9,369	42.1	0.45	17,274	75.8	0.59	4,802	21.0	0.31	2,008	8.3	0.19
MI: Metropolitan Detroit	42,235	574.7	2.87	10,874	138.9	1.37	3,292	45.4	0.81	5,997	82.9	1.10	1,780	23.9	0.58	808	10.1	0.37
Mississippi	28,407	543.8	3.31	6,230	111.0	1.43	2,766	53.7	1.05	5,147	97.8	1.39	1,345	26.0	0.73	713	13.0	0.50

Total Cancer Cases and Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates[1] for Commonly Diagnosed Cancers
Selected Areas in the NAACCR United States Combined[2], 2010-2014, Non-Hispanic White, Males
{Page 2 of 4}

	All Sites			Prostate			Colon & Rectum			Lung & Bronchus			Kidney & Renal Pelvis			Liver & IBD[3]		
	Count	*Rate*	*S.E.*[3]	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
Missouri	71,403	499.7	1.91	14,759	96.1	0.80	6,868	49.1	0.60	12,644	88.5	0.80	3,284	22.9	0.41	1,522	10.0	0.26
Montana	13,707	477.4	4.21	3,560	113.7	1.95	1,226	43.0	1.27	1,660	58.7	1.48	515	18.0	0.83	223	7.3	0.51
Nebraska	22,209	505.7	3.46	5,380	116.4	1.61	2,170	49.6	1.09	3,152	72.1	1.30	988	22.2	0.72	325	7.1	0.40
New Hampshire	19,094	527.2	3.95	4,864	121.7	1.80	1,449	40.9	1.11	2,526	71.8	1.47	756	20.4	0.77	326	8.2	0.47
New Jersey	91,954	565.0	1.91	22,800	131.7	0.89	7,865	48.8	0.56	11,355	69.6	0.66	3,731	22.7	0.38	1,739	10.1	0.25
New York	195,557	565.9	1.31	46,407	126.5	0.60	15,846	46.5	0.38	25,889	74.5	0.47	8,504	24.6	0.27	3,658	10.0	0.17
North Carolina	99,375	522.2	1.70	22,424	109.1	0.75	7,847	42.1	0.49	16,780	87.8	0.69	4,274	22.4	0.35	2,199	11.0	0.24
North Dakota	9,193	510.0	5.44	2,354	123.9	2.61	947	53.8	1.79	1,236	69.5	2.01	384	21.9	1.15	101	5.2	0.53
Ohio	135,141	494.8	1.38	28,942	98.8	0.59	12,706	47.4	0.43	22,859	83.6	0.56	5,950	21.8	0.29	2,501	8.7	0.18
Oklahoma	39,832	488.4	2.50	8,557	98.8	1.09	3,739	46.6	0.78	7,060	85.4	1.03	1,796	22.3	0.54	814	9.3	0.33
Oregon	46,205	472.0	2.26	10,680	100.4	1.00	3,848	40.4	0.67	6,175	63.4	0.83	2,026	20.7	0.48	1,232	11.2	0.33
Pennsylvania	169,294	534.6	1.33	36,108	106.4	0.57	15,486	49.5	0.41	24,941	77.9	0.50	7,100	22.7	0.28	3,381	10.1	0.18
Rhode Island	12,785	517.6	4.71	2,650	100.0	1.99	984	40.3	1.31	1,961	79.3	1.83	562	23.1	1.01	308	11.7	0.68
South Carolina	48,456	513.0	2.41	10,510	101.8	1.02	3,933	42.5	0.70	8,223	85.8	0.97	2,105	22.6	0.51	1,112	11.4	0.35
South Dakota	10,655	500.3	4.96	2,667	117.9	2.33	1,072	51.0	1.59	1,454	68.2	1.82	440	21.0	1.03	136	6.0	0.53
Tennessee	76,536	531.1	1.98	17,392	111.8	0.87	6,475	46.1	0.59	13,953	96.2	0.84	3,394	23.2	0.41	1,624	10.6	0.27
Texas	173,390	497.0	1.22	37,569	100.6	0.53	15,679	45.5	0.37	26,305	75.6	0.48	8,467	24.0	0.27	4,652	12.2	0.18
Utah	23,064	463.5	3.12	6,737	131.5	1.63	1,709	35.2	0.87	1,544	32.3	0.84	912	18.1	0.61	375	7.2	0.38
Vermont	8,892	492.1	5.41	1,935	97.6	2.29	705	39.4	1.54	1,289	71.4	2.05	370	20.9	1.13	170	8.6	0.69
Virginia	69,397	453.4	1.77	15,501	93.3	0.77	6,065	40.1	0.53	11,082	73.2	0.71	2,966	19.2	0.36	1,393	8.6	0.24
Washington	76,573	510.8	1.90	18,744	115.5	0.87	6,005	40.6	0.54	9,793	66.7	0.69	3,238	21.5	0.39	1,857	11.1	0.27
WA: Seattle/Puget Sound	53,261	539.7	2.41	13,323	125.3	1.12	4,049	41.4	0.67	6,422	67.2	0.86	2,201	22.2	0.49	1,276	11.6	0.33
West Virginia	28,396	525.7	3.22	5,708	97.9	1.33	2,813	53.0	1.03	5,467	99.6	1.38	1,243	23.2	0.68	525	9.3	0.42
Wisconsin	72,976	508.7	1.93	17,341	112.4	0.87	5,999	42.6	0.56	9,700	67.7	0.70	3,380	23.4	0.41	1,226	8.2	0.24
Wyoming	6,577	461.4	5.91	1,744	112.7	2.80	582	41.9	1.80	700	50.6	1.98	268	18.8	1.19	104	6.6	0.67

1. Rates are per 100,000 population and are age-adjusted by five-year age groups to the 2000 U.S. standard population based on single years of age.
2. See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.
3. Abbreviations: IBD - Intrahepatic Bile Duct; S.E. - Standard Error
- Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for the U.S. combined.

N.B. In areas with small Latino populations, methods to indirectly identify Latinos (like NHIAv2) can overestimate the ethnicity-specific counts of cancer cases. Also, even small errors in Latino population estimates can affect the magnitude of the cancer rates. Data Item 190 used in place of NHIA for MA and RI, see technical notes.

Total Cancer Cases and Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates[1] for Commonly Diagnosed Cancers
Selected Areas in the NAACCR United States Combined[2], 2010-2014, Non-Hispanic White, Males
{Page 3 of 4}

	Non-Hodgkin Lymphoma			Leukemia			Bladder			Stomach			Oral Cavity & Pharynx			Pancreas		
	Count	*Rate*	*S.E.*[3]	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
NAACCR U.S. Combined	**136,967**	**23.9**	**0.07**	**103,021**	**18.4**	**0.06**	**229,998**	**39.8**	**0.08**	**46,073**	**7.8**	**0.04**	**115,165**	**18.8**	**0.06**	**85,017**	**14.3**	**0.05**
Alabama	1,960	20.7	0.48	1,463	16.1	0.43	3,567	37.6	0.65	722	7.5	0.29	2,092	21.0	0.47	1,408	14.1	0.39
Alaska	239	20.7	1.50	158	14.6	1.28	388	37.3	2.12	77	6.9	0.87	223	15.7	1.18	152	14.2	1.31
Arizona	2,516	19.2	0.40	1,868	14.8	0.36	5,030	36.0	0.52	874	6.4	0.22	2,106	15.6	0.35	1,849	13.1	0.31
Arkansas	1,352	20.4	0.57	1,025	15.9	0.51	2,491	37.3	0.76	491	7.3	0.34	1,394	20.5	0.56	902	13.4	0.45
California	12,240	24.9	0.23	8,724	18.3	0.20	19,685	39.4	0.29	3,973	7.8	0.13	10,094	19.3	0.20	7,181	14.0	0.17
CA: Greater California	7,335	23.7	0.29	5,340	17.8	0.25	12,277	39.1	0.36	2,419	7.6	0.16	6,539	19.9	0.25	4,441	13.7	0.21
CA: Greater Bay Area	2,557	27.7	0.57	1,731	19.5	0.49	3,570	38.4	0.66	725	7.7	0.29	1,816	18.1	0.44	1,396	14.7	0.40
CA: Los Angeles	2,344	26.0	0.55	1,640	18.6	0.47	3,833	41.4	0.68	828	8.9	0.31	1,738	18.3	0.45	1,341	14.3	0.40
Colorado	2,138	21.8	0.49	1,783	18.7	0.46	3,231	34.2	0.62	648	6.6	0.27	1,757	16.1	0.40	1,293	12.9	0.37
Connecticut	2,016	26.0	0.60	1,520	19.8	0.52	3,959	49.5	0.80	810	10.0	0.36	1,459	17.4	0.47	1,223	15.0	0.44
Delaware	510	26.9	1.23	370	19.9	1.07	911	46.4	1.57	156	7.6	0.62	423	21.3	1.07	325	16.0	0.90
District of Columbia	114	23.8	2.40	58	12.0	1.66	121	26.3	2.47	28	6.5	1.28	67	14.1	1.78	61	13.3	1.77
Florida	8,896	21.4	0.24	6,936	17.0	0.22	16,827	36.6	0.29	2,881	6.5	0.13	9,718	23.3	0.25	6,305	14.0	0.18
Georgia	3,432	22.9	0.40	2,717	18.7	0.37	5,573	38.0	0.53	1,067	7.0	0.22	3,391	20.7	0.37	2,152	14.1	0.32
Hawaii	256	23.9	1.61	178	18.1	1.48	401	38.7	2.03	70	6.3	0.80	293	24.8	1.54	164	14.8	1.23
Idaho	815	21.8	0.78	710	19.6	0.75	1,411	38.5	1.04	253	6.8	0.44	675	17.2	0.68	538	14.1	0.62
Illinois	5,831	25.0	0.34	4,056	18.0	0.29	9,833	42.1	0.43	2,068	8.7	0.20	4,790	19.1	0.28	3,652	15.2	0.26
Indiana	3,351	23.4	0.41	2,394	17.2	0.36	5,384	37.8	0.53	1,057	7.2	0.23	2,807	18.1	0.35	2,121	14.3	0.32
Iowa	2,132	27.0	0.60	1,634	20.9	0.53	3,095	38.6	0.70	547	6.8	0.30	1,581	19.1	0.49	1,187	14.6	0.43
Kansas	1,534	24.3	0.63	1,270	20.4	0.58	2,558	39.9	0.80	466	7.2	0.34	1,227	18.3	0.54	850	13.1	0.46
Kentucky	2,605	25.4	0.51	2,030	20.5	0.47	4,109	41.3	0.66	938	9.1	0.31	2,442	22.1	0.46	1,585	15.1	0.39
Louisiana	1,973	25.8	0.59	1,398	18.7	0.51	2,965	38.7	0.73	655	8.4	0.34	1,820	22.2	0.53	1,220	15.5	0.45
Maine	891	22.8	0.79	760	20.1	0.75	1,867	47.7	1.13	289	7.1	0.43	783	18.6	0.69	581	14.4	0.61
Maryland	2,158	22.9	0.51	1,543	17.0	0.45	4,016	42.8	0.69	730	7.6	0.29	1,854	18.4	0.44	1,425	14.8	0.40
Massachusetts	3,473	23.9	0.42	2,372	16.7	0.35	6,238	42.6	0.55	1,277	8.7	0.25	2,842	18.1	0.35	2,125	14.1	0.31
Michigan	5,422	24.8	0.35	3,912	18.3	0.30	9,033	41.0	0.44	1,936	8.6	0.20	4,165	17.5	0.28	3,233	14.1	0.25
MI: Metropolitan Detroit	2,020	28.7	0.66	1,532	22.0	0.58	3,412	47.9	0.84	731	10.0	0.38	1,517	19.5	0.52	1,185	16.2	0.48
Mississippi	1,142	22.9	0.70	828	16.7	0.60	1,822	36.0	0.86	390	7.5	0.39	1,141	21.5	0.65	827	15.9	0.57

Total Cancer Cases and Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates[1] for Commonly Diagnosed Cancers
Selected Areas in the NAACCR United States Combined[2], 2010-2014, Non-Hispanic White, Males
{Page 4 of 4}

	Non-Hodgkin Lymphoma			Leukemia			Bladder			Stomach			Oral Cavity & Pharynx			Pancreas		
	Count	*Rate*	*S.E.*[3]	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
Missouri	3,159	22.9	0.42	2,403	17.6	0.37	4,904	35.4	0.51	1,002	7.0	0.23	2,804	18.9	0.36	1,996	14.0	0.32
Montana	642	22.9	0.94	570	20.6	0.89	1,008	35.9	1.16	155	5.5	0.45	461	15.7	0.76	356	12.3	0.68
Nebraska	1,072	25.1	0.78	781	18.6	0.68	1,640	38.3	0.96	322	7.4	0.42	761	16.9	0.63	603	13.8	0.57
New Hampshire	888	25.6	0.89	641	18.8	0.77	1,672	48.4	1.22	263	7.3	0.47	669	17.0	0.68	508	14.1	0.65
New Jersey	4,296	27.4	0.43	3,138	20.2	0.37	7,550	46.9	0.55	1,596	9.8	0.25	2,840	16.8	0.32	2,668	16.1	0.32
New York	9,480	28.3	0.30	7,823	23.6	0.27	16,558	48.4	0.38	3,398	9.8	0.17	5,923	16.7	0.22	5,776	16.6	0.22
North Carolina	4,124	22.5	0.36	3,198	17.9	0.32	7,161	38.8	0.47	1,380	7.3	0.20	3,850	19.5	0.32	2,647	13.9	0.28
North Dakota	407	22.8	1.16	376	21.4	1.13	664	37.3	1.47	142	8.0	0.68	322	17.4	1.00	229	12.9	0.87
Ohio	6,236	23.5	0.30	4,022	15.6	0.25	10,543	39.8	0.40	2,298	8.5	0.18	5,066	17.8	0.26	3,827	14.0	0.23
Oklahoma	1,763	22.3	0.54	1,318	17.2	0.48	2,803	35.0	0.67	588	7.2	0.30	1,688	20.4	0.51	1,040	12.8	0.40
Oregon	2,129	22.6	0.50	1,485	16.2	0.43	3,706	38.7	0.65	696	7.2	0.28	1,736	17.0	0.42	1,370	13.9	0.39
Pennsylvania	8,111	26.2	0.30	5,860	19.4	0.26	14,418	45.9	0.39	2,886	9.1	0.17	5,783	17.9	0.24	4,992	15.6	0.22
Rhode Island	646	26.9	1.09	426	18.3	0.92	1,163	47.0	1.41	250	10.1	0.65	433	16.8	0.83	374	14.8	0.78
South Carolina	1,926	21.2	0.50	1,610	18.4	0.48	3,605	39.8	0.68	701	7.5	0.29	1,991	20.7	0.48	1,318	14.2	0.40
South Dakota	500	24.5	1.12	414	20.2	1.02	754	36.2	1.34	138	6.3	0.55	415	18.9	0.96	284	13.1	0.79
Tennessee	3,078	22.2	0.41	2,513	18.6	0.38	5,131	37.0	0.53	1,040	7.4	0.24	3,064	20.5	0.38	1,900	13.2	0.31
Texas	7,931	23.5	0.27	6,373	19.5	0.25	11,165	33.4	0.32	2,380	6.9	0.14	7,490	20.4	0.24	4,861	13.9	0.20
Utah	1,147	23.4	0.71	913	18.3	0.62	1,438	31.1	0.83	244	5.0	0.33	648	12.7	0.51	607	12.6	0.52
Vermont	463	26.4	1.27	283	16.8	1.04	682	39.1	1.54	127	7.4	0.67	341	17.7	1.00	247	14.0	0.91
Virginia	3,241	21.9	0.40	2,060	14.5	0.33	4,929	33.9	0.50	1,085	7.2	0.22	2,876	17.9	0.34	1,996	13.2	0.30
Washington	3,737	25.8	0.43	2,820	19.9	0.39	5,662	39.5	0.54	1,159	7.8	0.24	2,882	18.2	0.35	2,105	14.1	0.32
WA: Seattle/Puget Sound	2,630	27.4	0.55	2,027	21.5	0.49	3,867	41.4	0.68	811	8.3	0.30	1,957	18.5	0.43	1,407	14.4	0.39
West Virginia	1,152	22.0	0.67	937	18.6	0.63	2,131	40.6	0.90	472	8.7	0.41	1,099	19.8	0.62	718	13.2	0.51
Wisconsin	3,560	25.6	0.44	3,108	22.7	0.42	5,692	41.0	0.55	1,242	8.8	0.25	2,661	17.7	0.35	2,099	14.6	0.32
Wyoming	283	20.5	1.26	242	17.0	1.14	504	37.5	1.73	106	7.7	0.78	218	14.6	1.03	137	9.5	0.84

1. Rates are per 100,000 population and are age-adjusted by five-year age groups to the 2000 U.S. standard population based on single years of age.
2. See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.
3. Abbreviations: S.E. - Standard Error

- Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for the U.S. combined.

N.B. In areas with small Latino populations, methods to indirectly identify Latinos (like NHIAv2) can overestimate the ethnicity-specific counts of cancer cases. Also, even small errors in Latino population estimates can affect the magnitude of the cancer rates. Data Item 190 used in place of NHIA for MA and RI, see technical notes.

Total Cancer Cases and Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates[1] for Commonly Diagnosed Cancers
Selected Areas in the NAACCR United States Combined[2], 2010-2014, Non-Hispanic White, Females
{Page 1 of 4}

	All Sites			Breast			Colon & Rectum			Thyroid			Uterus Corpus & Uterus NOS[3]			Lung & Bronchus		
	Count	*Rate*	*S.E.[3]*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
NAACCR U.S. Combined	**2,897,505**	**436.0**	**0.27**	**841,438**	**128.7**	**0.15**	**241,669**	**34.5**	**0.07**	**120,802**	**22.7**	**0.07**	**182,636**	**26.6**	**0.06**	**408,808**	**57.5**	**0.09**
Alabama	45,033	403.8	1.97	13,076	118.2	1.07	4,016	35.0	0.57	1,302	14.3	0.41	2,168	18.9	0.42	6,820	56.7	0.70
Alaska	4,678	408.2	6.35	1,508	125.9	3.45	375	33.4	1.84	191	15.7	1.18	341	27.0	1.56	564	53.8	2.42
Arizona	55,888	390.6	1.76	16,896	120.5	0.99	4,566	30.0	0.47	2,660	25.2	0.52	3,121	21.0	0.40	7,994	49.7	0.57
Arkansas	29,847	393.9	2.37	8,288	110.5	1.26	2,780	35.2	0.69	946	15.5	0.53	1,641	21.5	0.55	5,088	62.2	0.89
California	243,212	436.8	0.93	75,781	139.0	0.53	19,897	33.7	0.25	9,100	20.9	0.23	15,188	26.1	0.22	29,338	48.7	0.29
CA: Greater California	152,231	433.4	1.17	47,034	136.1	0.66	12,414	33.5	0.31	5,608	20.7	0.29	9,210	25.2	0.27	19,488	51.2	0.38
CA: Greater Bay Area	46,228	443.2	2.16	14,552	143.1	1.24	3,675	33.3	0.58	1,484	17.8	0.49	3,033	27.4	0.51	5,055	45.4	0.66
CA: Los Angeles	44,651	441.3	2.19	14,173	144.8	1.27	3,800	34.7	0.59	2,007	24.9	0.58	2,944	28.4	0.54	4,786	43.3	0.64
Colorado	44,852	397.7	1.94	14,484	127.1	1.09	3,450	30.4	0.53	2,386	23.8	0.50	2,746	22.7	0.45	4,883	42.5	0.62
Connecticut	43,718	466.7	2.36	13,019	143.3	1.32	3,409	33.5	0.61	2,130	29.9	0.69	2,889	29.3	0.56	5,999	59.7	0.79
Delaware	10,536	474.0	4.87	3,010	135.8	2.61	771	33.1	1.24	414	25.2	1.32	692	29.4	1.16	1,599	66.2	1.69
District of Columbia	2,017	401.6	9.46	763	157.6	5.95	131	26.7	2.46	151	25.0	2.34	120	24.0	2.28	181	37.8	2.94
Florida	193,430	419.3	1.04	53,659	121.3	0.57	15,804	31.5	0.27	5,392	17.2	0.26	10,643	22.8	0.24	32,402	62.2	0.36
Georgia	75,807	428.9	1.60	22,256	125.2	0.86	6,097	33.8	0.44	2,978	19.6	0.37	3,944	21.3	0.35	10,795	57.9	0.57
Hawaii	4,414	459.7	7.47	1,375	143.3	4.17	343	35.4	2.05	160	20.4	1.73	252	24.0	1.62	450	44.4	2.21
Idaho	16,735	413.6	3.29	4,936	121.9	1.79	1,323	32.3	0.91	871	25.3	0.88	1,014	24.0	0.78	2,031	47.7	1.08
Illinois	126,890	454.5	1.32	37,050	135.7	0.73	11,169	37.4	0.37	5,358	23.8	0.34	8,568	29.7	0.33	18,092	61.4	0.47
Indiana	73,597	430.1	1.63	20,471	120.9	0.87	6,800	38.2	0.48	2,669	18.8	0.38	4,873	27.6	0.41	11,090	61.9	0.60
Iowa	40,612	443.6	2.30	11,095	124.7	1.24	3,975	40.0	0.66	1,476	20.6	0.56	2,914	31.3	0.60	5,201	53.4	0.76
Kansas	31,511	429.4	2.52	9,020	124.4	1.36	2,791	36.0	0.71	1,502	25.1	0.68	1,854	24.8	0.60	4,154	53.1	0.84
Kentucky	58,249	479.0	2.04	15,099	124.3	1.04	5,272	42.2	0.59	2,329	22.5	0.48	3,265	26.3	0.47	10,334	80.9	0.81
Louisiana	38,390	428.5	2.25	10,881	121.8	1.20	3,534	38.5	0.66	1,723	22.4	0.56	1,656	18.1	0.46	5,500	58.0	0.79
Maine	20,631	456.0	3.31	5,641	126.5	1.76	1,610	34.4	0.89	784	22.1	0.84	1,484	31.1	0.84	3,136	65.4	1.19
Maryland	49,120	445.8	2.10	14,755	135.2	1.16	3,719	32.3	0.55	2,282	25.8	0.57	3,186	27.6	0.50	6,679	57.2	0.72
Massachusetts	81,887	459.7	1.68	24,646	141.8	0.94	6,330	33.5	0.44	4,149	29.0	0.47	5,491	29.8	0.41	11,934	63.5	0.60
Michigan	109,124	426.4	1.34	30,809	122.0	0.72	8,784	32.7	0.36	4,324	21.0	0.33	7,798	29.4	0.34	15,976	58.8	0.47
MI: Metropolitan Detroit	40,427	468.3	2.42	11,314	132.8	1.29	3,236	35.2	0.64	1,893	27.0	0.65	2,801	31.6	0.62	6,019	66.3	0.87
Mississippi	24,258	414.5	2.76	6,582	113.4	1.45	2,307	37.6	0.81	870	18.5	0.66	1,109	18.8	0.59	3,849	61.0	1.00

Total Cancer Cases and Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates[1] for Commonly Diagnosed Cancers
Selected Areas in the NAACCR United States Combined[2], 2010-2014, Non-Hispanic White, Females
{Page 2 of 4}

	All Sites			Breast			Colon & Rectum			Thyroid			Uterus Corpus & Uterus NOS[3]			Lung & Bronchus		
	Count	*Rate*	*S.E.*[3]	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
Missouri	69,758	429.8	1.68	20,169	126.0	0.92	6,135	36.2	0.48	2,492	18.7	0.39	4,370	26.3	0.41	11,102	64.7	0.62
Montana	12,394	418.2	3.93	3,613	122.6	2.14	1,012	32.9	1.07	511	21.8	1.02	796	25.4	0.94	1,635	51.8	1.31
Nebraska	20,669	417.8	3.03	6,026	123.7	1.66	2,002	38.0	0.88	850	21.4	0.76	1,369	27.3	0.76	2,623	50.0	1.00
New Hampshire	18,855	465.1	3.51	5,757	142.3	1.94	1,429	34.1	0.93	784	22.9	0.86	1,420	33.1	0.90	2,673	64.1	1.27
New Jersey	93,171	485.8	1.68	26,345	142.3	0.92	8,013	38.1	0.45	4,722	32.1	0.50	6,657	33.6	0.42	12,213	58.9	0.55
New York	195,568	488.1	1.16	54,058	139.6	0.63	15,687	36.4	0.30	9,754	31.1	0.33	13,416	32.7	0.29	26,970	62.8	0.39
North Carolina	94,102	429.8	1.45	28,293	130.3	0.80	7,200	32.0	0.39	3,587	19.8	0.34	5,548	24.4	0.34	13,898	59.4	0.51
North Dakota	8,029	412.8	4.82	2,274	122.2	2.69	820	39.3	1.43	396	24.8	1.30	500	25.3	1.17	968	47.0	1.55
Ohio	135,034	428.7	1.21	38,491	123.7	0.65	11,766	35.3	0.34	5,765	22.6	0.31	9,536	29.4	0.31	19,727	59.4	0.43
Oklahoma	36,919	402.4	2.17	10,453	114.8	1.17	3,343	34.9	0.62	1,342	18.0	0.51	1,955	20.9	0.49	5,670	57.7	0.78
Oregon	45,732	425.1	2.08	13,795	128.5	1.15	3,487	30.9	0.55	1,671	19.6	0.50	3,004	26.6	0.51	6,213	54.1	0.70
Pennsylvania	169,465	463.8	1.19	46,936	131.8	0.64	14,759	37.1	0.32	8,946	32.0	0.36	12,343	33.0	0.31	22,071	56.0	0.39
Rhode Island	13,963	473.0	4.23	3,868	135.6	2.29	1,133	34.8	1.10	663	29.0	1.19	1,010	33.2	1.08	2,095	66.2	1.49
South Carolina	44,749	425.1	2.09	13,544	128.8	1.15	3,542	32.6	0.56	1,654	19.4	0.50	2,416	22.2	0.47	6,615	58.3	0.73
South Dakota	9,893	431.1	4.55	2,937	132.1	2.56	960	38.0	1.29	383	21.0	1.12	675	29.1	1.16	1,218	50.0	1.47
Tennessee	70,086	427.5	1.66	19,895	121.8	0.89	6,044	35.8	0.47	2,812	20.6	0.40	3,919	23.4	0.38	11,041	63.3	0.61
Texas	158,721	410.4	1.06	47,036	122.5	0.58	13,062	32.6	0.29	6,612	20.3	0.26	8,313	20.8	0.23	22,117	53.8	0.37
Utah	20,470	376.7	2.67	6,265	116.8	1.50	1,467	27.2	0.72	1,580	29.3	0.75	1,426	25.5	0.69	1,272	23.4	0.66
Vermont	8,806	438.3	4.87	2,599	130.5	2.67	699	33.2	1.30	304	19.2	1.16	701	33.2	1.30	1,247	59.2	1.72
Virginia	70,635	408.8	1.58	22,334	130.0	0.90	5,672	32.0	0.44	2,969	20.4	0.39	4,531	24.9	0.38	9,781	54.2	0.56
Washington	74,159	450.0	1.71	22,925	139.3	0.95	5,568	33.1	0.46	2,816	20.3	0.40	4,603	26.3	0.40	9,567	55.7	0.58
WA: Seattle/Puget Sound	50,966	464.5	2.13	15,861	144.5	1.19	3,731	33.4	0.57	1,899	20.1	0.48	3,167	27.0	0.50	6,318	55.9	0.72
West Virginia	26,730	447.7	2.85	6,838	115.1	1.45	2,575	41.2	0.84	1,183	24.7	0.75	1,865	30.7	0.74	4,297	67.1	1.04
Wisconsin	69,488	435.1	1.71	20,182	129.1	0.94	5,583	32.9	0.46	2,575	19.9	0.41	4,977	29.8	0.43	9,022	53.8	0.58
Wyoming	5,673	390.3	5.37	1,705	116.1	2.92	458	31.2	1.50	284	22.5	1.39	329	21.7	1.24	684	45.4	1.77

1. Rates are per 100,000 population and are age-adjusted by five-year age groups to the 2000 U.S. standard population based on single years of age.
2. See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.
3. Abbreviations: NOS - Not Otherwise Specified; S.E. - Standard Error
- Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for the U.S. combined.

N.B. In areas with small Latino populations, methods to indirectly identify Latinos (like NHIAv2) can overestimate the ethnicity-specific counts of cancer cases. Also, even small errors in Latino population estimates can affect the magnitude of the cancer rates. Data Item 190 used in place of NHIA for MA and RI, see technical notes.

Total Cancer Cases and Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates[1] for Commonly Diagnosed Cancers
Selected Areas in the NAACCR United States Combined[2], 2010-2014, Non-Hispanic White, Females
{Page 3 of 4}

	Non-Hodgkin Lymphoma			Kidney & Renal Pelvis			Cervix			Ovary			Leukemia			Pancreas		
	Count	*Rate*	*S.E.*[3]	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
NAACCR U.S. Combined	**112,230**	**16.4**	**0.05**	**75,325**	**11.3**	**0.04**	**36,848**	**7.0**	**0.04**	**79,140**	**12.0**	**0.04**	**73,124**	**11.1**	**0.04**	**78,507**	**10.8**	**0.04**
Alabama	1,709	14.7	0.37	1,370	12.3	0.35	757	8.7	0.33	1,361	12.3	0.34	1,057	9.9	0.32	1,232	10.3	0.30
Alaska	164	14.4	1.20	121	9.9	0.96	71	6.2	0.76	148	13.3	1.17	109	10.7	1.07	122	10.9	1.05
Arizona	2,033	13.6	0.32	1,419	9.8	0.28	615	5.8	0.25	1,648	11.4	0.30	1,261	9.3	0.29	1,596	9.9	0.26
Arkansas	1,183	15.2	0.45	925	12.4	0.43	493	8.7	0.41	763	10.3	0.39	687	9.3	0.37	749	9.0	0.34
California	9,290	16.3	0.18	5,309	9.5	0.14	2,864	6.7	0.13	6,951	12.6	0.16	5,897	10.7	0.15	6,484	10.5	0.13
CA: Greater California	5,585	15.5	0.22	3,534	10.0	0.18	1,860	7.0	0.17	4,276	12.3	0.20	3,693	10.7	0.19	4,038	10.4	0.17
CA: Greater Bay Area	1,890	18.0	0.43	923	8.7	0.30	459	5.5	0.27	1,338	12.8	0.37	1,124	11.0	0.35	1,220	10.3	0.31
CA: Los Angeles	1,809	17.2	0.42	848	8.4	0.30	542	6.8	0.31	1,335	13.5	0.39	1,071	10.6	0.35	1,225	10.7	0.32
Colorado	1,678	14.7	0.37	1,018	9.0	0.29	563	5.7	0.25	1,296	11.4	0.33	1,211	11.1	0.33	1,128	9.7	0.30
Connecticut	1,661	17.2	0.44	925	9.9	0.34	428	5.9	0.31	1,182	12.4	0.38	1,152	12.2	0.39	1,254	12.0	0.35
Delaware	425	18.5	0.94	255	11.7	0.78	147	8.6	0.76	263	11.9	0.77	235	10.8	0.76	268	10.8	0.67
District of Columbia	57	11.5	1.63	31	6.0	1.15	22	4.5	0.99	60	12.4	1.67	34	6.1	1.16	44	9.0	1.43
Florida	7,047	14.5	0.19	4,474	9.7	0.16	2,634	8.6	0.18	5,330	11.8	0.18	4,724	10.5	0.17	5,449	10.2	0.14
Georgia	2,800	15.7	0.30	1,893	10.6	0.25	1,119	7.6	0.24	2,148	12.2	0.27	1,957	11.4	0.26	1,921	10.3	0.24
Hawaii	174	17.4	1.43	87	8.8	1.02	51	6.7	1.00	107	12.0	1.26	117	13.1	1.33	107	10.3	1.04
Idaho	638	15.6	0.63	434	10.5	0.52	201	5.9	0.43	476	11.5	0.54	522	12.9	0.58	487	11.4	0.53
Illinois	4,926	17.1	0.25	3,481	12.4	0.22	1,628	7.1	0.18	3,383	12.2	0.22	3,000	10.8	0.21	3,405	11.1	0.20
Indiana	2,856	16.3	0.31	2,225	13.0	0.28	1,052	7.5	0.24	1,947	11.4	0.27	1,837	11.0	0.26	1,997	10.9	0.25
Iowa	1,741	18.4	0.46	1,104	12.0	0.38	480	7.0	0.33	1,088	11.9	0.38	1,162	12.4	0.38	1,146	11.2	0.34
Kansas	1,282	16.8	0.49	930	12.9	0.44	395	6.9	0.36	847	11.7	0.42	912	12.3	0.42	802	10.1	0.37
Kentucky	2,123	17.2	0.38	1,721	14.5	0.36	893	8.9	0.31	1,285	10.6	0.30	1,532	12.8	0.34	1,407	11.0	0.30
Louisiana	1,688	18.4	0.46	1,484	16.5	0.44	565	7.8	0.34	911	10.2	0.35	1,031	11.8	0.38	1,129	11.8	0.36
Maine	840	18.3	0.66	505	11.2	0.52	220	6.2	0.44	480	10.6	0.50	552	12.4	0.56	620	12.6	0.52
Maryland	1,838	16.4	0.40	1,127	10.2	0.32	524	5.7	0.26	1,257	11.4	0.33	1,217	11.0	0.33	1,276	10.8	0.31
Massachusetts	3,075	16.7	0.31	1,916	10.8	0.26	675	4.7	0.19	2,125	11.9	0.27	1,714	9.9	0.25	2,222	11.2	0.25
Michigan	4,550	17.4	0.27	2,933	11.5	0.22	1,244	6.2	0.19	3,156	12.5	0.23	2,798	11.1	0.22	3,008	10.8	0.20
MI: Metropolitan Detroit	1,679	19.0	0.48	1,063	12.3	0.39	447	6.5	0.32	1,248	14.5	0.43	1,097	12.9	0.41	1,177	12.5	0.38
Mississippi	949	15.6	0.52	842	14.5	0.52	387	8.6	0.46	605	10.5	0.44	582	10.1	0.44	730	11.4	0.43

Total Cancer Cases and Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates[1] for Commonly Diagnosed Cancers
Selected Areas in the NAACCR United States Combined[2], 2010-2014, Non-Hispanic White, Females
{Page 4 of 4}

	Non-Hodgkin Lymphoma			Kidney & Renal Pelvis			Cervix			Ovary			Leukemia			Pancreas		
	Count	*Rate*	*S.E.*[3]	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
Missouri	2,595	15.6	0.32	2,025	12.5	0.29	1,065	8.3	0.26	1,700	10.5	0.26	1,768	11.0	0.27	1,847	10.6	0.25
Montana	489	16.3	0.77	292	10.1	0.62	149	5.9	0.52	330	10.8	0.62	325	11.1	0.64	323	10.1	0.58
Nebraska	866	16.8	0.59	556	11.3	0.50	273	7.1	0.45	547	11.1	0.50	537	10.6	0.48	551	10.3	0.45
New Hampshire	731	17.6	0.67	433	10.6	0.53	157	4.5	0.38	468	11.5	0.55	417	10.8	0.56	440	10.4	0.51
New Jersey	3,788	19.1	0.33	2,095	10.8	0.25	1,049	6.8	0.23	2,593	13.5	0.28	2,331	12.2	0.27	2,689	12.4	0.25
New York	8,035	19.3	0.22	4,698	11.7	0.18	2,072	6.3	0.15	5,423	13.7	0.19	5,609	13.8	0.20	5,648	12.5	0.17
North Carolina	3,337	14.9	0.27	2,516	11.6	0.24	1,192	6.8	0.20	2,496	11.4	0.24	2,329	10.9	0.24	2,321	9.9	0.21
North Dakota	380	18.3	0.98	219	11.0	0.77	67	4.3	0.55	190	10.1	0.77	216	10.3	0.74	204	9.3	0.68
Ohio	5,108	15.7	0.23	3,746	12.0	0.20	1,909	7.5	0.18	3,696	11.7	0.20	2,894	9.4	0.18	3,752	10.9	0.18
Oklahoma	1,400	14.6	0.40	1,132	12.3	0.38	599	8.6	0.37	1,061	11.5	0.37	984	10.8	0.36	999	10.0	0.32
Oregon	1,782	16.1	0.40	1,116	10.4	0.33	538	6.5	0.29	1,286	12.0	0.35	1,007	9.7	0.32	1,246	10.8	0.32
Pennsylvania	6,923	18.0	0.23	4,257	11.7	0.19	1,899	6.7	0.16	4,632	12.8	0.20	4,326	12.0	0.19	4,788	11.5	0.17
Rhode Island	567	18.3	0.81	349	12.2	0.69	152	6.9	0.60	331	11.2	0.65	346	11.9	0.70	329	10.0	0.57
South Carolina	1,630	15.1	0.39	1,204	11.4	0.34	603	7.4	0.32	1,106	10.6	0.33	1,094	10.7	0.34	1,171	10.3	0.31
South Dakota	386	16.5	0.88	234	10.2	0.70	112	6.0	0.60	247	10.5	0.70	273	11.8	0.75	271	10.4	0.66
Tennessee	2,602	15.4	0.31	2,012	12.3	0.28	1,080	8.1	0.26	1,905	11.8	0.28	1,730	10.9	0.27	1,833	10.5	0.25
Texas	6,206	15.7	0.20	4,914	12.6	0.18	2,636	8.5	0.17	4,415	11.5	0.18	4,501	11.8	0.18	4,252	10.2	0.16
Utah	832	15.1	0.53	506	9.2	0.41	240	4.6	0.30	642	11.8	0.47	635	11.5	0.46	514	9.4	0.42
Vermont	360	17.6	0.96	185	9.1	0.70	78	4.7	0.56	245	12.3	0.82	195	10.3	0.78	228	11.0	0.75
Virginia	2,568	14.6	0.30	1,677	9.8	0.25	803	5.6	0.20	2,009	11.6	0.26	1,527	9.2	0.24	1,800	9.8	0.23
Washington	2,848	17.0	0.33	1,767	10.8	0.27	873	6.5	0.23	2,081	12.6	0.28	1,921	11.8	0.28	1,969	11.3	0.26
WA: Seattle/Puget Sound	1,996	17.9	0.41	1,189	10.8	0.33	614	6.6	0.28	1,435	12.9	0.35	1,355	12.7	0.36	1,295	11.2	0.32
West Virginia	998	16.2	0.53	838	13.8	0.50	468	9.9	0.48	763	13.0	0.49	640	11.0	0.46	660	10.2	0.41
Wisconsin	2,866	17.3	0.34	1,886	11.8	0.28	717	5.5	0.21	1,973	12.5	0.29	2,076	13.0	0.30	1,958	11.2	0.26
Wyoming	206	13.8	0.99	139	9.5	0.83	89	7.2	0.80	184	12.9	0.99	143	10.1	0.88	131	8.3	0.74

1. Rates are per 100,000 population and are age-adjusted by five-year age groups to the 2000 U.S. standard population based on single years of age.
2. See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.
3. Abbreviations: S.E. - Standard Error

\- Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for the U.S. combined.

N.B. In areas with small Latino populations, methods to indirectly identify Latinos (like NHIAv2) can overestimate the ethnicity-specific counts of cancer cases. Also, even small errors in Latino population estimates can affect the magnitude of the cancer rates. Data Item 190 used in place of NHIA for MA and RI, see technical notes.

NAACCR United States Combined Incidence 2010-2014, The Non-Hispanic Black Population

UNITED STATES

III-77 Combined Incidence – Non-Hispanic Black

Total Cancer Cases and Average Annual Age-adjusted Cancer Incidence Rates*
NAACCR United States Combined, 2010-2014, Non-Hispanic Black

	MALES (NON-HISPANIC BLACK)					FEMALES (NON-HISPANIC BLACK)				
SITES	*Count*	*Adj. Rate 2000 U.S.*	*S.E.*	*Adj. Rate 1996 Cdn*	*S.E.*	*Count*	*Adj. Rate 2000 U.S.*	*S.E.*	*Adj. Rate 1996 Cdn*	*S.E.*
All Sites (invasive & in situ bladder)	424,657	561.0	0.92	540.4	0.87	408,856	407.5	0.65	393.6	0.63
Oral Cavity & Pharynx	12,247	14.8	0.14	14.4	0.14	5,377	5.2	0.07	5.0	0.07
Lip	89	0.1	0.01	0.1	0.01	67	0.1	0.01	0.1	0.01
Tongue	2,806	3.4	0.07	3.3	0.07	1,135	1.1	0.03	1.0	0.03
Major salivary gland	939	1.2	0.04	1.1	0.04	1,084	1.1	0.03	1.1	0.03
Floor of mouth	733	0.9	0.03	0.9	0.03	298	0.3	0.02	0.3	0.02
Gum & other mouth	1,444	1.8	0.05	1.8	0.05	1,060	1.0	0.03	1.0	0.03
Nasopharynx	879	1.0	0.04	1.0	0.04	375	0.4	0.02	0.4	0.02
Oropharynx	989	1.2	0.04	1.2	0.04	310	0.3	0.02	0.3	0.02
Hypopharynx	1,487	1.8	0.05	1.8	0.05	298	0.3	0.02	0.3	0.02
Digestive System	92,303	123.4	0.44	117.8	0.41	80,982	81.7	0.29	77.5	0.28
Esophagus	5,478	7.2	0.10	6.9	0.10	2,339	2.3	0.05	2.2	0.05
Stomach	9,880	14.3	0.15	13.4	0.14	7,547	7.8	0.09	7.4	0.09
Small intestine	3,283	4.4	0.08	4.3	0.08	3,287	3.3	0.06	3.2	0.06
Colon excl. rectum	29,694	41.4	0.26	39.3	0.24	31,868	32.3	0.18	30.6	0.17
Rectum & rectosigmoid	11,695	15.0	0.15	14.4	0.14	9,561	9.4	0.10	9.0	0.09
Anus, anal canal & anorectum	1,660	2.0	0.05	2.0	0.05	1,792	1.7	0.04	1.7	0.04
Liver	14,363	16.2	0.14	15.9	0.14	4,569	4.3	0.07	4.1	0.06
Gallbladder	904	1.4	0.05	1.3	0.05	2,001	2.0	0.05	1.9	0.04
Pancreas	12,381	17.3	0.17	16.4	0.15	14,183	14.7	0.13	13.8	0.12
Respiratory System	69,677	97.7	0.40	93.2	0.37	51,750	52.4	0.24	50.1	0.22
Larynx	6,771	8.7	0.11	8.4	0.11	1,802	1.7	0.04	1.7	0.04
Lung & bronchus	62,072	88.0	0.38	83.7	0.35	49,347	50.1	0.23	47.8	0.22
Bones & Joints	803	0.9	0.03	0.9	0.03	739	0.7	0.03	0.7	0.03
Soft Tissue (incl. heart)	2,992	3.7	0.07	3.6	0.07	3,283	3.2	0.06	3.2	0.06
Skin (excl. basal & sq.)	1,612	2.1	0.06	2.1	0.05	1,817	1.8	0.04	1.8	0.04
Melanomas of the skin	795	1.1	0.04	1.1	0.04	955	1.0	0.03	0.9	0.03
Breast (excl. *in situ*)	1,368	1.9	0.06	1.8	0.05	127,722	125.5	0.36	122.4	0.35
Female Genital System	-	-	-	-	-	49,246	48.2	0.22	47.4	0.22
Cervix uteri	-	-	-	-	-	9,517	9.5	0.10	9.5	0.10
Uterus Corpus	-	-	-	-	-	24,720	23.8	0.15	23.5	0.15
Uterus, NOS[1]	-	-	-	-	-	1,778	1.8	0.04	1.7	0.04
Ovary	-	-	-	-	-	9,430	9.4	0.10	9.0	0.09
Male Genital System	149,683	189.5	0.52	185.2	0.50	-	-	-	-	-
Prostate	147,464	186.8	0.52	182.5	0.50	-	-	-	-	-
Testis	1,375	1.5	0.04	1.6	0.04	-	-	-	-	-
Penis	667	1.0	0.04	0.9	0.04	-	-	-	-	-
Urinary System	32,996	45.4	0.27	43.3	0.25	19,785	20.1	0.15	19.2	0.14
Bladder (incl. *in situ*)	12,822	19.8	0.19	18.4	0.17	6,327	6.7	0.09	6.2	0.08
Kidney & renal pelvis	19,663	24.8	0.19	24.2	0.18	12,994	12.9	0.12	12.5	0.11
Ureter	201	0.3	0.02	0.3	0.02	185	0.2	0.01	0.2	0.01
Eye & Orbit	240	0.3	0.02	0.3	0.02	237	0.2	0.02	0.2	0.01
Brain & Other Nervous System	4,125	4.9	0.08	4.8	0.08	3,646	3.6	0.06	3.5	0.06
Brain	3,798	4.5	0.08	4.4	0.07	3,251	3.2	0.06	3.1	0.06
Endocrine System	4,062	4.9	0.08	4.8	0.08	15,149	14.8	0.12	14.9	0.12
Thyroid gland	3,179	3.8	0.07	3.8	0.07	14,196	13.8	0.12	14.0	0.12
Lymphomas	16,366	20.4	0.17	19.9	0.16	14,569	14.5	0.12	14.2	0.12
Hodgkin Lymphoma	2,944	3.2	0.06	3.3	0.06	2,461	2.4	0.05	2.4	0.05
Non-Hodgkin Lymphoma	13,422	17.2	0.16	16.6	0.15	12,108	12.1	0.11	11.8	0.11
Multiple Myeloma	11,087	15.7	0.16	14.9	0.15	11,306	11.5	0.11	11.0	0.11
Leukemias	10,136	13.7	0.15	13.1	0.14	8,730	8.8	0.10	8.5	0.09
Acute lymphocytic	1,112	1.1	0.04	1.1	0.03	871	0.8	0.03	0.8	0.03
Chronic lymphocytic	3,132	4.6	0.09	4.3	0.08	2,284	2.4	0.05	2.2	0.05
Acute myeloid	3,057	4.2	0.08	4.0	0.08	3,096	3.1	0.06	3.0	0.06
Chronic myeloid	1,595	2.1	0.05	2.0	0.05	1,404	1.4	0.04	1.4	0.04
Other leukemias	1,240	1.8	0.05	1.7	0.05	1,075	1.1	0.03	1.0	0.03
Mesothelioma	553	0.9	0.04	0.8	0.04	225	0.2	0.02	0.2	0.02
Kaposi Sarcoma	1,457	1.6	0.04	1.7	0.05	159	0.2	0.01	0.2	0.01
Ill-defined & Unknown	12,950	19.1	0.18	17.9	0.16	14,134	14.7	0.13	13.7	0.12
Selected tumors (non-invasive)										
Brain & ONS[1] (*benign/borderline*)	6,373	8.3	0.11	8.0	0.10	14,831	14.9	0.12	14.4	0.12
Breast (*in situ*)	200	0.3	0.02	0.2	0.02	33,270	32.3	0.18	31.6	0.18

* Rates are per 100,000 population and are age-adjusted by five-year age groups to the 2000 U.S. standard population based on single years of age and to the 1996 Canadian population standard.

- Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for All Sites combined.

[1] Abbreviations: NOS - Not Otherwise Specified; ONS - Other Nervous System

N.B. In areas with small Latino populations, methods to indirectly identify Latinos (like NHIAv2) can overestimate the ethnicity-specific counts of cancer cases. Also, even small errors in Latino population estimates can affect the magnitude of the cancer rates. Data Item 190 used in place of NHIA for MA and RI, see technical notes.

Total Cancer Cases and Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates[1] for Commonly Diagnosed Cancers
Selected Areas in the NAACCR United States Combined[2], 2010-2014, Non-Hispanic Black, Males
{Page 1 of 4}

	All Sites			Prostate			Colon & Rectum			Lung & Bronchus			Kidney & Renal Pelvis			Liver & IBD[3]		
	Count	*Rate*	*S.E.*[3]	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
NAACCR U.S. Combined	**424,657**	**561.0**	**0.92**	**147,464**	**186.8**	**0.52**	**41,389**	**56.4**	**0.30**	**62,072**	**88.0**	**0.38**	**19,663**	**24.8**	**0.19**	**15,084**	**17.2**	**0.15**
Alabama	14,427	571.3	5.11	5,168	197.6	2.95	1,541	63.8	1.75	2,325	95.8	2.14	619	23.9	1.02	365	12.5	0.70
Alaska	215	479.0	40.65	87	167.2	21.69	21	45.5	12.17	27	82.6	18.87	8	12.9	5.22	9	16.6	6.49
Arizona	2,088	409.6	9.97	654	120.5	5.16	194	40.3	3.28	293	65.9	4.26	126	23.5	2.33	84	14.9	1.75
Arkansas	5,005	575.5	8.78	1,658	178.8	4.71	498	59.1	2.85	951	114.9	4.03	198	22.7	1.73	132	13.0	1.21
California	27,420	530.6	3.38	9,571	174.1	1.87	2,721	54.7	1.11	3,389	71.2	1.29	1,320	24.4	0.71	1,074	18.5	0.59
CA: Greater California	10,915	518.0	5.39	3,856	172.8	2.99	1,090	54.3	1.80	1,350	71.5	2.11	546	24.1	1.11	435	18.3	0.95
CA: Greater Bay Area	5,984	562.5	7.67	2,049	179.3	4.16	525	51.5	2.38	767	77.1	2.95	289	27.1	1.68	288	23.3	1.43
CA: Los Angeles	10,509	528.1	5.33	3,660	173.1	2.95	1,105	57.3	1.78	1,269	67.8	1.97	484	23.4	1.10	351	16.1	0.89
Colorado	1,998	471.3	11.75	689	157.1	6.69	198	48.9	3.87	220	60.1	4.46	104	23.2	2.43	98	18.9	2.08
Connecticut	3,683	551.2	9.71	1,312	187.0	5.47	338	52.9	3.10	458	75.6	3.78	169	21.8	1.75	110	14.9	1.50
Delaware	2,345	601.6	13.31	916	217.0	7.56	184	50.9	4.05	294	81.0	5.06	124	31.5	3.02	84	18.9	2.20
District of Columbia	4,590	609.0	9.27	1,453	188.5	5.10	449	61.9	3.01	679	90.1	3.56	198	26.1	1.91	254	30.1	1.94
Florida	28,210	491.1	3.10	10,261	173.6	1.80	2,857	51.3	1.02	3,655	67.5	1.18	1,070	17.8	0.57	787	12.0	0.45
Georgia	29,714	572.2	3.73	10,844	202.3	2.16	2,996	58.4	1.21	4,063	85.5	1.51	1,297	23.2	0.72	807	13.0	0.50
Hawaii	253	487.2	36.03	103	171.9	18.99	25	49.6	12.12	20	47.9	12.81	10	16.3	5.79	12	18.2	5.56
Idaho	68	404.8	61.13	25	132.8	31.33	-	-	-	-	-	-	-	-	-	8	59.5	28.79
Illinois	21,532	587.8	4.21	6,960	181.5	2.28	2,394	67.6	1.45	3,465	99.6	1.78	1,039	27.2	0.88	654	16.0	0.65
Indiana	5,685	523.7	7.42	1,613	141.6	3.75	575	54.1	2.41	980	97.7	3.33	321	26.9	1.59	223	17.1	1.20
Iowa	757	555.7	23.29	220	162.4	12.29	74	50.6	6.95	124	96.5	9.86	45	32.3	5.58	35	21.1	4.03
Kansas	1,892	583.9	14.53	627	189.3	8.13	190	62.8	4.94	317	101.9	6.19	83	23.9	2.80	65	16.8	2.21
Kentucky	4,054	610.3	10.40	1,150	166.1	5.28	413	63.2	3.39	729	119.0	4.78	235	31.9	2.21	142	18.1	1.62
Louisiana	17,930	629.1	5.05	5,883	200.8	2.80	1,905	67.0	1.65	2,968	110.1	2.18	795	27.6	1.04	675	19.9	0.81
Maine	111	381.8	40.72	35	124.3	23.08	7	18.7	7.29	7	30.7	12.73	-	-	-	7	26.7	11.82
Maryland	18,642	540.5	4.26	7,117	194.2	2.46	1,646	50.5	1.35	2,350	74.8	1.67	804	22.5	0.85	753	18.8	0.73
Massachusetts	4,459	525.5	8.45	1,720	194.7	5.00	375	45.3	2.52	506	67.8	3.24	182	20.8	1.65	186	19.6	1.54
Michigan	17,119	594.9	4.79	5,890	196.2	2.69	1,541	55.2	1.48	2,603	96.8	2.00	921	31.3	1.08	655	19.7	0.81
MI: Metropolitan Detroit	13,027	640.8	5.89	4,580	215.4	3.33	1,181	60.0	1.83	1,925	101.5	2.43	701	33.8	1.33	508	21.6	1.00
Mississippi	12,804	617.5	5.90	4,352	203.4	3.32	1,477	73.0	2.06	2,264	114.6	2.61	600	28.7	1.26	229	9.4	0.67

Total Cancer Cases and Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates[1] for Commonly Diagnosed Cancers
Selected Areas in the NAACCR United States Combined[2], 2010-2014, Non-Hispanic Black, Males
{Page 2 of 4}

	All Sites			Prostate			Colon & Rectum			Lung & Bronchus			Kidney & Renal Pelvis			Liver & IBD[3]		
	Count	*Rate*	*S.E.*[3]	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
Missouri	7,583	571.9	7.01	2,228	159.2	3.57	745	60.3	2.36	1,339	106.0	3.08	360	25.9	1.45	366	23.6	1.30
Montana	50	588.5	96.16	17	190.9	49.43	-	-	-	9	129.0	47.41	-	-	-	-	-	-
Nebraska	902	634.5	23.26	308	209.4	12.99	106	80.2	8.66	125	99.2	9.63	53	36.2	5.49	35	21.8	3.94
New Hampshire	132	433.9	41.92	51	164.5	25.39	15	48.3	13.53	12	39.6	12.22	-	-	-	6	23.5	10.48
New Jersey	13,041	563.9	5.21	5,024	210.9	3.12	1,217	53.9	1.63	1,581	74.5	1.98	597	24.4	1.04	390	14.8	0.79
New York	34,705	582.2	3.28	13,992	226.1	2.00	3,034	53.4	1.02	3,888	69.1	1.17	1,420	22.9	0.63	1,246	18.8	0.56
North Carolina	24,593	575.7	3.95	8,548	191.0	2.21	2,226	53.4	1.22	4,019	100.2	1.70	1,116	25.4	0.81	734	14.6	0.57
North Dakota	48	441.7	88.78	14	120.7	42.10	6	59.2	32.38	7	100.7	47.14	-	-	-	-	-	-
Ohio	15,777	538.1	4.49	5,258	171.8	2.48	1,458	51.1	1.41	2,658	96.1	1.95	769	25.2	0.94	607	17.8	0.75
Oklahoma	3,159	597.7	11.61	1,073	200.3	6.66	296	54.8	3.49	522	103.5	4.91	144	27.2	2.45	99	15.8	1.73
Oregon	802	517.0	20.53	260	162.9	11.32	74	51.8	6.67	114	81.5	8.63	42	23.8	4.11	50	30.7	4.86
Pennsylvania	16,420	605.0	4.97	5,152	181.5	2.65	1,457	55.4	1.53	2,476	99.1	2.09	894	31.0	1.08	900	28.9	1.01
Rhode Island	499	467.4	22.79	148	126.9	11.27	38	40.5	7.09	68	71.9	9.46	20	18.5	4.56	22	14.6	3.26
South Carolina	15,438	566.6	4.90	5,428	189.4	2.75	1,461	55.1	1.55	2,368	90.8	2.01	639	22.8	0.96	387	12.4	0.67
South Dakota	65	275.4	40.23	10	33.7	10.92	7	29.7	11.68	8	45.0	21.99	6	22.1	9.29	-	-	-
Tennessee	11,207	568.3	5.91	3,734	183.0	3.28	1,091	58.2	1.94	1,897	103.8	2.63	503	24.6	1.20	382	16.1	0.89
Texas	29,597	546.5	3.51	8,950	160.1	1.86	3,209	59.6	1.17	4,540	92.8	1.52	1,496	25.9	0.73	1,498	22.7	0.63
Utah	238	557.0	44.31	87	209.8	26.75	29	66.1	15.26	23	60.8	15.08	14	32.0	10.24	16	33.7	9.67
Vermont	33	277.3	54.76	14	115.1	35.02	-	-	-	-	-	-	-	-	-	-	-	-
Virginia	17,857	543.6	4.34	6,409	182.5	2.42	1,595	50.1	1.34	2,671	88.9	1.84	900	26.0	0.91	542	13.9	0.63
Washington	2,736	533.6	11.65	926	169.2	6.25	224	44.2	3.35	314	74.6	4.73	161	27.8	2.44	138	23.9	2.28
WA: Seattle/Puget Sound	2,487	547.8	12.54	854	174.2	6.68	197	44.6	3.61	284	77.6	5.16	142	27.7	2.60	132	26.4	2.57
West Virginia	854	526.8	19.70	281	160.4	10.33	91	56.0	6.44	131	92.1	8.79	46	25.7	4.02	38	18.9	3.27
Wisconsin	3,889	682.1	12.25	1,235	209.9	6.63	375	67.8	3.94	605	119.0	5.39	192	30.1	2.40	171	24.9	2.02
Wyoming	31	186.4	34.82	9	51.9	17.96	-	-	-	-	-	-	-	-	-	-	-	-

1. Rates are per 100,000 population and are age-adjusted by five-year age groups to the 2000 U.S. standard population based on single years of age.
2. See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.
3. Abbreviations: IBD - Intrahepatic Bile Duct; S.E. - Standard Error

\- Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for the U.S. combined.

N.B. In areas with small Latino populations, methods to indirectly identify Latinos (like NHIAv2) can overestimate the ethnicity-specific counts of cancer cases. Also, even small errors in Latino population estimates can affect the magnitude of the cancer rates. Data Item 190 used in place of NHIA for MA and RI, see technical notes.

Total Cancer Cases and Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates[1] for Commonly Diagnosed Cancers
Selected Areas in the NAACCR United States Combined[2], 2010-2014, Non-Hispanic Black, Males
{Page 3 of 4}

	Non-Hodgkin Lymphoma			Leukemia			Bladder			Stomach			Oral Cavity & Pharynx			Pancreas		
	Count	*Rate*	*S.E.*[3]	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
NAACCR U.S. Combined	**13,422**	**17.2**	**0.16**	**10,136**	**13.7**	**0.15**	**12,822**	**19.8**	**0.19**	**9,880**	**14.3**	**0.15**	**12,247**	**14.8**	**0.14**	**12,381**	**17.3**	**0.17**
Alabama	352	13.9	0.79	291	12.0	0.76	424	18.9	0.98	343	15.1	0.88	439	15.9	0.80	409	16.8	0.90
Alaska	9	25.5	10.18	6	6.6	2.75	10	34.7	12.52	-	-	-	-	-	-	-	-	-
Arizona	79	13.7	1.72	35	6.2	1.15	69	17.4	2.35	42	9.6	1.67	40	6.9	1.18	66	13.5	1.81
Arkansas	139	15.9	1.45	111	14.0	1.45	141	19.9	1.80	115	14.9	1.50	145	14.5	1.27	141	17.0	1.56
California	943	17.7	0.61	744	14.9	0.57	1,031	23.2	0.76	609	12.8	0.55	743	13.5	0.52	821	16.9	0.62
CA: Greater California	367	16.2	0.92	315	14.9	0.91	399	23.0	1.25	219	11.2	0.82	275	12.0	0.78	316	16.3	1.00
CA: Greater Bay Area	231	20.9	1.44	156	16.0	1.34	221	23.7	1.68	130	13.2	1.22	167	15.1	1.22	193	19.0	1.44
CA: Los Angeles	345	17.6	0.98	273	14.2	0.89	411	23.1	1.18	260	14.1	0.91	301	14.4	0.86	312	16.4	0.96
Colorado	78	16.0	1.94	54	13.8	2.16	73	19.0	2.49	55	14.2	2.14	48	10.0	1.53	54	13.1	1.96
Connecticut	125	18.2	1.74	83	11.7	1.37	126	22.2	2.12	100	16.9	1.79	98	13.3	1.43	120	18.9	1.86
Delaware	78	20.3	2.46	54	14.2	2.05	84	25.6	3.03	55	16.0	2.33	55	12.5	1.77	57	15.1	2.16
District of Columbia	166	22.3	1.78	81	11.8	1.34	145	20.0	1.71	106	15.2	1.52	151	19.0	1.60	165	21.8	1.75
Florida	1,000	16.6	0.56	806	13.6	0.51	727	15.1	0.59	749	14.4	0.56	797	12.9	0.48	740	13.8	0.54
Georgia	1,021	18.0	0.63	716	15.0	0.64	795	19.0	0.76	661	14.4	0.64	811	13.7	0.53	798	16.5	0.66
Hawaii	9	19.9	7.73	8	16.2	6.85	14	35.6	10.59	-	-	-	7	16.5	7.30	9	17.7	6.82
Idaho	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Illinois	621	16.5	0.69	544	15.4	0.70	661	20.2	0.83	531	15.8	0.72	749	19.3	0.74	629	18.0	0.76
Indiana	178	16.4	1.33	137	12.0	1.12	181	19.6	1.54	144	14.9	1.32	167	14.3	1.18	191	19.5	1.50
Iowa	28	18.6	4.09	25	18.0	4.24	25	26.2	5.96	9	7.2	2.82	20	11.5	2.74	26	21.8	4.71
Kansas	39	11.1	1.90	48	15.8	2.47	50	17.2	2.64	29	8.9	1.81	55	16.3	2.37	49	17.0	2.63
Kentucky	141	21.2	1.95	107	17.0	1.79	140	24.5	2.24	81	12.6	1.52	138	18.7	1.72	117	17.1	1.72
Louisiana	552	18.9	0.86	365	13.2	0.74	477	19.4	0.96	454	17.1	0.86	550	17.4	0.79	484	17.9	0.88
Maine	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Maryland	530	15.2	0.71	447	13.8	0.70	650	22.3	0.94	369	11.9	0.67	464	12.6	0.62	569	17.5	0.80
Massachusetts	170	18.3	1.51	103	11.2	1.20	119	16.2	1.60	125	16.3	1.58	125	14.0	1.34	122	16.2	1.57
Michigan	535	18.7	0.85	372	13.4	0.74	554	21.5	0.96	358	13.3	0.74	523	16.4	0.74	526	19.1	0.88
MI: Metropolitan Detroit	394	19.4	1.02	296	15.4	0.94	417	22.9	1.17	265	13.7	0.88	409	18.5	0.95	410	20.7	1.07
Mississippi	313	14.6	0.89	306	15.0	0.93	305	16.6	1.03	296	14.9	0.94	412	17.6	0.92	355	18.5	1.07

Total Cancer Cases and Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates[1] for Commonly Diagnosed Cancers
Selected Areas in the NAACCR United States Combined[2], 2010-2014, Non-Hispanic Black, Males
{Page 4 of 4}

	Non-Hodgkin Lymphoma			Leukemia			Bladder			Stomach			Oral Cavity & Pharynx			Pancreas		
	Count	*Rate*	*S.E.*[3]	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
Missouri	248	17.6	1.18	192	15.8	1.22	206	18.2	1.35	173	13.5	1.10	235	16.2	1.12	263	21.4	1.41
Montana	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Nebraska	23	16.8	3.95	20	14.6	3.75	21	14.7	3.49	18	12.8	3.27	32	17.5	3.19	24	19.3	4.40
New Hampshire	-	-	-	-	-	-	7	38.0	15.27	-	-	-	-	-	-	-	-	-
New Jersey	401	16.6	0.87	322	14.4	0.85	440	22.0	1.11	284	13.1	0.83	323	13.0	0.76	399	17.4	0.92
New York	1,186	19.2	0.58	817	14.0	0.51	1,010	19.5	0.65	858	15.8	0.57	811	12.7	0.46	945	16.4	0.56
North Carolina	665	15.5	0.64	527	12.6	0.59	758	20.6	0.81	512	12.7	0.61	844	18.0	0.66	717	17.5	0.71
North Dakota	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Ohio	498	16.9	0.79	325	11.2	0.66	549	21.0	0.94	364	13.3	0.73	430	13.5	0.68	451	16.4	0.81
Oklahoma	83	15.7	1.90	78	15.5	1.96	97	19.8	2.20	75	15.1	1.91	100	16.5	1.74	94	17.7	1.98
Oregon	25	15.6	3.57	20	14.4	3.58	25	20.8	4.64	19	12.5	3.18	17	10.2	2.58	18	12.3	3.25
Pennsylvania	586	21.0	0.91	352	12.8	0.72	593	24.8	1.07	337	13.4	0.77	489	17.8	0.84	489	18.7	0.89
Rhode Island	21	20.6	4.80	15	15.2	4.35	17	16.2	4.25	17	16.9	4.48	17	15.9	4.15	19	19.4	4.80
South Carolina	390	14.6	0.78	338	13.1	0.77	371	16.1	0.90	427	16.9	0.88	532	17.5	0.80	519	20.3	0.96
South Dakota	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee	340	16.0	0.94	244	12.8	0.90	258	15.8	1.08	242	13.2	0.93	310	13.5	0.82	326	17.5	1.07
Texas	998	17.0	0.60	816	15.1	0.59	825	18.9	0.72	720	14.8	0.61	780	13.1	0.51	854	16.3	0.62
Utah	6	14.9	7.52	12	19.2	7.59	-	-	-	-	-	-	-	-	-	-	-	-
Vermont	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Virginia	564	17.5	0.78	375	11.9	0.66	594	21.1	0.93	398	13.9	0.74	538	15.0	0.69	582	19.3	0.86
Washington	117	20.2	2.12	82	17.3	2.21	95	21.6	2.54	61	14.7	2.11	80	13.5	1.70	79	17.3	2.19
WA: Seattle/Puget Sound	103	20.5	2.31	79	19.3	2.52	84	20.5	2.57	51	13.8	2.18	69	12.2	1.63	70	17.2	2.33
West Virginia	23	14.2	3.22	29	19.4	3.95	28	20.4	4.21	21	13.1	3.14	26	17.6	3.76	21	13.3	3.14
Wisconsin	117	20.2	2.08	113	20.7	2.23	110	22.6	2.39	98	18.3	2.06	126	19.5	1.91	120	22.2	2.29
Wyoming	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-

1. Rates are per 100,000 population and are age-adjusted by five-year age groups to the 2000 U.S. standard population based on single years of age.
2. See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.
3. Abbreviations: S.E. - Standard Error

\- Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for the U.S. combined.

N.B. In areas with small Latino populations, methods to indirectly identify Latinos (like NHIAv2) can overestimate the ethnicity-specific counts of cancer cases. Also, even small errors in Latino population estimates can affect the magnitude of the cancer rates. Data Item 190 used in place of NHIA for MA and RI, see technical notes.

Total Cancer Cases and Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates[1] for Commonly Diagnosed Cancers
Selected Areas in the NAACCR United States Combined[2], 2010-2014, Non-Hispanic Black, Females
{Page 1 of 4}

	All Sites			Breast			Colon & Rectum			Thyroid			Uterus Corpus & Uterus NOS[3]			Lung & Bronchus		
	Count	*Rate*	*S.E.[3]*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
NAACCR U.S. Combined	**408,856**	**407.5**	**0.65**	**127,722**	**125.5**	**0.36**	**41,429**	**41.7**	**0.21**	**14,196**	**13.8**	**0.12**	**26,498**	**25.6**	**0.16**	**49,347**	**50.1**	**0.23**
Alabama	12,919	379.3	3.41	4,306	124.9	1.95	1,504	44.5	1.17	339	9.9	0.55	750	21.7	0.81	1,312	38.6	1.09
Alaska	153	341.1	29.20	60	133.7	18.13	12	23.6	7.18	8	15.5	5.81	-	-	-	11	28.5	8.78
Arizona	1,944	357.1	8.40	607	107.8	4.51	189	36.0	2.72	114	18.9	1.80	132	23.2	2.09	229	44.5	3.05
Arkansas	4,225	376.3	5.94	1,300	114.2	3.25	498	44.6	2.05	113	9.7	0.93	239	20.7	1.38	532	48.3	2.15
California	25,327	407.4	2.61	8,101	129.0	1.46	2,634	42.7	0.85	776	12.7	0.47	1,678	25.7	0.64	3,066	49.8	0.92
CA: Greater California	9,284	396.9	4.24	3,012	124.9	2.34	942	41.8	1.40	329	13.1	0.74	579	23.4	1.00	1,099	49.4	1.54
CA: Greater Bay Area	5,316	422.5	5.93	1,660	131.3	3.30	556	43.9	1.91	105	8.7	0.87	352	27.0	1.47	697	55.4	2.15
CA: Los Angeles	10,724	410.5	4.02	3,429	131.4	2.28	1,136	43.1	1.29	342	14.1	0.78	747	27.4	1.02	1,269	47.7	1.35
Colorado	1,616	361.1	9.22	545	119.1	5.22	147	34.6	2.92	80	16.2	1.84	87	19.2	2.12	183	44.1	3.35
Connecticut	3,729	406.5	6.78	1,158	122.9	3.67	339	37.5	2.08	173	18.4	1.41	266	27.6	1.72	395	44.4	2.28
Delaware	2,183	426.8	9.38	686	130.6	5.11	181	35.9	2.74	94	17.6	1.85	157	29.9	2.45	276	57.3	3.55
District of Columbia	4,880	468.4	6.91	1,438	141.6	3.85	508	47.9	2.19	170	18.1	1.44	385	35.2	1.84	622	57.5	2.35
Florida	26,659	360.4	2.25	8,145	108.3	1.22	2,673	36.8	0.73	910	11.9	0.40	1,953	25.6	0.59	2,595	35.9	0.72
Georgia	28,852	394.7	2.41	9,577	126.3	1.33	3,036	42.6	0.80	926	11.5	0.39	1,846	24.6	0.59	2,862	41.3	0.80
Hawaii	119	330.1	33.58	49	120.5	18.63	6	21.1	9.83	-	-	-	6	17.4	7.69	9	38.0	13.17
Idaho	40	359.4	62.51	12	111.2	35.98	-	-	-	-	-	-	-	-	-	-	-	-
Illinois	22,347	440.3	2.99	6,725	131.8	1.63	2,469	48.9	1.00	650	12.8	0.51	1,457	27.5	0.73	3,215	63.5	1.14
Indiana	6,119	424.6	5.55	1,916	130.6	3.05	619	43.8	1.80	152	10.0	0.83	341	23.1	1.28	875	62.3	2.15
Iowa	665	448.9	18.27	162	105.1	8.60	65	47.3	6.11	17	9.5	2.37	45	30.1	4.65	102	74.4	7.68
Kansas	1,642	432.7	10.88	482	126.2	5.85	152	40.5	3.34	77	19.6	2.27	95	25.1	2.62	235	63.7	4.22
Kentucky	3,805	450.1	7.48	1,090	127.0	3.94	421	49.6	2.48	133	15.4	1.36	200	23.5	1.70	640	78.1	3.17
Louisiana	15,700	424.0	3.47	4,964	132.5	1.93	1,852	50.5	1.20	526	14.0	0.62	840	21.9	0.78	1,848	50.4	1.20
Maine	73	309.6	38.77	15	61.7	16.51	-	-	-	-	-	-	6	26.3	11.50	10	48.9	16.56
Maryland	19,352	404.4	2.99	6,513	132.4	1.68	1,763	37.5	0.92	760	15.2	0.56	1,310	26.4	0.75	2,213	47.4	1.04
Massachusetts	4,288	389.3	6.05	1,281	114.7	3.26	401	36.3	1.85	268	22.9	1.42	276	24.5	1.50	481	45.9	2.13
Michigan	16,294	430.7	3.45	4,828	127.1	1.87	1,610	42.6	1.08	529	14.1	0.63	1,058	26.8	0.84	2,395	63.4	1.33
MI: Metropolitan Detroit	12,547	447.8	4.09	3,722	132.4	2.22	1,275	45.6	1.30	433	15.7	0.77	794	27.1	0.99	1,822	64.6	1.55
Mississippi	11,176	406.6	3.94	3,396	121.0	2.12	1,434	52.6	1.42	350	12.2	0.66	639	22.8	0.93	1,271	47.8	1.37

Total Cancer Cases and Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates[1] for Commonly Diagnosed Cancers
Selected Areas in the NAACCR United States Combined[2], 2010-2014, Non-Hispanic Black, Females
{Page 2 of 4}

	All Sites			Breast			Colon & Rectum			Thyroid			Uterus Corpus & Uterus NOS[3]			Lung & Bronchus		
	Count	*Rate*	*S.E.*[3]	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
Missouri	8,048	446.7	5.07	2,431	133.4	2.75	785	44.0	1.60	227	12.3	0.83	497	26.9	1.23	1,214	68.2	2.00
Montana	17	309.8	80.22	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Nebraska	728	434.6	16.61	213	127.8	8.99	70	43.5	5.34	31	15.9	2.93	34	19.1	3.37	107	65.4	6.49
New Hampshire	64	228.6	29.86	21	77.9	17.61	6	22.6	9.65	-	-	-	6	21.9	9.47	7	26.2	10.05
New Jersey	13,628	417.7	3.63	4,176	125.9	1.97	1,394	43.3	1.18	459	14.1	0.66	949	28.3	0.93	1,595	49.3	1.25
New York	34,561	404.0	2.20	10,215	119.5	1.20	3,302	38.6	0.68	1,356	16.0	0.44	2,851	32.1	0.61	3,722	43.1	0.72
North Carolina	23,728	410.8	2.72	7,837	134.1	1.55	2,209	38.4	0.83	817	14.1	0.50	1,519	25.5	0.67	2,725	47.6	0.93
North Dakota	27	344.7	85.69	6	55.5	22.84	-	-	-	-	-	-	-	-	-	-	-	-
Ohio	15,483	405.8	3.31	4,740	123.8	1.83	1,445	37.9	1.01	602	16.3	0.68	919	23.3	0.78	2,424	63.8	1.31
Oklahoma	2,720	406.4	8.01	839	122.9	4.35	271	40.2	2.51	84	11.8	1.32	164	23.6	1.89	372	57.8	3.09
Oregon	608	409.8	17.12	192	127.1	9.40	54	38.8	5.45	28	17.4	3.35	37	24.0	4.05	77	55.4	6.51
Pennsylvania	17,364	470.9	3.63	4,810	130.8	1.92	1,581	42.6	1.09	819	22.7	0.81	1,121	29.4	0.89	2,573	70.0	1.40
Rhode Island	504	370.9	16.85	156	113.4	9.25	41	30.7	4.88	36	24.4	4.13	32	23.9	4.32	66	49.6	6.22
South Carolina	14,121	391.2	3.37	4,573	125.7	1.91	1,364	37.2	1.03	461	13.0	0.62	992	26.5	0.86	1,536	42.3	1.11
South Dakota	39	216.0	35.63	16	93.4	23.62	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee	10,947	405.6	3.99	3,506	126.3	2.19	1,123	42.8	1.31	292	10.5	0.62	617	22.0	0.92	1,357	52.3	1.47
Texas	28,925	403.9	2.45	8,919	120.3	1.31	3,021	42.3	0.80	1,004	12.9	0.42	1,682	22.8	0.58	3,336	49.4	0.89
Utah	128	348.1	33.30	34	90.3	16.21	16	50.6	13.28	7	14.0	5.81	-	-	-	13	39.8	11.72
Vermont	23	253.0	58.19	8	75.1	28.27	-	-	-	-	-	-	-	-	-	-	-	-
Virginia	16,923	397.8	3.12	5,807	134.3	1.80	1,696	40.2	1.00	536	12.5	0.55	950	21.8	0.72	2,005	48.0	1.09
Washington	1,978	392.0	9.31	656	125.7	5.15	152	30.0	2.57	96	15.9	1.69	98	19.5	2.07	216	48.1	3.44
WA: Seattle/Puget Sound	1,837	407.6	10.06	612	130.3	5.54	141	31.5	2.80	91	16.9	1.84	92	20.6	2.27	203	51.5	3.80
West Virginia	653	383.1	15.38	200	120.1	8.74	64	37.5	4.81	25	15.6	3.21	36	19.3	3.32	90	51.3	5.51
Wisconsin	3,502	484.6	8.55	1,001	133.5	4.37	308	43.4	2.58	127	15.4	1.42	205	28.0	2.03	517	76.9	3.53
Wyoming	30	189.0	35.67	7	48.8	19.06	-	-	-	-	-	-	-	-	-	-	-	-

1. Rates are per 100,000 population and are age-adjusted by five-year age groups to the 2000 U.S. standard population based on single years of age.
2. See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.
3. Abbreviations: NOS - Not Otherwise Specified; S.E. - Standard Error

\- Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for the U.S. combined.

N.B. In areas with small Latino populations, methods to indirectly identify Latinos (like NHIAv2) can overestimate the ethnicity-specific counts of cancer cases. Also, even small errors in Latino population estimates can affect the magnitude of the cancer rates. Data Item 190 used in place of NHIA for MA and RI, see technical notes.

Total Cancer Cases and Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates[1] for Commonly Diagnosed Cancers
Selected Areas in the NAACCR United States Combined[2], 2010-2014, Non-Hispanic Black, Females
{Page 3 of 4}

	Non-Hodgkin Lymphoma			Kidney & Renal Pelvis			Cervix			Ovary			Leukemia			Pancreas		
	Count	*Rate*	*S.E.*[3]	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
NAACCR U.S. Combined	**12,108**	**12.1**	**0.11**	**12,994**	**12.9**	**0.12**	**9,517**	**9.5**	**0.10**	**9,430**	**9.4**	**0.10**	**8,730**	**8.8**	**0.10**	**14,183**	**14.7**	**0.13**
Alabama	332	9.8	0.55	398	11.4	0.58	322	9.9	0.56	341	10.0	0.56	252	7.5	0.48	496	15.0	0.69
Alaska	7	19.9	7.91	6	12.5	5.39	-	-	-	-	-	-	-	-	-	-	-	-
Arizona	51	9.5	1.38	66	12.0	1.53	43	7.2	1.11	44	8.4	1.32	43	8.0	1.29	67	13.8	1.75
Arkansas	105	9.5	0.95	147	13.3	1.13	135	11.8	1.04	103	9.3	0.94	75	6.7	0.79	182	17.2	1.31
California	784	12.9	0.47	774	12.3	0.45	465	7.6	0.36	605	9.7	0.40	520	8.6	0.38	857	14.0	0.49
CA: Greater California	282	12.4	0.76	319	13.6	0.78	190	7.6	0.57	204	8.6	0.62	186	8.0	0.60	301	13.5	0.80
CA: Greater Bay Area	171	13.8	1.08	160	12.7	1.02	65	5.5	0.69	137	11.1	0.97	103	8.5	0.86	200	16.1	1.16
CA: Los Angeles	331	13.0	0.72	295	11.1	0.66	210	8.7	0.62	264	10.1	0.63	231	9.2	0.62	355	13.5	0.72
Colorado	54	11.9	1.66	50	11.0	1.60	29	5.9	1.12	35	8.2	1.43	39	8.3	1.37	56	13.8	1.89
Connecticut	105	11.6	1.16	94	10.1	1.06	76	8.6	1.00	89	9.8	1.06	82	9.2	1.03	108	12.5	1.23
Delaware	60	11.5	1.52	67	13.4	1.67	51	9.7	1.38	45	8.8	1.34	45	9.0	1.37	80	16.2	1.86
District of Columbia	135	13.4	1.19	133	12.5	1.11	108	11.8	1.18	125	12.2	1.13	76	7.1	0.84	175	16.1	1.25
Florida	896	12.0	0.41	695	9.4	0.36	811	10.8	0.39	696	9.3	0.36	673	9.1	0.36	888	12.7	0.43
Georgia	896	12.2	0.42	874	11.9	0.42	656	8.5	0.34	734	10.0	0.38	667	9.5	0.38	938	14.1	0.48
Hawaii	-	-	-	-	-	-	-	-	-	7	17.1	6.53	-	-	-	-	-	-
Idaho	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Illinois	583	11.7	0.49	665	13.0	0.51	540	10.8	0.47	501	9.9	0.45	465	9.3	0.44	743	14.9	0.56
Indiana	195	13.7	1.00	182	12.6	0.96	141	9.4	0.80	121	8.3	0.77	118	8.2	0.78	217	15.6	1.08
Iowa	23	13.5	2.92	37	25.8	4.40	13	7.1	2.04	13	9.1	2.67	16	11.0	2.92	24	18.4	3.92
Kansas	49	12.6	1.84	52	14.0	1.98	26	7.2	1.43	22	5.9	1.28	43	10.9	1.71	43	11.5	1.78
Kentucky	94	10.6	1.12	145	17.0	1.45	70	8.1	0.98	67	8.1	1.01	79	9.4	1.09	127	15.7	1.43
Louisiana	440	12.1	0.59	578	15.5	0.66	412	11.6	0.58	329	8.8	0.50	317	8.7	0.50	518	14.4	0.65
Maine	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Maryland	595	12.4	0.52	577	12.1	0.52	362	7.6	0.41	451	9.3	0.45	392	8.4	0.44	647	14.2	0.58
Massachusetts	144	12.9	1.10	122	11.0	1.01	91	8.1	0.86	91	8.0	0.85	84	7.5	0.83	133	12.6	1.11
Michigan	459	12.2	0.58	531	14.1	0.62	299	8.2	0.48	367	9.6	0.51	334	8.9	0.49	623	16.6	0.68
MI: Metropolitan Detroit	349	12.7	0.69	406	14.5	0.74	241	9.0	0.59	286	10.1	0.61	274	10.0	0.62	473	16.8	0.79
Mississippi	296	10.9	0.65	458	16.8	0.81	334	12.0	0.67	238	8.7	0.58	236	8.6	0.57	367	14.0	0.75

Total Cancer Cases and Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates[1] for Commonly Diagnosed Cancers
Selected Areas in the NAACCR United States Combined[2], 2010-2014, Non-Hispanic Black, Females
{Page 4 of 4}

	Non-Hodgkin Lymphoma			Kidney & Renal Pelvis			Cervix			Ovary			Leukemia			Pancreas		
	Count	*Rate*	*S.E.*[3]	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
Missouri	224	12.7	0.86	288	15.9	0.95	181	10.0	0.76	152	8.4	0.69	203	11.4	0.81	282	16.0	0.97
Montana	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Nebraska	24	15.0	3.14	25	15.6	3.22	19	9.8	2.28	8	4.8	1.75	16	7.7	2.02	27	17.3	3.41
New Hampshire	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey	406	12.4	0.62	381	11.7	0.61	354	10.7	0.58	341	10.5	0.57	292	9.4	0.56	482	15.1	0.70
New York	1,110	13.0	0.40	871	10.3	0.35	897	10.9	0.37	873	10.2	0.35	737	8.8	0.33	1,221	14.4	0.42
North Carolina	692	12.1	0.47	849	14.6	0.51	469	8.2	0.38	530	9.3	0.41	486	8.7	0.40	842	15.0	0.53
North Dakota	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Ohio	423	11.1	0.55	491	12.9	0.59	269	7.2	0.45	300	7.9	0.47	301	7.8	0.46	546	14.5	0.63
Oklahoma	79	12.3	1.42	101	14.9	1.52	50	7.5	1.09	70	10.7	1.32	66	10.1	1.28	96	14.3	1.51
Oregon	14	9.1	2.57	29	19.2	3.67	9	5.3	1.79	17	11.4	2.85	16	10.1	2.63	19	13.2	3.10
Pennsylvania	525	14.1	0.63	554	15.0	0.65	386	10.8	0.56	353	9.6	0.52	355	9.7	0.52	624	17.0	0.69
Rhode Island	17	13.6	3.38	12	9.0	2.65	7	5.0	1.91	9	6.3	2.17	13	8.5	2.40	14	10.9	2.97
South Carolina	357	10.1	0.55	449	12.4	0.60	306	8.9	0.52	321	9.0	0.51	293	8.3	0.50	522	14.8	0.66
South Dakota	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee	315	11.6	0.67	418	15.4	0.78	306	11.1	0.65	256	9.3	0.60	259	9.8	0.63	362	14.3	0.77
Texas	883	12.3	0.43	1,053	14.8	0.47	797	10.6	0.38	670	9.4	0.37	646	9.2	0.37	1,007	15.2	0.50
Utah	7	16.0	6.97	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Vermont	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Virginia	507	12.0	0.54	598	13.9	0.58	328	7.9	0.44	369	8.7	0.46	314	7.6	0.44	627	15.0	0.61
Washington	84	16.5	1.91	62	11.9	1.58	29	5.7	1.12	41	7.9	1.34	47	9.6	1.49	69	15.9	2.01
WA: Seattle/Puget Sound	72	15.7	1.97	56	12.3	1.72	25	5.6	1.20	40	8.4	1.44	45	10.3	1.64	64	16.3	2.15
West Virginia	20	11.6	2.65	22	13.2	2.90	8	5.0	1.79	21	12.0	2.70	14	7.6	2.08	26	14.7	2.95
Wisconsin	104	14.8	1.52	125	17.8	1.66	102	12.7	1.29	62	7.5	1.00	101	14.2	1.50	112	16.8	1.66
Wyoming	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-

1. Rates are per 100,000 population and are age-adjusted by five-year age groups to the 2000 U.S. standard population based on single years of age.
2. See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.
3. Abbreviations: S.E. - Standard Error

\- Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for the U.S. combined.

N.B. In areas with small Latino populations, methods to indirectly identify Latinos (like NHIAv2) can overestimate the ethnicity-specific counts of cancer cases. Also, even small errors in Latino population estimates can affect the magnitude of the cancer rates. Data Item 190 used in place of NHIA for MA and RI, see technical notes.

Race-Specific Population Indicators (2010-2014)

Combined Population:	
All Races	65,999,047
White	-
Black	-
American Indian/Alaskan Native	-
Asian/Pacific Islander	-

Percentage of Canada Population Reflected in the Combined Cancer Statistics:	
All Races	38.0%
White	-
Black	-
American Indian/Alaskan Native	-
Asian/Pacific Islander	-

Number of Cases in Combined Cancer Statistics:	
All Races	341,437
White	-
Black	-
American Indian/Alaskan Native	-
Asian/Pacific Islander	-

Race-Specific Population Counts by Year

Race	2010	2011	2012	2013	2014
All Races	12,872,897	13,001,982	13,180,172	13,372,001	13,571,995

Data Quality Indicators (2010-2014)

Indicators (%)	2010	2011	2012	2013	2014
Missing Race	na	na	na	na	na
Missing Sex	0.0	0.0	0.0	0.0	0.0
Missing County	0.0	0.0	0.0	0.0	0.0
Missing Age	0.0	0.0	0.0	0.0	0.0
Death Certificate Only Cases	0.7	0.5	0.6	0.4	0.3

Site-Specific Microscopic Confirmation (2010-2014)

Cancer Site	% Micro. Confirm	Cancer Site	% Micro. Confirm
All Sites	91.6	Prostate	94.6
Oral Cavity & Pharynx	97.8	Testis	99.3
Esophagus	95.8	Bladder (incl. *in situ*)	96.9
Stomach	95.2	Kidney & Renal Pelvis	86.6
Colon & Rectum	95.3	Brain & Other Nervous System	83.7
Liver	46.5	Thyroid Gland	99.3
Pancreas	65.8	Hodgkin Lymphoma	99.6
Lung & Bronchus	82.0	Non-Hodgkin Lymphoma	97.2
Melanomas of the Skin	99.5	Multiple Myeloma	91.2
Female Breast (excl. *in situ*)	98.9	Leukemias	96.2
Cervix uteri	98.8	Brain & ONS[1] *benign/borderline*	59.9
Uterus Corpus & Uterus NOS[1]	98.9	Breast *in situ*	100.0
Ovary	90.4		

[1] Abbreviations: NOS - Not Otherwise Specified; ONS - Other Nervous System

NAACCR Canadian Combined Incidence 2010-2014, All Races Combined

CANADA

III-89 Combined Incidence – All Races

Total Cancer Cases and Average Annual Age-adjusted Cancer Incidence Rates*
NAACCR Canadian Combined, 2010-2014, All Races

	MALES (ALL RACES)					FEMALES (ALL RACES)				
SITES	*Count*	*Adj. Rate 2000 U.S.*	*S.E.*	*Adj. Rate 1996 Cdn*	*S.E.*	*Count*	*Adj. Rate 2000 U.S.*	*S.E.*	*Adj. Rate 1996 Cdn*	*S.E.*
All Sites (invasive & in situ bladder)	176,555	488.9	1.19	464.7	1.12	164,882	410.6	1.03	394.9	1.00
Oral Cavity & Pharynx	5,512	14.4	0.20	13.9	0.19	2,425	6.0	0.12	5.8	0.12
Lip	526	1.5	0.07	1.4	0.06	184	0.4	0.03	0.4	0.03
Tongue	1,592	4.0	0.10	3.9	0.10	684	1.7	0.07	1.6	0.06
Major salivary gland	386	1.1	0.06	1.0	0.05	343	0.9	0.05	0.9	0.05
Floor of mouth	307	0.8	0.04	0.7	0.04	136	0.3	0.03	0.3	0.03
Gum & other mouth	562	1.5	0.07	1.4	0.06	490	1.2	0.06	1.1	0.05
Nasopharynx	350	0.9	0.05	0.9	0.05	149	0.4	0.03	0.4	0.03
Oropharynx	240	0.6	0.04	0.6	0.04	65	0.2	0.02	0.1	0.02
Hypopharynx	342	0.9	0.05	0.9	0.05	67	0.2	0.02	0.2	0.02
Digestive System	41,674	116.0	0.58	109.2	0.54	31,630	76.6	0.44	71.9	0.42
Esophagus	2,835	7.7	0.15	7.3	0.14	830	2.0	0.07	1.9	0.07
Stomach	3,686	10.4	0.18	9.8	0.16	1,846	4.5	0.11	4.2	0.10
Small intestine	860	2.4	0.08	2.3	0.08	674	1.7	0.06	1.6	0.06
Colon excl. rectum	15,264	43.3	0.36	40.6	0.33	14,394	34.8	0.29	32.4	0.28
Rectum & rectosigmoid	9,444	25.7	0.27	24.5	0.26	5,435	13.3	0.18	12.7	0.18
Anus, anal canal & anorectum	414	1.1	0.06	1.1	0.05	808	2.0	0.07	1.9	0.07
Liver	2,774	7.2	0.14	6.9	0.13	899	2.2	0.07	2.1	0.07
Gallbladder	335	1.0	0.05	0.9	0.05	608	1.5	0.06	1.4	0.06
Pancreas	4,378	12.3	0.19	11.5	0.18	4,175	10.0	0.16	9.3	0.15
Respiratory System	24,908	70.1	0.45	65.9	0.42	22,521	54.9	0.37	52.3	0.35
Larynx	1,524	4.1	0.11	3.9	0.10	273	0.7	0.04	0.6	0.04
Lung & bronchus	22,999	64.9	0.44	61.0	0.41	22,022	53.6	0.37	51.1	0.35
Bones & Joints	346	1.0	0.06	1.0	0.05	296	0.9	0.05	0.8	0.05
Soft Tissue (incl. heart)	1,208	3.5	0.10	3.3	0.10	916	2.4	0.08	2.3	0.08
Skin (excl. basal & sq.)	7,660	21.4	0.25	20.4	0.24	6,626	17.1	0.21	16.7	0.21
Melanomas of the skin	6,951	19.3	0.24	18.5	0.23	6,104	15.8	0.21	15.5	0.20
Breast (excl. *in situ*)	364	1.0	0.05	1.0	0.05	46,172	115.6	0.55	112.4	0.54
Female Genital System	-	-	-	-	-	19,574	48.9	0.36	47.9	0.35
Cervix uteri	-	-	-	-	-	2,679	7.6	0.15	7.8	0.16
Uterus Corpus	-	-	-	-	-	10,205	24.6	0.25	24.1	0.24
Uterus, NOS[1]	-	-	-	-	-	209	0.5	0.03	0.5	0.03
Ovary	-	-	-	-	-	4,563	11.4	0.17	11.0	0.17
Male Genital System	45,111	120.0	0.58	116.3	0.56	-	-	-	-	-
Prostate	42,600	112.5	0.56	108.5	0.53	-	-	-	-	-
Testis	2,008	6.0	0.14	6.5	0.15	-	-	-	-	-
Penis	389	1.1	0.06	1.0	0.05	-	-	-	-	-
Urinary System	20,091	56.8	0.41	53.2	0.38	8,099	19.8	0.22	18.8	0.21
Bladder (incl. *in situ*)	12,665	36.7	0.33	33.9	0.30	3,994	9.6	0.15	9.0	0.15
Kidney & renal pelvis	6,996	18.9	0.23	18.2	0.22	3,867	9.6	0.16	9.3	0.15
Ureter	266	0.8	0.05	0.7	0.04	175	0.4	0.03	0.4	0.03
Eye & Orbit	307	0.9	0.05	0.8	0.05	267	0.7	0.04	0.7	0.04
Brain & Other Nervous System	2,682	7.6	0.15	7.4	0.15	1,909	5.1	0.12	5.0	0.12
Brain	2,583	7.3	0.15	7.1	0.14	1,819	4.9	0.12	4.7	0.12
Endocrine System	2,317	6.4	0.14	6.4	0.14	5,888	16.4	0.22	16.6	0.22
Thyroid gland	2,034	5.6	0.13	5.6	0.13	5,637	15.7	0.21	16.0	0.22
Lymphomas	8,915	25.2	0.27	24.2	0.26	7,125	18.0	0.22	17.3	0.21
Hodgkin Lymphoma	890	2.6	0.09	2.7	0.09	677	2.0	0.08	2.0	0.08
Non-Hodgkin Lymphoma	8,025	22.6	0.26	21.5	0.24	6,448	16.0	0.20	15.3	0.20
Multiple Myeloma	2,560	7.1	0.14	6.7	0.13	1,875	4.6	0.11	4.3	0.10
Leukemias	6,279	18.0	0.23	17.0	0.22	4,121	10.6	0.17	10.1	0.16
Acute lymphocytic	459	1.5	0.07	1.5	0.07	375	1.3	0.07	1.2	0.07
Chronic lymphocytic	2,859	7.9	0.15	7.4	0.14	1,634	3.9	0.10	3.7	0.09
Acute myeloid	1,573	4.6	0.12	4.3	0.11	1,230	3.2	0.09	3.1	0.09
Chronic myeloid	763	2.2	0.08	2.1	0.08	467	1.2	0.06	1.1	0.05
Other leukemias	625	1.8	0.07	1.7	0.07	415	1.0	0.05	0.9	0.05
Mesothelioma	830	2.5	0.09	2.3	0.08	181	0.4	0.03	0.4	0.03
Kaposi Sarcoma	109	0.3	0.03	0.3	0.03	23	0.1	0.01	0.1	0.01
Ill-defined & Unknown	5,682	16.8	0.23	15.3	0.21	5,234	12.6	0.18	11.5	0.16
Selected tumors (non-invasive)										
Brain & ONS[1] (*benign/borderline*)	1,396	4.0	0.11	3.9	0.11	2,482	6.4	0.13	6.2	0.13
Breast (*in situ*)	21	0.1	0.01	0.1	0.01	6,878	17.4	0.21	17.1	0.21

* Rates are per 100,000 population and are age-adjusted by five-year age groups to the 2000 U.S. standard population based on single years of age and to the 1996 Canadian population standard.

\- Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for All Sites combined.

[1] Abbreviations: NOS - Not Otherwise Specified; ONS - Other Nervous System

Total Cancer Cases and Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates[1] for the Five Most Commonly Diagnosed Cancers
Selected Areas in the NAACCR Canadian Combined[3], 2010-2014, All Races, Males

	All Sites			Prostate			Colon & Rectum			Lung & Bronchus			Bladder			Non-Hodgkin Lymphoma		
	Count	*Rate*	*S.E.*[4]	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
NAACCR Canadian Combined	**176,555**	**488.9**	**1.19**	**42,600**	**112.5**	**0.56**	**24,708**	**69.0**	**0.45**	**22,999**	**64.9**	**0.44**	**12,665**	**36.7**	**0.33**	**8,025**	**22.6**	**0.26**
Alberta	43,725	490.4	2.44	11,240	121.1	1.19	5,606	64.1	0.89	5,065	60.7	0.88	3,048	37.0	0.69	1,994	22.4	0.52
British Columbia	62,484	461.3	1.88	15,257	107.5	0.89	8,552	63.4	0.70	7,598	56.2	0.66	4,820	36.7	0.54	2,875	21.7	0.41
Manitoba	16,246	495.0	3.96	3,399	100.1	1.75	2,366	72.6	1.52	2,203	67.9	1.47	1,018	31.7	1.01	771	23.6	0.87
New Brunswick	12,570	531.1	4.89	3,011	116.8	2.19	1,627	69.3	1.77	2,064	88.5	2.00	943	41.3	1.38	554	24.8	1.09
Newfoundland and Labrador	9,106	549.3	6.04	2,231	123.1	2.72	1,612	99.2	2.59	1,341	82.1	2.34	565	36.1	1.60	380	23.3	1.25
Northwest Territory	356	484.0	30.31	55	74.5	11.57	68	96.0	13.58	53	87.8	13.70	7	7.6	3.15	19	25.0	6.92
Nova Scotia	15,801	544.9	4.47	3,492	112.5	1.95	2,300	79.9	1.72	2,453	85.5	1.77	1,117	39.8	1.22	738	26.4	1.00
Prince Edward Island	2,356	541.5	11.53	594	126.8	5.37	335	77.4	4.36	359	82.0	4.46	142	33.5	2.90	93	22.3	2.39
Saskatchewan	13,911	473.9	4.08	3,321	109.1	1.92	2,242	76.3	1.64	1,863	64.4	1.51	1,005	35.2	1.12	601	21.0	0.87

Total Cancer Cases and Average Annual Age-adjusted (1996 Canadian Standard) Cancer Incidence Rates[2] for the Five Most Commonly Diagnosed Cancers
Selected Areas in the NAACCR Canadian Combined[3], 2010-2014, All Races, Males

	All Sites			Prostate			Colon & Rectum			Lung & Bronchus			Bladder			Non-Hodgkin Lymphoma		
	Count	*Rate*	*S.E.*[4]	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
NAACCR Canadian Combined	**176,555**	**464.7**	**1.12**	**42,600**	**108.5**	**0.53**	**24,708**	**65.1**	**0.42**	**22,999**	**61.0**	**0.41**	**12,665**	**33.9**	**0.30**	**8,025**	**21.5**	**0.24**
Alberta	43,725	466.4	2.29	11,240	117.0	1.13	5,606	60.4	0.83	5,065	57.0	0.82	3,048	34.3	0.63	1,994	21.3	0.49
British Columbia	62,484	437.4	1.77	15,257	103.3	0.84	8,552	59.6	0.65	7,598	52.6	0.61	4,820	33.7	0.49	2,875	20.7	0.39
Manitoba	16,246	470.4	3.75	3,399	96.0	1.67	2,366	68.5	1.43	2,203	64.0	1.38	1,018	29.6	0.94	771	22.3	0.82
New Brunswick	12,570	506.9	4.62	3,011	113.9	2.11	1,627	65.8	1.66	2,064	83.2	1.85	943	38.4	1.27	554	23.5	1.02
Newfoundland and Labrador	9,106	524.3	5.66	2,231	119.2	2.58	1,612	93.9	2.40	1,341	77.5	2.16	565	33.5	1.44	380	22.6	1.21
Northwest Territory	356	460.2	27.75	55	72.0	10.91	68	90.8	12.42	53	82.6	12.46	7	7.8	3.24	19	23.4	6.16
Nova Scotia	15,801	517.4	4.20	3,492	108.5	1.86	2,300	75.2	1.59	2,453	80.2	1.64	1,117	36.8	1.12	738	25.2	0.95
Prince Edward Island	2,356	514.5	10.83	594	121.8	5.06	335	72.9	4.04	359	77.6	4.17	142	31.1	2.65	93	21.4	2.29
Saskatchewan	13,911	451.6	3.90	3,321	105.2	1.85	2,242	72.3	1.55	1,863	61.1	1.43	1,005	32.4	1.04	601	19.8	0.82

1. Rates are per 100,000 population and are age-adjusted by five-year age groups to the 2000 U.S. standard population based on single years of age.
2. Rates are per 100,000 and are age-adjusted to the 1996 Canadian population standard.
3. See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.
4. Abbreviations: S.E. - Standard Error

- Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for the Canadian combined.

Total Cancer Cases and Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates[1] for the Five Most Commonly Diagnosed Cancers
Selected Areas in the NAACCR Canadian Combined[3], 2010-2014, All Races, Females

	All Sites			Breast			Lung & Bronchus			Colon & Rectum			Uterus Corpus & Uterus, NOS[4]			Non-Hodgkin Lymphoma		
	Count	*Rate*	*S.E.*[4]	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
NAACCR Canadian Combined	**164,882**	**410.6**	**1.03**	**46,172**	**115.6**	**0.55**	**22,022**	**53.6**	**0.37**	**19,829**	**48.1**	**0.35**	**10,414**	**25.1**	**0.25**	**6,448**	**16.0**	**0.20**
Alberta	40,556	407.1	2.05	11,844	117.8	1.10	5,104	52.6	0.75	4,317	43.5	0.67	2,597	24.9	0.50	1,513	15.4	0.40
British Columbia	58,176	393.4	1.67	16,780	114.7	0.90	7,566	49.4	0.57	6,916	45.1	0.55	3,635	23.9	0.40	2,332	15.5	0.33
Manitoba	16,333	432.5	3.46	4,342	117.0	1.82	2,310	59.3	1.26	1,940	49.3	1.15	1,199	31.6	0.93	637	16.6	0.67
New Brunswick	11,058	417.1	4.09	2,889	109.4	2.10	1,594	57.3	1.46	1,287	46.6	1.32	669	24.0	0.95	474	17.7	0.84
Newfoundland and Labrador	8,046	449.4	5.16	2,052	113.0	2.57	939	50.0	1.66	1,280	70.4	2.01	545	29.3	1.29	311	17.3	1.01
Northwest Territory	350	433.6	25.55	112	123.7	12.69	34	58.8	10.78	63	79.9	11.10	22	25.2	5.90	10	11.3	3.86
Nova Scotia	14,945	447.3	3.78	3,835	116.2	1.94	2,303	66.0	1.40	1,977	56.7	1.31	847	24.2	0.86	565	16.8	0.73
Prince Edward Island	2,105	418.2	9.39	605	118.4	4.94	287	55.1	3.29	278	54.6	3.35	99	19.3	2.01	80	16.2	1.87
Saskatchewan	13,313	406.6	3.62	3,713	115.6	1.95	1,885	56.2	1.32	1,771	51.6	1.26	801	24.4	0.88	526	15.8	0.71

Total Cancer Cases and Average Annual Age-adjusted (1996 Canadian Standard) Cancer Incidence Rates[2] for the Five Most Commonly Diagnosed Cancers
Selected Areas in the NAACCR Canadian Combined[3], 2010-2014, All Races, Females

	All Sites			Breast			Lung & Bronchus			Colon & Rectum			Uterus Corpus & Uterus, NOS[4]			Non-Hodgkin Lymphoma		
	Count	*Rate*	*S.E.*[4]	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
NAACCR Canadian Combined	**164,882**	**394.9**	**1.00**	**46,172**	**112.4**	**0.54**	**22,022**	**51.1**	**0.35**	**19,829**	**45.1**	**0.33**	**10,414**	**24.6**	**0.25**	**6,448**	**15.3**	**0.20**
Alberta	40,556	392.1	1.99	11,844	114.7	1.07	5,104	50.2	0.72	4,317	40.8	0.63	2,597	24.5	0.49	1,513	14.6	0.38
British Columbia	58,176	377.8	1.62	16,780	111.6	0.89	7,566	46.8	0.55	6,916	42.1	0.52	3,635	23.3	0.40	2,332	14.8	0.32
Manitoba	16,333	415.8	3.38	4,342	113.6	1.78	2,310	56.6	1.21	1,940	46.3	1.09	1,199	31.0	0.92	637	15.8	0.65
New Brunswick	11,058	401.6	3.99	2,889	106.2	2.06	1,594	54.7	1.39	1,287	43.5	1.25	669	23.5	0.93	474	17.1	0.82
Newfoundland and Labrador	8,046	434.1	5.02	2,052	109.8	2.51	939	48.0	1.59	1,280	66.5	1.90	545	28.6	1.26	311	16.8	0.98
Northwest Territory	350	418.1	24.31	112	120.7	12.31	34	55.3	10.01	63	76.3	10.44	22	24.5	5.70	10	11.0	3.71
Nova Scotia	14,945	430.1	3.68	3,835	113.2	1.91	2,303	62.7	1.33	1,977	53.4	1.24	847	23.8	0.85	565	16.1	0.70
Prince Edward Island	2,105	401.2	9.10	605	114.4	4.81	287	52.6	3.15	278	51.1	3.18	99	18.9	1.97	80	15.8	1.85
Saskatchewan	13,313	392.2	3.55	3,713	112.3	1.92	1,885	54.0	1.28	1,771	48.8	1.22	801	23.8	0.87	526	15.0	0.69

1. Rates are per 100,000 population and are age-adjusted by five-year age groups to the 2000 U.S. standard population based on single years of age.
2. Rates are per 100,000 and are age-adjusted to the 1996 Canadian population standard.
3. See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.
4. Abbreviations: NOS - Not Otherwise Specified; S.E. - Standard Error

- Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for the Canadian combined.

Race-Specific Population Indicators (2010-2014)

Combined Population:	
All Races	1,585,379,059
White	-
Black	-
American Indian/Alaskan Native	-
Asian/Pacific Islander	-

Percentage of North America Population Reflected in the Combined Cancer Statistics:	
All Races	90.9%
White	-
Black	-
American Indian/Alaskan Native	-
Asian/Pacific Islander	-

Number of Cases in Combined Cancer Statistics:	
All Races	
White	8,016,815
Black	-
American Indian/Alaskan Native	-
Asian/Pacific Islander	-
	-

Race-Specific Population Counts by Year

Race	2010	2011	2012	2013	2014
All Races	312,140,676	314,575,675	317,062,686	319,501,599	322,098,423

Data Quality Indicators (2010-2014)

Indicators (%)	2010	2011	2012	2013	2014
Missing Race	na	na	na	na	na
Missing Sex	0.0	0.0	0.0	0.0	0.0
Missing County	0.0	0.0	0.0	0.0	0.0
Missing Age	0.0	0.0	0.0	0.0	0.0
Death Certificate Only Cases	1.7	1.6	1.6	1.6	1.8

Site-Specific Microscopic Confirmation (2010-2014)

Cancer Site	% Micro. Confirm	Cancer Site	% Micro. Confirm
All Sites	93.0	Prostate	96.6
Oral Cavity & Pharynx	97.5	Testis	99.1
Esophagus	94.7	Bladder (incl. *in situ*)	97.7
Stomach	96.2	Kidney & Renal Pelvis	89.7
Colon & Rectum	96.1	Brain & Other Nervous System	84.7
Liver	57.6	Thyroid Gland	99.4
Pancreas	81.3	Hodgkin Lymphoma	98.3
Lung & Bronchus	87.8	Non-Hodgkin Lymphoma	96.4
Melanomas of the Skin	99.1	Multiple Myeloma	88.5
Female Breast (excl. *in situ*)	98.5	Leukemias	89.0
Cervix uteri	97.2	Brain & ONS[1] *benign/borderline*	44.5
Uterus Corpus & Uterus NOS[1]	98.5	Breast *in situ*	100.0
Ovary	92.0		

[1] Abbreviations: NOS - Not Otherwise Specified; ONS - Other Nervous System

NAACCR North America Combined Incidence 2010-2014, All Races Combined

NORTH AMERICA

III-95 Combined Incidence – All Races

Total Cancer Cases and Average Annual Age-adjusted Cancer Incidence Rates*
NAACCR North America Combined, 2010-2014, All Races

	MALES (ALL RACES)					FEMALES (ALL RACES)				
SITES	*Count*	*Adj. Rate 2000 U.S.*	*S.E.*	*Adj. Rate 1996 Cdn*	*S.E.*	*Count*	*Adj. Rate 2000 U.S.*	*S.E.*	*Adj. Rate 1996 Cdn*	*S.E.*
All Sites (invasive & in situ bladder)	4,070,764	501.5	0.25	479.1	0.24	3,946,051	417.7	0.21	403.3	0.21
Oral Cavity & Pharynx	147,015	17.2	0.05	16.6	0.04	60,660	6.3	0.03	6.1	0.03
Lip	7,478	1.0	0.01	0.9	0.01	2,834	0.3	0.01	0.3	0.01
Tongue	45,150	5.2	0.02	5.1	0.02	17,891	1.9	0.01	1.8	0.01
Major salivary gland	12,738	1.6	0.01	1.5	0.01	9,019	1.0	0.01	1.0	0.01
Floor of mouth	7,042	0.8	0.01	0.8	0.01	3,178	0.3	0.01	0.3	0.01
Gum & other mouth	15,427	1.9	0.02	1.8	0.01	12,278	1.3	0.01	1.2	0.01
Nasopharynx	6,594	0.8	0.01	0.8	0.01	2,678	0.3	0.01	0.3	0.01
Oropharynx	7,278	0.8	0.01	0.8	0.01	2,257	0.2	0.00	0.2	0.00
Hypopharynx	9,370	1.1	0.01	1.1	0.01	2,216	0.2	0.00	0.2	0.00
Digestive System	798,031	98.5	0.11	93.3	0.11	640,662	65.8	0.08	62.0	0.08
Esophagus	65,650	8.0	0.03	7.6	0.03	17,711	1.8	0.01	1.7	0.01
Stomach	73,503	9.3	0.03	8.7	0.03	44,667	4.6	0.02	4.4	0.02
Small intestine	21,765	2.7	0.02	2.6	0.02	19,420	2.0	0.01	2.0	0.01
Colon excl. rectum	253,538	32.1	0.07	30.2	0.06	256,974	26.4	0.05	24.6	0.05
Rectum & rectosigmoid	122,002	14.8	0.04	14.2	0.04	86,127	9.0	0.03	8.6	0.03
Anus, anal canal & anorectum	12,080	1.5	0.01	1.4	0.01	20,813	2.1	0.02	2.1	0.01
Liver	95,312	10.9	0.04	10.4	0.03	32,000	3.3	0.02	3.1	0.02
Gallbladder	6,414	0.8	0.01	0.8	0.01	13,627	1.4	0.01	1.3	0.01
Pancreas	113,477	14.2	0.04	13.3	0.04	109,137	11.0	0.03	10.3	0.03
Respiratory System	637,727	79.8	0.10	75.5	0.10	533,777	54.8	0.08	52.2	0.07
Larynx	50,182	5.9	0.03	5.7	0.03	12,912	1.3	0.01	1.3	0.01
Lung & bronchus	578,125	72.7	0.10	68.6	0.09	515,051	52.8	0.07	50.3	0.07
Bones & Joints	8,567	1.1	0.01	1.1	0.01	6,868	0.8	0.01	0.8	0.01
Soft Tissue (incl. heart)	30,880	3.9	0.02	3.8	0.02	24,813	2.8	0.02	2.7	0.02
Skin (excl. basal & sq.)	225,639	28.5	0.06	27.1	0.06	158,652	17.5	0.04	17.2	0.04
Melanomas of the skin	208,393	26.2	0.06	25.0	0.06	147,410	16.3	0.04	16.1	0.04
Breast (excl. *in situ*)	10,273	1.3	0.01	1.2	0.01	1,153,821	123.3	0.12	120.0	0.11
Female Genital System	-	-	-	-	-	465,193	49.3	0.07	48.3	0.07
Cervix uteri	-	-	-	-	-	63,013	7.6	0.03	7.7	0.03
Uterus Corpus	-	-	-	-	-	243,735	25.0	0.05	24.5	0.05
Uterus, NOS[1]	-	-	-	-	-	8,187	0.8	0.01	0.8	0.01
Ovary	-	-	-	-	-	107,897	11.5	0.04	11.1	0.03
Male Genital System	1,037,592	121.6	0.12	119.0	0.12	-	-	-	-	-
Prostate	986,455	114.9	0.12	112.0	0.11	-	-	-	-	-
Testis	42,384	5.5	0.03	5.9	0.03	-	-	-	-	-
Penis	6,890	0.9	0.01	0.8	0.01	-	-	-	-	-
Urinary System	463,772	58.9	0.09	55.3	0.08	199,364	20.7	0.05	19.7	0.05
Bladder (incl. *in situ*)	275,044	35.9	0.07	33.2	0.06	87,596	8.9	0.03	8.3	0.03
Kidney & renal pelvis	178,896	21.7	0.05	21.0	0.05	106,317	11.2	0.04	10.9	0.03
Ureter	6,208	0.8	0.01	0.8	0.01	3,754	0.4	0.01	0.4	0.01
Eye & Orbit	7,713	1.0	0.01	0.9	0.01	6,534	0.7	0.01	0.7	0.01
Brain & Other Nervous System	61,909	7.7	0.03	7.5	0.03	49,328	5.5	0.03	5.4	0.03
Brain	58,553	7.3	0.03	7.1	0.03	45,758	5.1	0.02	5.0	0.02
Endocrine System	64,157	7.9	0.03	7.9	0.03	181,522	21.6	0.05	21.9	0.05
Thyroid gland	57,702	7.1	0.03	7.1	0.03	175,611	20.9	0.05	21.3	0.05
Lymphomas	205,021	25.9	0.06	24.9	0.06	169,643	18.1	0.04	17.5	0.04
Hodgkin Lymphoma	23,845	3.0	0.02	3.1	0.02	19,123	2.3	0.02	2.4	0.02
Non-Hodgkin Lymphoma	181,176	22.9	0.05	21.8	0.05	150,520	15.8	0.04	15.1	0.04
Multiple Myeloma	64,121	8.0	0.03	7.6	0.03	51,423	5.3	0.02	5.0	0.02
Leukemias	136,922	17.6	0.05	16.6	0.05	99,642	10.7	0.03	10.2	0.03
Acute lymphocytic	13,876	1.8	0.02	1.8	0.02	10,880	1.4	0.01	1.4	0.01
Chronic lymphocytic	49,449	6.2	0.03	5.8	0.03	32,157	3.2	0.02	3.0	0.02
Acute myeloid	39,671	5.1	0.03	4.9	0.02	32,387	3.5	0.02	3.4	0.02
Chronic myeloid	17,528	2.3	0.02	2.2	0.02	12,857	1.4	0.01	1.3	0.01
Other leukemias	16,398	2.1	0.02	2.0	0.02	11,361	1.2	0.01	1.1	0.01
Mesothelioma	12,875	1.7	0.02	1.6	0.01	3,989	0.4	0.01	0.4	0.01
Kaposi Sarcoma	5,151	0.7	0.01	0.7	0.01	674	0.1	0.00	0.1	0.00
Ill-defined & Unknown	153,399	20.1	0.05	18.5	0.05	139,486	14.1	0.04	13.0	0.04
Selected tumors (non-invasive)										
Brain & ONS[1] (*benign/borderline*)	64,860	8.2	0.03	7.9	0.03	131,751	14.3	0.04	13.8	0.04
Breast (*in situ*)	1,143	0.1	0.00	0.1	0.00	277,885	30.0	0.06	29.4	0.06

* Rates are per 100,000 population and are age-adjusted by five-year age groups to the 2000 U.S. standard population based on single years of age and to the 1996 Canadian population standard.

- Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for All Sites combined.

[1] Abbreviations: NOS - Not Otherwise Specified; ONS - Other Nervous System

Total Cancer Cases and Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates[1] for the Five Most Commonly Diagnosed Cancers
Selected Areas in the NAACCR North American Combined[2], 2010-2014, All Races, Males

	All Sites			Prostate			Lung & Bronchus			Colon & Rectum			Bladder			Melanoma of the Skin		
	Count	*Rate*	*S.E.*[3]	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
NAACCR U.S. Combined	**3,894,209**	**502.1**	**0.26**	**943,855**	**115.0**	**0.12**	**555,126**	**73.0**	**0.10**	**350,832**	**45.9**	**0.08**	**262,379**	**35.8**	**0.07**	**201,442**	**26.6**	**0.06**
Alabama	67,581	533.2	2.11	17,371	129.3	1.01	11,675	92.5	0.88	6,498	52.2	0.67	4,058	34.0	0.55	3,304	27.2	0.49
Alaska	6,949	447.7	6.08	1,548	91.4	2.63	937	67.5	2.49	707	46.6	1.99	451	33.8	1.78	248	15.2	1.07
Arizona	72,737	413.8	1.56	14,985	80.8	0.67	9,842	56.0	0.57	6,778	39.2	0.48	5,534	32.2	0.44	4,429	25.7	0.39
Arkansas	42,168	528.0	2.63	10,204	120.7	1.22	7,946	99.0	1.13	3,937	50.1	0.82	2,745	36.0	0.70	1,921	25.1	0.59
California	407,224	459.3	0.74	100,652	109.3	0.35	42,877	51.0	0.25	37,576	42.5	0.22	25,904	31.4	0.20	25,399	28.9	0.19
CA: Greater California	231,016	468.0	1.00	55,682	108.4	0.47	25,441	54.1	0.35	20,841	42.4	0.30	15,128	32.9	0.27	15,910	32.6	0.26
CA: Greater Bay Area	80,433	465.2	1.68	20,916	115.8	0.82	7,967	49.0	0.56	7,006	40.6	0.50	4,915	30.4	0.44	5,443	31.8	0.44
CA: Los Angeles	95,589	434.0	1.44	23,993	105.9	0.70	9,449	45.6	0.48	9,717	44.2	0.46	5,855	28.7	0.38	4,043	18.5	0.30
Colorado	55,158	445.4	1.98	14,649	110.9	0.96	5,581	48.3	0.67	4,577	37.9	0.58	3,696	32.7	0.56	3,422	27.8	0.49
Connecticut	50,632	526.7	2.40	12,169	118.8	1.10	6,578	69.7	0.88	4,205	44.4	0.70	4,342	46.9	0.73	2,566	27.4	0.55
Delaware	14,378	570.7	4.87	3,837	141.6	2.33	2,025	81.3	1.85	1,069	43.5	1.36	1,017	42.2	1.35	969	39.5	1.30
District of Columbia	7,245	534.4	6.49	2,188	159.7	3.52	883	67.4	2.34	641	48.7	1.99	297	23.3	1.39	168	12.4	0.99
Florida	287,296	479.4	0.91	65,177	103.6	0.41	43,754	71.4	0.34	25,244	42.5	0.27	20,090	33.2	0.24	17,089	29.2	0.23
Georgia	118,328	534.1	1.63	30,481	129.3	0.77	17,952	84.6	0.66	10,783	48.9	0.49	6,580	33.3	0.43	7,314	33.9	0.42
Hawaii	17,112	444.1	3.48	3,582	90.1	1.55	2,178	56.9	1.25	2,003	51.3	1.18	893	23.7	0.81	1,023	26.5	0.85
Idaho	19,579	477.5	3.50	5,187	119.7	1.70	2,219	55.7	1.21	1,612	39.8	1.01	1,479	37.9	1.01	1,311	32.2	0.91
Illinois	164,223	520.1	1.32	39,845	119.4	0.61	24,277	79.2	0.52	16,315	52.5	0.42	11,139	37.5	0.36	7,142	23.0	0.28
Indiana	81,526	494.3	1.78	16,713	95.7	0.76	14,434	88.9	0.76	7,932	48.8	0.56	5,747	36.7	0.50	3,539	21.8	0.38
Iowa	44,898	529.6	2.54	10,064	112.2	1.14	6,611	78.3	0.98	4,310	51.7	0.80	3,190	38.4	0.69	2,394	29.1	0.61
Kansas	38,405	521.8	2.72	9,518	124.1	1.30	5,306	73.1	1.02	3,439	47.4	0.82	2,748	38.8	0.75	2,167	30.2	0.66
Kentucky	67,517	590.8	2.35	13,778	113.0	0.99	13,203	116.3	1.04	6,687	59.3	0.75	4,304	40.2	0.63	3,403	30.7	0.54
Louisiana	65,307	576.5	2.33	17,311	144.4	1.13	10,066	90.8	0.94	6,303	56.0	0.73	3,530	33.5	0.58	2,482	23.0	0.48
Maine	21,320	517.6	3.65	4,491	99.8	1.53	3,464	83.9	1.46	1,717	42.7	1.06	1,892	47.5	1.12	1,134	28.1	0.86
Maryland	72,723	502.5	1.92	20,406	131.5	0.95	9,265	66.6	0.71	6,025	42.6	0.56	4,916	37.0	0.54	4,032	28.5	0.46
Massachusetts	87,754	504.1	1.74	21,277	114.2	0.80	11,876	70.3	0.66	7,327	42.9	0.51	6,718	40.5	0.50	4,332	25.4	0.40
Michigan	138,639	516.4	1.42	36,195	126.3	0.68	20,448	77.2	0.55	11,508	44.0	0.42	9,972	39.1	0.40	5,904	22.7	0.30
MI: Metropolitan Detroit	57,850	583.8	2.50	16,298	155.2	1.25	8,152	85.0	0.97	4,722	48.5	0.73	3,965	42.4	0.69	2,041	21.2	0.48
Mississippi	41,771	558.7	2.82	10,748	135.5	1.34	7,466	101.4	1.21	4,300	58.2	0.92	2,145	30.9	0.69	1,662	23.1	0.59
Missouri	80,984	502.0	1.80	17,428	101.0	0.78	14,218	89.0	0.76	7,820	49.7	0.57	5,198	33.8	0.48	3,831	24.3	0.40
Montana	14,771	485.9	4.13	3,856	116.6	1.93	1,810	60.9	1.47	1,334	44.2	1.25	1,056	36.0	1.13	897	30.3	1.05
Nebraska	24,322	507.5	3.31	5,977	119.6	1.58	3,374	71.8	1.26	2,375	49.9	1.04	1,704	37.1	0.91	1,122	23.5	0.71
New Hampshire	19,813	527.5	3.88	5,094	123.5	1.79	2,581	71.2	1.45	1,504	40.9	1.09	1,714	48.1	1.20	1,218	33.3	0.99
New Jersey	122,895	543.0	1.58	33,215	139.4	0.78	14,382	65.6	0.56	10,728	47.9	0.47	8,887	41.5	0.45	6,416	28.7	0.37
New York	270,901	546.4	1.07	71,412	136.8	0.52	33,864	69.9	0.39	22,839	46.9	0.32	19,314	41.0	0.30	11,099	22.8	0.22

Total Cancer Cases and Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates[1] for the Five Most Commonly Diagnosed Cancers
Selected Areas in the NAACCR North American Combined[2], 2010-2014, All Races, Males

	All Sites			Prostate			Lung & Bronchus			Colon & Rectum			Bladder			Melanoma of the Skin		
	Count	*Rate*	*S.E.*[3]	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
North Carolina	130,183	527.9	1.51	32,878	125.0	0.71	21,415	88.5	0.62	10,565	43.7	0.44	8,158	35.5	0.40	7,449	31.4	0.37
North Dakota	9,651	510.4	5.31	2,446	123.2	2.55	1,308	70.8	1.99	1,000	53.8	1.74	683	37.0	1.44	466	24.9	1.18
Ohio	157,260	504.6	1.30	37,060	111.8	0.59	25,857	83.7	0.53	14,626	47.9	0.41	11,494	38.6	0.37	7,519	24.9	0.29
Oklahoma	49,300	502.0	2.31	11,133	108.3	1.05	8,448	86.6	0.96	4,705	48.6	0.72	3,158	33.6	0.61	2,276	23.9	0.51
Oregon	50,882	467.9	2.13	11,791	101.1	0.96	6,714	63.1	0.79	4,313	40.8	0.64	3,946	38.0	0.62	3,324	31.1	0.55
Pennsylvania	196,353	543.9	1.25	45,104	117.5	0.56	28,146	78.3	0.47	17,716	49.9	0.38	15,434	43.9	0.36	10,098	28.7	0.29
Rhode Island	14,563	518.5	4.40	3,240	108.5	1.95	2,139	77.7	1.72	1,135	41.2	1.25	1,234	45.1	1.31	781	28.5	1.04
South Carolina	65,603	521.3	2.11	16,502	121.4	0.98	10,738	85.7	0.86	5,531	44.7	0.62	4,026	34.7	0.57	3,466	28.4	0.50
South Dakota	11,379	498.4	4.77	2,815	116.8	2.24	1,568	69.7	1.79	1,165	51.6	1.54	771	35.1	1.28	558	24.9	1.08
Tennessee	89,477	529.2	1.83	21,512	119.1	0.84	16,028	95.7	0.78	7,724	46.8	0.55	5,466	34.5	0.48	4,232	26.0	0.41
Texas	262,427	465.1	0.94	58,205	99.4	0.43	36,319	68.0	0.37	26,047	46.3	0.30	13,966	27.4	0.24	9,825	17.7	0.19
Utah	25,518	457.4	2.94	7,404	130.6	1.56	1,717	32.7	0.81	1,942	35.5	0.82	1,515	29.8	0.78	2,643	47.7	0.95
Vermont	9,122	493.3	5.36	1,989	98.4	2.28	1,325	71.8	2.04	721	39.3	1.52	691	38.9	1.52	715	39.5	1.54
Virginia	93,384	460.8	1.56	23,473	107.6	0.72	14,291	73.1	0.63	8,205	41.1	0.47	5,791	31.1	0.42	4,861	24.6	0.36
Washington	88,331	502.1	1.74	21,829	116.3	0.81	11,011	65.5	0.64	7,123	40.9	0.50	6,175	37.6	0.49	5,344	30.5	0.43
WA: Seattle/Puget Sound	62,082	525.1	2.18	15,504	123.3	1.03	7,398	66.1	0.79	4,930	41.8	0.61	4,246	38.9	0.62	3,978	33.7	0.55
West Virginia	29,544	523.7	3.14	6,057	99.6	1.31	5,639	98.7	1.35	2,939	53.1	1.01	2,175	39.9	0.88	1,350	25.3	0.71
Wisconsin	80,116	515.8	1.86	19,236	116.0	0.86	10,633	69.5	0.69	6,648	43.6	0.55	5,912	39.9	0.53	4,173	27.3	0.43
Wyoming	6,960	447.2	5.57	1,823	108.5	2.64	738	49.4	1.89	634	41.7	1.72	524	36.0	1.63	425	28.5	1.43
NAACCR Canadian Combined	**176,555**	**488.9**	**1.19**	**42,600**	**112.5**	**0.56**	**22,999**	**64.9**	**0.44**	**24,708**	**69.0**	**0.45**	**12,665**	**36.7**	**0.33**	**6,951**	**19.3**	**0.24**
Alberta	43,725	490.4	2.44	11,240	121.1	1.19	5,065	60.7	0.88	5,606	64.1	0.89	3,048	37.0	0.69	1,585	17.4	0.45
British Columbia	62,484	461.3	1.88	15,257	107.5	0.89	7,598	56.2	0.66	8,552	63.4	0.70	4,820	36.7	0.54	2,875	21.4	0.41
Manitoba	16,246	495.0	3.96	3,399	100.1	1.75	2,203	67.9	1.47	2,366	72.6	1.52	1,018	31.7	1.01	510	15.7	0.71
New Brunswick	12,570	531.1	4.89	3,011	116.8	2.19	2,064	88.5	2.00	1,627	69.3	1.77	943	41.3	1.38	430	18.7	0.93
Newfoundland and Labrador	9,106	549.3	6.04	2,231	123.1	2.72	1,341	82.1	2.34	1,612	99.2	2.59	565	36.1	1.60	253	15.8	1.04
Northwest Territory	356	484.0	30.31	55	74.5	11.57	53	87.8	13.70	68	96.0	13.58	7	7.6	3.15	9	8.7	3.16
Nova Scotia	15,801	544.9	4.47	3,492	112.5	1.95	2,453	85.5	1.77	2,300	79.9	1.72	1,117	39.8	1.22	767	26.9	1.01
Prince Edward Island	2,356	541.5	11.53	594	126.8	5.37	359	82.0	4.46	335	77.4	4.36	142	33.5	2.90	116	28.1	2.70
Saskatchewan	13,911	473.9	4.08	3,321	109.1	1.92	1,863	64.4	1.51	2,242	76.3	1.64	1,005	35.2	1.12	406	13.9	0.70
NAACCR North American Combined	**4,070,764**	**501.5**	**0.25**	**986,455**	**114.9**	**0.12**	**578,125**	**72.7**	**0.10**	**375,540**	**46.9**	**0.08**	**275,044**	**35.9**	**0.07**	**208,393**	**26.2**	**0.06**

1. Rates are per 100,000 population and are age-adjusted by five-year age groups to the 2000 U.S. standard population based on single years of age.
2. See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.
3. Abbreviations: S.E. - Standard Error

\- Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for the combined.

Total Cancer Cases and Average Annual Age-adjusted (1996 Canadian Standard) Cancer Incidence Rates[1] for the Five Most Commonly Diagnosed Cancers
Selected Areas in the NAACCR North American Combined[2], 2010-2014, All Races, Males

	All Sites			Prostate			Lung & Bronchus			Colon & Rectum			Bladder			Melanoma of the Skin		
	Count	*Rate*	*S.E.*[3]	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
NAACCR U.S. Combined	**3,894,209**	**479.8**	**0.25**	**943,855**	**112.2**	**0.12**	**555,126**	**69.0**	**0.09**	**350,832**	**43.3**	**0.07**	**262,379**	**33.1**	**0.07**	**201,442**	**25.3**	**0.06**
Alabama	67,581	510.1	2.00	17,371	125.4	0.96	11,675	87.9	0.82	6,498	49.5	0.63	4,058	31.7	0.50	3,304	25.9	0.46
Alaska	6,949	426.5	5.58	1,548	88.7	2.46	937	63.3	2.25	707	43.9	1.81	451	31.7	1.61	248	14.8	1.01
Arizona	72,737	394.6	1.48	14,985	78.5	0.64	9,842	52.7	0.53	6,778	37.1	0.45	5,534	29.7	0.40	4,429	24.4	0.37
Arkansas	42,168	506.4	2.50	10,204	117.8	1.18	7,946	94.4	1.07	3,937	47.6	0.77	2,745	33.4	0.64	1,921	23.8	0.55
California	407,224	438.0	0.70	100,652	106.6	0.34	42,877	47.7	0.23	37,576	40.1	0.21	25,904	28.9	0.18	25,399	27.4	0.17
CA: Greater California	231,016	446.1	0.94	55,682	105.6	0.45	25,441	50.6	0.32	20,841	40.1	0.28	15,128	30.2	0.25	15,910	30.9	0.25
CA: Greater Bay Area	80,433	443.5	1.59	20,916	113.0	0.80	7,967	45.6	0.52	7,006	38.2	0.47	4,915	27.9	0.40	5,443	30.2	0.42
CA: Los Angeles	95,589	414.7	1.37	23,993	103.5	0.68	9,449	42.7	0.45	9,717	41.9	0.43	5,855	26.4	0.35	4,043	17.5	0.28
Colorado	55,158	425.0	1.86	14,649	108.0	0.92	5,581	45.2	0.62	4,577	35.6	0.54	3,696	30.2	0.51	3,422	26.6	0.47
Connecticut	50,632	502.9	2.29	12,169	116.5	1.08	6,578	65.8	0.83	4,205	41.7	0.66	4,342	43.3	0.67	2,566	25.9	0.52
Delaware	14,378	543.8	4.60	3,837	138.1	2.25	2,025	76.4	1.71	1,069	40.8	1.27	1,017	38.7	1.22	969	37.5	1.23
District of Columbia	7,245	516.0	6.21	2,188	155.6	3.40	883	64.6	2.22	641	46.4	1.88	297	21.8	1.29	168	12.0	0.94
Florida	287,296	459.4	0.87	65,177	101.5	0.40	43,754	67.7	0.33	25,244	40.4	0.26	20,090	30.8	0.22	17,089	27.6	0.22
Georgia	118,328	510.9	1.53	30,481	126.1	0.74	17,952	80.0	0.61	10,783	46.4	0.46	6,580	30.8	0.39	7,314	32.2	0.39
Hawaii	17,112	424.6	3.32	3,582	87.1	1.48	2,178	53.5	1.17	2,003	49.1	1.12	893	22.0	0.75	1,023	25.7	0.82
Idaho	19,579	455.2	3.30	5,187	115.8	1.63	2,219	52.1	1.12	1,612	37.6	0.95	1,479	35.1	0.92	1,311	31.1	0.87
Illinois	164,223	496.5	1.25	39,845	116.5	0.59	24,277	74.7	0.49	16,315	49.4	0.39	11,139	34.8	0.33	7,142	21.9	0.26
Indiana	81,526	472.0	1.68	16,713	92.9	0.73	14,434	84.3	0.71	7,932	46.1	0.53	5,747	34.0	0.45	3,539	20.8	0.36
Iowa	44,898	505.2	2.43	10,064	109.2	1.10	6,611	73.9	0.92	4,310	48.5	0.75	3,190	35.5	0.64	2,394	27.9	0.59
Kansas	38,405	499.3	2.59	9,518	120.7	1.26	5,306	69.1	0.96	3,439	44.8	0.78	2,748	36.0	0.70	2,167	28.9	0.63
Kentucky	67,517	564.7	2.21	13,778	109.9	0.95	13,203	110.3	0.98	6,687	56.2	0.70	4,304	37.3	0.58	3,403	29.4	0.51
Louisiana	65,307	553.2	2.21	17,311	141.1	1.09	10,066	86.3	0.88	6,303	53.3	0.69	3,530	31.3	0.54	2,482	21.8	0.45
Maine	21,320	493.4	3.47	4,491	97.3	1.47	3,464	79.2	1.37	1,717	40.2	1.00	1,892	44.1	1.03	1,134	26.8	0.82
Maryland	72,723	480.4	1.82	20,406	128.8	0.92	9,265	62.8	0.67	6,025	40.1	0.53	4,916	34.0	0.49	4,032	27.1	0.44
Massachusetts	87,754	482.1	1.66	21,277	111.8	0.78	11,876	66.2	0.62	7,327	40.3	0.48	6,718	37.6	0.47	4,332	24.2	0.38
Michigan	138,639	493.8	1.35	36,195	123.4	0.66	20,448	73.0	0.52	11,508	41.4	0.39	9,972	36.3	0.37	5,904	21.7	0.29
MI: Metropolitan Detroit	57,850	558.5	2.38	16,298	151.7	1.22	8,152	80.2	0.91	4,722	45.6	0.68	3,965	39.3	0.64	2,041	20.1	0.46
Mississippi	41,771	536.4	2.67	10,748	132.5	1.30	7,466	96.4	1.13	4,300	55.5	0.86	2,145	28.7	0.63	1,662	22.1	0.55
Missouri	80,984	480.3	1.71	17,428	98.8	0.76	14,218	84.4	0.72	7,820	46.8	0.54	5,198	31.4	0.44	3,831	23.3	0.38
Montana	14,771	465.5	3.93	3,856	113.7	1.86	1,810	57.0	1.36	1,334	42.2	1.18	1,056	33.3	1.04	897	29.3	1.02
Nebraska	24,322	485.3	3.17	5,977	116.1	1.53	3,374	67.9	1.19	2,375	47.2	0.99	1,704	34.4	0.84	1,122	22.7	0.69
New Hampshire	19,813	504.2	3.69	5,094	121.1	1.74	2,581	66.7	1.34	1,504	38.3	1.02	1,714	44.7	1.10	1,218	31.7	0.93
New Jersey	122,895	518.3	1.50	33,215	136.0	0.76	14,382	61.8	0.52	10,728	45.1	0.44	8,887	38.3	0.41	6,416	27.2	0.35
New York	270,901	522.2	1.02	71,412	133.6	0.51	33,864	65.9	0.36	22,839	44.2	0.30	19,314	37.9	0.28	11,099	21.7	0.21

Total Cancer Cases and Average Annual Age-adjusted (1996 Canadian Standard) Cancer Incidence Rates[1] for the Five Most Commonly Diagnosed Cancers
Selected Areas in the NAACCR North American Combined[2], 2010-2014, All Races, Males

	All Sites			Prostate			Lung & Bronchus			Colon & Rectum			Bladder			Melanoma of the Skin		
	Count	*Rate*	*S.E.*[3]	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
North Carolina	130,183	505.3	1.42	32,878	121.7	0.68	21,415	83.9	0.58	10,565	41.4	0.41	8,158	32.9	0.37	7,449	29.9	0.35
North Dakota	9,651	488.6	5.09	2,446	120.1	2.48	1,308	66.5	1.87	1,000	51.3	1.66	683	34.1	1.33	466	24.3	1.16
Ohio	157,260	482.8	1.24	37,060	109.1	0.58	25,857	79.3	0.50	14,626	45.2	0.38	11,494	35.8	0.34	7,519	23.8	0.28
Oklahoma	49,300	480.0	2.19	11,133	104.9	1.00	8,448	82.1	0.90	4,705	46.2	0.68	3,158	31.3	0.56	2,276	22.7	0.48
Oregon	50,882	447.1	2.02	11,791	98.6	0.92	6,714	59.3	0.73	4,313	38.4	0.60	3,946	35.1	0.57	3,324	29.9	0.53
Pennsylvania	196,353	519.9	1.20	45,104	114.7	0.55	28,146	74.0	0.45	17,716	46.9	0.36	15,434	40.7	0.33	10,098	27.3	0.28
Rhode Island	14,563	494.6	4.20	3,240	106.1	1.90	2,139	73.0	1.61	1,135	38.5	1.17	1,234	41.9	1.22	781	27.3	1.00
South Carolina	65,603	497.9	1.98	16,502	118.3	0.93	10,738	81.1	0.80	5,531	42.4	0.58	4,026	31.9	0.51	3,466	27.1	0.47
South Dakota	11,379	476.4	4.56	2,815	114.0	2.18	1,568	65.1	1.67	1,165	48.7	1.46	771	32.4	1.18	558	24.1	1.05
Tennessee	89,477	507.1	1.73	21,512	116.2	0.80	16,028	90.9	0.73	7,724	44.4	0.51	5,466	32.1	0.44	4,232	24.8	0.39
Texas	262,427	443.5	0.88	58,205	96.2	0.41	36,319	64.1	0.34	26,047	43.9	0.28	13,966	25.3	0.22	9,825	16.8	0.17
Utah	25,518	438.2	2.79	7,404	127.2	1.50	1,717	30.8	0.75	1,942	33.6	0.77	1,515	27.6	0.72	2,643	45.5	0.90
Vermont	9,122	471.7	5.09	1,989	95.6	2.19	1,325	67.9	1.91	721	37.0	1.42	691	36.3	1.40	715	37.8	1.46
Virginia	93,384	439.9	1.47	23,473	105.1	0.70	14,291	68.9	0.59	8,205	38.8	0.44	5,791	28.7	0.38	4,861	23.3	0.34
Washington	88,331	478.4	1.64	21,829	113.4	0.78	11,011	61.3	0.59	7,123	38.5	0.46	6,175	34.6	0.45	5,344	29.3	0.41
WA: Seattle/Puget Sound	62,082	499.7	2.05	15,504	120.1	0.99	7,398	61.8	0.73	4,930	39.3	0.57	4,246	35.6	0.56	3,978	32.3	0.52
West Virginia	29,544	501.0	2.98	6,057	96.8	1.26	5,639	93.9	1.27	2,939	50.3	0.95	2,175	37.3	0.81	1,350	24.2	0.68
Wisconsin	80,116	492.5	1.77	19,236	113.1	0.83	10,633	65.6	0.65	6,648	41.0	0.51	5,912	36.9	0.49	4,173	26.1	0.41
Wyoming	6,960	429.0	5.28	1,823	106.2	2.55	738	46.5	1.75	634	39.5	1.61	524	33.7	1.50	425	27.3	1.36
NAACCR Canadian Combined	**176,555**	**464.7**	**1.12**	**42,600**	**108.5**	**0.53**	**22,999**	**61.0**	**0.41**	**24,708**	**65.1**	**0.42**	**12,665**	**33.9**	**0.30**	**6,951**	**18.5**	**0.23**
Alberta	43,725	466.4	2.29	11,240	117.0	1.13	5,065	57.0	0.82	5,606	60.4	0.83	3,048	34.3	0.63	1,585	16.5	0.43
British Columbia	62,484	437.4	1.77	15,257	103.3	0.84	7,598	52.6	0.61	8,552	59.6	0.65	4,820	33.7	0.49	2,875	20.4	0.39
Manitoba	16,246	470.4	3.75	3,399	96.0	1.67	2,203	64.0	1.38	2,366	68.5	1.43	1,018	29.6	0.94	510	14.8	0.67
New Brunswick	12,570	506.9	4.62	3,011	113.9	2.11	2,064	83.2	1.85	1,627	65.8	1.66	943	38.4	1.27	430	18.0	0.89
Newfoundland and Labrador	9,106	524.3	5.66	2,231	119.2	2.58	1,341	77.5	2.16	1,612	93.9	2.40	565	33.5	1.44	253	15.3	1.00
Northwest Territory	356	460.2	27.75	55	72.0	10.91	53	82.6	12.46	68	90.8	12.42	7	7.8	3.24	9	8.6	3.19
Nova Scotia	15,801	517.4	4.20	3,492	108.5	1.86	2,453	80.2	1.64	2,300	75.2	1.59	1,117	36.8	1.12	767	25.9	0.96
Prince Edward Island	2,356	514.5	10.83	594	121.8	5.06	359	77.6	4.17	335	72.9	4.04	142	31.1	2.65	116	27.0	2.59
Saskatchewan	13,911	451.6	3.90	3,321	105.2	1.85	1,863	61.1	1.43	2,242	72.3	1.55	1,005	32.4	1.04	406	13.3	0.67
NAACCR North American Combined	**4,070,764**	**479.1**	**0.24**	**986,455**	**112.0**	**0.11**	**578,125**	**68.6**	**0.09**	**375,540**	**44.3**	**0.07**	**275,044**	**33.2**	**0.06**	**208,393**	**25.0**	**0.06**

1. Rates are per 100,000 and are age-adjusted to the 1996 Canadian population standard.
2. See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.
3. Abbreviations: S.E. - Standard Error

\- Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for the combined.

Total Cancer Cases and Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates[1] for the Five Most Commonly Diagnosed Cancers
Selected Areas in the NAACCR North American Combined[2], 2010-2014, All Races, Females

	All Sites			Breast			Lung & Bronchus			Colon & Rectum			Uterus Corpus & Uterus, NOS[3]			Thyroid		
	Count	*Rate*	*S.E.*[3]	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
NAACCR U.S. Combined	**3,781,169**	**418.0**	**0.22**	**1,107,649**	**123.6**	**0.12**	**493,029**	**52.8**	**0.08**	**323,272**	**34.8**	**0.06**	**241,508**	**25.9**	**0.05**	**169,974**	**21.1**	**0.05**
Alabama	59,329	394.9	1.66	17,811	119.6	0.92	8,235	52.1	0.58	5,651	36.9	0.50	2,968	19.1	0.36	1,712	13.0	0.32
Alaska	6,568	411.9	5.38	2,085	125.4	2.90	773	53.3	2.04	620	40.2	1.71	431	24.6	1.26	299	17.5	1.05
Arizona	72,444	375.0	1.42	21,537	112.4	0.79	9,264	45.4	0.48	5,908	29.8	0.40	4,260	21.6	0.34	3,979	23.7	0.39
Arkansas	36,111	398.1	2.15	10,122	112.7	1.15	5,818	60.7	0.81	3,456	37.1	0.64	1,997	21.6	0.49	1,180	15.1	0.45
California	404,208	388.5	0.62	125,201	120.7	0.35	41,600	39.9	0.20	34,469	32.6	0.18	26,391	24.7	0.15	18,747	19.1	0.14
CA: Greater California	225,193	395.6	0.85	69,313	122.1	0.47	24,875	43.4	0.28	18,789	32.6	0.24	13,933	23.7	0.20	10,305	19.4	0.19
CA: Greater Bay Area	79,420	392.4	1.42	25,087	124.6	0.80	7,912	39.0	0.45	6,667	32.3	0.40	5,363	25.7	0.36	3,212	16.9	0.30
CA: Los Angeles	99,469	370.0	1.19	30,772	114.6	0.66	8,801	32.9	0.36	9,004	32.9	0.35	7,095	25.8	0.31	5,229	20.0	0.28
Colorado	54,644	390.2	1.71	17,541	123.7	0.95	5,751	41.6	0.56	4,259	30.8	0.48	3,321	22.3	0.40	3,154	23.5	0.43
Connecticut	52,718	458.4	2.06	15,767	139.2	1.14	6,767	56.8	0.71	4,159	34.3	0.55	3,483	29.2	0.50	2,827	29.1	0.56
Delaware	13,375	456.4	4.05	3,886	133.1	2.20	1,930	62.9	1.45	986	32.9	1.07	890	29.0	1.00	578	23.6	1.01
District of Columbia	7,606	446.1	5.24	2,389	143.5	3.00	845	49.5	1.74	690	40.3	1.57	549	31.2	1.36	396	22.5	1.18
Florida	266,096	396.9	0.80	75,212	115.5	0.44	38,674	53.1	0.27	22,825	32.3	0.22	15,788	23.1	0.19	9,412	17.6	0.19
Georgia	110,666	410.9	1.25	33,614	123.5	0.68	14,072	52.2	0.45	9,578	35.6	0.37	6,133	21.6	0.28	4,359	16.6	0.26
Hawaii	17,260	407.4	3.21	5,670	136.0	1.87	1,717	38.1	0.94	1,599	36.5	0.94	1,237	30.0	0.88	831	23.2	0.83
Idaho	18,042	410.5	3.12	5,295	120.5	1.69	2,117	46.8	1.03	1,437	32.5	0.88	1,111	24.2	0.74	980	24.9	0.81
Illinois	165,887	437.9	1.10	48,665	130.0	0.60	22,372	57.8	0.39	14,948	38.1	0.32	11,336	29.0	0.28	7,370	21.7	0.26
Indiana	82,508	425.6	1.51	23,084	120.1	0.81	12,158	60.8	0.56	7,666	38.5	0.45	5,390	27.0	0.38	3,015	17.8	0.33
Iowa	42,359	439.5	2.21	11,520	122.8	1.19	5,377	53.0	0.74	4,114	39.8	0.64	3,042	31.2	0.58	1,575	20.1	0.52
Kansas	35,882	430.0	2.33	10,182	123.5	1.26	4,596	52.8	0.79	3,129	36.1	0.66	2,104	24.7	0.55	1,833	25.3	0.61
Kentucky	63,000	472.7	1.92	16,413	123.3	0.98	11,053	79.7	0.77	5,774	42.4	0.57	3,521	25.8	0.44	2,526	21.6	0.44
Louisiana	55,904	420.8	1.81	16,330	123.2	0.99	7,513	54.9	0.64	5,551	41.4	0.57	2,610	19.1	0.38	2,371	19.4	0.41
Maine	21,056	454.0	3.25	5,743	125.5	1.72	3,189	65.3	1.18	1,638	34.3	0.88	1,506	30.9	0.82	822	22.3	0.82
Maryland	74,065	421.0	1.58	23,092	131.0	0.88	9,245	52.0	0.55	5,920	33.3	0.44	4,867	26.4	0.39	3,554	22.0	0.38
Massachusetts	94,120	449.3	1.50	28,058	136.1	0.83	13,038	60.5	0.54	7,338	33.7	0.40	6,255	28.9	0.37	5,322	29.1	0.41
Michigan	131,536	426.3	1.21	37,301	122.2	0.65	18,930	58.8	0.43	10,849	34.0	0.33	9,267	28.9	0.31	5,218	19.8	0.28
MI: Metropolitan Detroit	55,534	458.5	2.00	15,788	131.6	1.07	8,076	64.8	0.73	4,704	37.5	0.56	3,787	30.3	0.50	2,511	23.6	0.48
Mississippi	35,981	407.4	2.19	10,104	115.5	1.17	5,172	56.3	0.79	3,786	41.8	0.69	1,778	19.6	0.47	1,261	15.9	0.46
Missouri	79,928	427.6	1.55	23,218	125.9	0.85	12,540	64.3	0.58	7,077	36.6	0.45	4,997	26.1	0.38	2,866	17.8	0.34
Montana	13,450	424.7	3.80	3,893	123.1	2.06	1,791	53.8	1.30	1,117	34.1	1.05	860	25.9	0.92	561	21.6	0.95
Nebraska	22,633	416.0	2.84	6,538	121.8	1.55	2,834	50.2	0.96	2,170	38.2	0.84	1,470	26.6	0.71	975	21.0	0.69
New Hampshire	19,517	463.1	3.42	5,914	140.4	1.88	2,726	63.3	1.24	1,481	34.2	0.91	1,466	32.8	0.88	845	23.3	0.83
New Jersey	125,460	452.9	1.31	36,075	132.0	0.71	15,020	52.5	0.44	10,947	37.8	0.37	8,874	31.1	0.34	6,800	27.9	0.35
New York	273,599	451.0	0.88	76,791	129.0	0.48	34,081	54.2	0.30	22,514	35.7	0.24	19,341	31.0	0.23	15,178	28.3	0.23

Total Cancer Cases and Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Rates[1] for the Five Most Commonly Diagnosed Cancers
Selected Areas in the NAACCR North American Combined[2], 2010-2014, All Races, Females

	All Sites			Breast			Lung & Bronchus			Colon & Rectum			Uterus Corpus & Uterus, NOS[3]			Thyroid		
	Count	*Rate*	*S.E.*[3]	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
North Carolina	124,067	421.4	1.22	37,914	129.4	0.68	17,109	56.1	0.43	9,839	33.0	0.34	7,459	24.4	0.29	4,854	18.3	0.27
North Dakota	8,513	414.5	4.67	2,397	121.4	2.58	1,048	48.9	1.54	872	40.0	1.40	527	25.4	1.14	426	24.4	1.23
Ohio	154,863	424.4	1.11	44,328	122.9	0.60	22,438	59.0	0.40	13,624	35.8	0.31	10,697	28.4	0.28	6,693	21.7	0.27
Oklahoma	46,231	412.3	1.96	13,103	117.8	1.05	6,873	58.7	0.72	4,236	36.9	0.58	2,496	21.7	0.45	1,731	17.5	0.43
Oregon	50,591	419.2	1.92	15,188	126.0	1.06	6,696	53.4	0.67	3,881	31.2	0.52	3,342	26.5	0.47	2,010	19.6	0.45
Pennsylvania	195,598	461.3	1.08	53,851	129.8	0.58	25,224	56.1	0.36	16,964	37.5	0.30	14,004	32.1	0.28	10,515	30.5	0.31
Rhode Island	15,625	457.9	3.80	4,331	130.3	2.04	2,265	63.7	1.37	1,271	34.7	1.01	1,117	31.9	0.98	784	27.1	1.00
South Carolina	60,486	411.9	1.72	18,597	127.2	0.96	8,268	53.7	0.60	5,051	33.8	0.49	3,505	22.7	0.39	2,222	17.2	0.38
South Dakota	10,661	428.1	4.31	3,167	130.7	2.41	1,324	50.8	1.42	1,041	38.8	1.25	708	28.2	1.09	412	19.8	1.01
Tennessee	82,826	418.7	1.49	23,909	121.1	0.80	12,555	61.0	0.55	7,307	36.4	0.43	4,633	22.7	0.34	3,231	18.5	0.33
Texas	251,958	380.2	0.77	74,269	111.5	0.41	29,302	44.7	0.26	21,373	32.2	0.22	14,447	21.1	0.18	11,944	18.3	0.17
Utah	23,175	373.9	2.49	7,012	114.7	1.39	1,449	24.1	0.64	1,686	27.7	0.68	1,643	25.9	0.65	1,856	28.8	0.68
Vermont	9,030	437.4	4.78	2,662	130.1	2.62	1,285	59.7	1.71	710	33.0	1.28	710	32.8	1.27	316	19.0	1.13
Virginia	94,849	398.7	1.32	30,271	126.9	0.74	12,257	50.8	0.47	7,934	33.1	0.38	5,915	23.8	0.31	4,277	19.6	0.31
Washington	86,489	437.9	1.53	26,686	134.9	0.85	10,713	53.7	0.53	6,534	32.9	0.42	5,342	25.6	0.36	3,679	20.4	0.34
WA: Seattle/Puget Sound	60,793	450.7	1.87	18,947	139.9	1.04	7,271	54.0	0.65	4,519	33.4	0.51	3,756	26.3	0.44	2,586	20.6	0.41
West Virginia	27,673	443.8	2.77	7,112	114.8	1.42	4,416	66.2	1.01	2,664	40.9	0.81	1,935	30.4	0.72	1,226	24.3	0.72
Wisconsin	76,465	437.3	1.63	21,974	127.9	0.89	9,884	54.8	0.56	6,139	33.6	0.44	5,428	29.9	0.41	2,947	19.7	0.37
Wyoming	6,117	382.5	5.04	1,827	113.5	2.74	725	44.1	1.67	490	30.5	1.41	357	21.5	1.18	305	21.3	1.26
NAACCR Canadian Combined	**164,882**	**410.6**	**1.03**	**46,172**	**115.6**	**0.55**	**22,022**	**53.6**	**0.37**	**19,829**	**48.1**	**0.35**	**10,414**	**25.1**	**0.25**	**5,637**	**15.7**	**0.21**
Alberta	40,556	407.1	2.05	11,844	117.8	1.10	5,104	52.6	0.75	4,317	43.5	0.67	2,597	24.9	0.50	1,781	17.8	0.43
British Columbia	58,176	393.4	1.67	16,780	114.7	0.90	7,566	49.4	0.57	6,916	45.1	0.55	3,635	23.9	0.40	1,500	11.7	0.31
Manitoba	16,333	432.5	3.46	4,342	117.0	1.82	2,310	59.3	1.26	1,940	49.3	1.15	1,199	31.6	0.93	539	16.5	0.73
New Brunswick	11,058	417.1	4.09	2,889	109.4	2.10	1,594	57.3	1.46	1,287	46.6	1.32	669	24.0	0.95	497	22.1	1.03
Newfoundland and Labrador	8,046	449.4	5.16	2,052	113.0	2.57	939	50.0	1.66	1,280	70.4	2.01	545	29.3	1.29	433	27.3	1.37
Northwest Territory	350	433.6	25.55	112	123.7	12.69	34	58.8	10.78	63	79.9	11.10	22	25.2	5.90	11	9.7	2.94
Nova Scotia	14,945	447.3	3.78	3,835	116.2	1.94	2,303	66.0	1.40	1,977	56.7	1.31	847	24.2	0.86	478	17.2	0.82
Prince Edward Island	2,105	418.2	9.39	605	118.4	4.94	287	55.1	3.29	278	54.6	3.35	99	19.3	2.01	41	10.1	1.64
Saskatchewan	13,313	406.6	3.62	3,713	115.6	1.95	1,885	56.2	1.32	1,771	51.6	1.26	801	24.4	0.88	357	12.8	0.70
NAACCR North American Combined	**3,946,051**	**417.7**	**0.21**	**1,153,821**	**123.3**	**0.12**	**515,051**	**52.8**	**0.07**	**343,101**	**35.4**	**0.06**	**251,922**	**25.8**	**0.05**	**175,611**	**20.9**	**0.05**

1. Rates are per 100,000 population and are age-adjusted by five-year age groups to the 2000 U.S. standard population based on single years of age.
2. See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.
3. Abbreviations: NOS - Not Otherwise Specified; S.E. - Standard Error

- Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for the combined.

Total Cancer Cases and Average Annual Age-adjusted (1996 Canadian Standard) Cancer Incidence Rates[1] for the Five Most Commonly Diagnosed Cancers
Selected Areas in the NAACCR North American Combined[2], 2010-2014, All Races, Females

	All Sites			Breast			Lung & Bronchus			Colon & Rectum			Uterus Corpus & Uterus, NOS[3]			Thyroid		
	Count	*Rate*	*S.E.*[3]	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
NAACCR U.S. Combined	**3,781,169**	**403.7**	**0.21**	**1,107,649**	**120.4**	**0.12**	**493,029**	**50.3**	**0.07**	**323,272**	**32.8**	**0.06**	**241,508**	**25.4**	**0.05**	**169,974**	**21.5**	**0.05**
Alabama	59,329	382.0	1.62	17,811	116.6	0.90	8,235	50.1	0.56	5,651	35.0	0.48	2,968	18.6	0.35	1,712	13.2	0.33
Alaska	6,568	397.8	5.14	2,085	122.0	2.80	773	50.6	1.91	620	38.3	1.61	431	24.2	1.23	299	17.9	1.07
Arizona	72,444	361.6	1.38	21,537	109.4	0.77	9,264	43.0	0.45	5,908	28.0	0.37	4,260	21.2	0.33	3,979	24.2	0.40
Arkansas	36,111	385.4	2.09	10,122	109.6	1.13	5,818	58.4	0.78	3,456	35.3	0.62	1,997	21.1	0.49	1,180	15.4	0.47
California	404,208	374.3	0.60	125,201	117.5	0.34	41,600	37.6	0.19	34,469	30.6	0.17	26,391	24.2	0.15	18,747	19.4	0.14
CA: Greater California	225,193	381.1	0.82	69,313	118.8	0.46	24,875	41.0	0.26	18,789	30.6	0.23	13,933	23.3	0.20	10,305	19.7	0.20
CA: Greater Bay Area	79,420	377.6	1.37	25,087	121.5	0.78	7,912	36.5	0.42	6,667	30.3	0.38	5,363	25.0	0.35	3,212	17.2	0.31
CA: Los Angeles	99,469	356.7	1.15	30,772	111.6	0.65	8,801	31.1	0.34	9,004	31.0	0.33	7,095	25.4	0.31	5,229	20.3	0.29
Colorado	54,644	376.2	1.65	17,541	120.4	0.93	5,751	39.3	0.53	4,259	28.8	0.45	3,321	21.8	0.39	3,154	24.0	0.44
Connecticut	52,718	442.6	2.02	15,767	135.6	1.12	6,767	54.0	0.68	4,159	32.1	0.53	3,483	28.6	0.50	2,827	29.6	0.59
Delaware	13,375	441.3	3.96	3,886	129.5	2.15	1,930	59.9	1.39	986	30.9	1.02	890	28.6	0.99	578	24.1	1.05
District of Columbia	7,606	431.5	5.08	2,389	139.6	2.93	845	47.0	1.65	690	38.3	1.50	549	30.4	1.32	396	23.0	1.19
Florida	266,096	383.3	0.78	75,212	112.5	0.43	38,674	50.7	0.26	22,825	30.5	0.21	15,788	22.7	0.19	9,412	17.9	0.19
Georgia	110,666	397.0	1.21	33,614	120.3	0.67	14,072	49.8	0.42	9,578	33.8	0.35	6,133	21.2	0.28	4,359	16.9	0.26
Hawaii	17,260	395.1	3.15	5,670	133.0	1.84	1,717	36.2	0.90	1,599	34.6	0.91	1,237	29.5	0.87	831	23.8	0.86
Idaho	18,042	396.2	3.03	5,295	117.1	1.65	2,117	44.5	0.98	1,437	30.5	0.83	1,111	23.7	0.73	980	25.5	0.84
Illinois	165,887	422.6	1.07	48,665	126.6	0.59	22,372	55.0	0.38	14,948	35.8	0.30	11,336	28.5	0.27	7,370	22.1	0.26
Indiana	82,508	411.5	1.48	23,084	117.1	0.79	12,158	58.3	0.54	7,666	36.2	0.43	5,390	26.4	0.37	3,015	18.2	0.34
Iowa	42,359	424.3	2.18	11,520	119.4	1.17	5,377	50.6	0.71	4,114	37.1	0.61	3,042	30.5	0.57	1,575	20.5	0.54
Kansas	35,882	416.3	2.29	10,182	120.3	1.24	4,596	50.5	0.76	3,129	34.1	0.64	2,104	24.2	0.54	1,833	25.7	0.62
Kentucky	63,000	457.5	1.88	16,413	120.3	0.97	11,053	76.6	0.74	5,774	40.1	0.54	3,521	25.3	0.44	2,526	22.1	0.46
Louisiana	55,904	407.2	1.77	16,330	120.1	0.97	7,513	52.6	0.62	5,551	39.4	0.54	2,610	18.8	0.38	2,371	19.6	0.42
Maine	21,056	438.7	3.19	5,743	122.4	1.70	3,189	62.3	1.13	1,638	32.2	0.84	1,506	30.3	0.81	822	22.8	0.85
Maryland	74,065	406.4	1.54	23,092	127.6	0.86	9,245	49.3	0.52	5,920	31.3	0.42	4,867	25.9	0.38	3,554	22.3	0.39
Massachusetts	94,120	434.0	1.47	28,058	132.7	0.82	13,038	57.6	0.52	7,338	31.5	0.39	6,255	28.4	0.37	5,322	29.7	0.42
Michigan	131,536	412.0	1.18	37,301	119.1	0.64	18,930	56.2	0.42	10,849	32.0	0.32	9,267	28.4	0.30	5,218	20.3	0.29
MI: Metropolitan Detroit	55,534	442.1	1.95	15,788	128.1	1.05	8,076	61.6	0.70	4,704	35.2	0.53	3,787	29.7	0.50	2,511	24.0	0.50
Mississippi	35,981	394.4	2.14	10,104	112.8	1.15	5,172	54.1	0.76	3,786	39.8	0.66	1,778	19.2	0.47	1,261	16.2	0.47
Missouri	79,928	413.2	1.52	23,218	122.6	0.83	12,540	61.5	0.56	7,077	34.4	0.42	4,997	25.6	0.37	2,866	18.1	0.35
Montana	13,450	410.6	3.72	3,893	119.7	2.02	1,791	51.0	1.23	1,117	32.2	1.01	860	25.3	0.90	561	22.1	0.99
Nebraska	22,633	402.0	2.79	6,538	118.6	1.53	2,834	48.1	0.93	2,170	35.8	0.80	1,470	26.1	0.70	975	21.4	0.71
New Hampshire	19,517	447.8	3.35	5,914	136.9	1.85	2,726	60.4	1.19	1,481	31.9	0.87	1,466	32.3	0.87	845	23.7	0.86
New Jersey	125,460	437.0	1.28	36,075	128.6	0.70	15,020	49.9	0.42	10,947	35.4	0.35	8,874	30.5	0.33	6,800	28.4	0.36
New York	273,599	435.3	0.86	76,791	125.6	0.47	34,081	51.6	0.29	22,514	33.5	0.23	19,341	30.4	0.22	15,178	28.8	0.24

Total Cancer Cases and Average Annual Age-adjusted (1996 Canadian Standard) Cancer Incidence Rates[1] for the Five Most Commonly Diagnosed Cancers
Selected Areas in the NAACCR North American Combined[2], 2010-2014, All Races, Females

	All Sites			Breast			Lung & Bronchus			Colon & Rectum			Uterus Corpus & Uterus, NOS[3]			Thyroid		
	Count	*Rate*	*S.E.*[3]	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
North Carolina	124,067	407.4	1.19	37,914	126.0	0.66	17,109	53.7	0.42	9,839	31.2	0.32	7,459	23.9	0.28	4,854	18.6	0.27
North Dakota	8,513	401.9	4.61	2,397	118.1	2.54	1,048	47.1	1.50	872	37.8	1.35	527	24.9	1.13	426	24.9	1.27
Ohio	154,863	410.5	1.09	44,328	119.6	0.59	22,438	56.4	0.39	13,624	33.6	0.30	10,697	27.8	0.28	6,693	22.2	0.28
Oklahoma	46,231	399.2	1.91	13,103	114.7	1.03	6,873	56.3	0.69	4,236	35.0	0.55	2,496	21.3	0.44	1,731	17.8	0.44
Oregon	50,591	404.8	1.87	15,188	122.7	1.03	6,696	50.6	0.63	3,881	29.2	0.49	3,342	26.0	0.46	2,010	20.0	0.46
Pennsylvania	195,598	446.0	1.06	53,851	126.4	0.57	25,224	53.6	0.35	16,964	35.2	0.29	14,004	31.5	0.28	10,515	31.1	0.32
Rhode Island	15,625	443.3	3.74	4,331	127.1	2.02	2,265	60.7	1.32	1,271	32.5	0.97	1,117	31.3	0.97	784	27.7	1.04
South Carolina	60,486	398.2	1.67	18,597	123.8	0.94	8,268	51.3	0.57	5,051	32.0	0.46	3,505	22.3	0.39	2,222	17.5	0.39
South Dakota	10,661	413.3	4.23	3,167	127.4	2.38	1,324	48.4	1.37	1,041	36.3	1.19	708	27.5	1.08	412	20.2	1.04
Tennessee	82,826	405.5	1.45	23,909	118.2	0.79	12,555	58.6	0.53	7,307	34.5	0.41	4,633	22.3	0.34	3,231	18.9	0.34
Texas	251,958	367.0	0.74	74,269	108.6	0.40	29,302	42.4	0.25	21,373	30.4	0.21	14,447	20.8	0.18	11,944	18.6	0.17
Utah	23,175	362.2	2.41	7,012	111.5	1.35	1,449	22.9	0.61	1,686	26.2	0.65	1,643	25.5	0.64	1,856	29.5	0.69
Vermont	9,030	422.3	4.68	2,662	126.9	2.58	1,285	57.0	1.63	710	30.5	1.20	710	32.0	1.25	316	19.7	1.19
Virginia	94,849	384.8	1.28	30,271	123.6	0.73	12,257	48.3	0.44	7,934	31.2	0.36	5,915	23.3	0.31	4,277	20.0	0.31
Washington	86,489	421.8	1.48	26,686	131.2	0.83	10,713	50.7	0.50	6,534	30.9	0.39	5,342	25.0	0.35	3,679	20.7	0.35
WA: Seattle/Puget Sound	60,793	433.8	1.81	18,947	136.2	1.02	7,271	50.8	0.61	4,519	31.4	0.48	3,756	25.7	0.43	2,586	20.9	0.42
West Virginia	27,673	430.3	2.72	7,112	111.9	1.39	4,416	63.5	0.98	2,664	38.5	0.78	1,935	30.2	0.72	1,226	24.8	0.75
Wisconsin	76,465	422.5	1.59	21,974	124.7	0.87	9,884	52.5	0.54	6,139	31.3	0.42	5,428	29.2	0.41	2,947	20.1	0.39
Wyoming	6,117	371.2	4.92	1,827	110.4	2.68	725	42.0	1.60	490	29.0	1.35	357	21.1	1.16	305	21.9	1.31
NAACCR Canadian Combined	**164,882**	**394.9**	**1.00**	**46,172**	**112.4**	**0.54**	**22,022**	**51.1**	**0.35**	**19,829**	**45.1**	**0.33**	**10,414**	**24.6**	**0.25**	**5,637**	**16.0**	**0.22**
Alberta	40,556	392.1	1.99	11,844	114.7	1.07	5,104	50.2	0.72	4,317	40.8	0.63	2,597	24.5	0.49	1,781	18.1	0.44
British Columbia	58,176	377.8	1.62	16,780	111.6	0.89	7,566	46.8	0.55	6,916	42.1	0.52	3,635	23.3	0.40	1,500	11.9	0.32
Manitoba	16,333	415.8	3.38	4,342	113.6	1.78	2,310	56.6	1.21	1,940	46.3	1.09	1,199	31.0	0.92	539	16.7	0.75
New Brunswick	11,058	401.6	3.99	2,889	106.2	2.06	1,594	54.7	1.39	1,287	43.5	1.25	669	23.5	0.93	497	22.3	1.06
Newfoundland and Labrador	8,046	434.1	5.02	2,052	109.8	2.51	939	48.0	1.59	1,280	66.5	1.90	545	28.6	1.26	433	27.8	1.42
Northwest Territory	350	418.1	24.31	112	120.7	12.31	34	55.3	10.01	63	76.3	10.44	22	24.5	5.70	11	9.9	3.04
Nova Scotia	14,945	430.1	3.68	3,835	113.2	1.91	2,303	62.7	1.33	1,977	53.4	1.24	847	23.8	0.85	478	17.5	0.85
Prince Edward Island	2,105	401.2	9.10	605	114.4	4.81	287	52.6	3.15	278	51.1	3.18	99	18.9	1.97	41	10.2	1.68
Saskatchewan	13,313	392.2	3.55	3,713	112.3	1.92	1,885	54.0	1.28	1,771	48.8	1.22	801	23.8	0.87	357	13.0	0.72
NAACCR North American Combined	**3,946,051**	**403.3**	**0.21**	**1,153,821**	**120.0**	**0.11**	**515,051**	**50.3**	**0.07**	**343,101**	**33.3**	**0.06**	**251,922**	**25.3**	**0.05**	**175,611**	**21.3**	**0.05**

1. Rates are per 100,000 and are age-adjusted to the 1996 Canadian population standard.
2. See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.
3. Abbreviations: NOS - Not Otherwise Specified; S.E. - Standard Error

\- Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for the combined.

NAACCR U.S. Combined Pediatric Cancer Incidence
2010-2014

Total Cancer Cases and Average Annual Age-specific Incidence Rates[1] for Pediatric Cancers
NAACCR United States Combined[2], 2010-2014, All Races, Males

	Ages 0-4			Ages 5-9			Ages 10-14			Ages 15-19		
Cancer Types	*Count*	*Rate*	*S.E.*[3]	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
All Sites	**12,245**	**248.0**	**2.24**	**7,095**	**140.6**	**1.67**	**7,514**	**147.1**	**1.70**	**12,217**	**229.4**	**2.08**
I. Leukemias	4,223	85.5	1.32	2,393	47.4	0.97	1,848	36.2	0.84	2,122	39.8	0.86
I(a). Lymphoid leukemias	3,259	66.0	1.16	1,970	39.0	0.88	1,237	24.2	0.69	1,162	21.8	0.64
I(b). Acute myeloid leukemias	585	11.8	0.49	235	4.7	0.30	356	7.0	0.37	537	10.1	0.44
II. Lymphomas & reticuloendothelial neoplasms	1,023	20.7	0.65	1,261	25.0	0.70	1,654	32.4	0.80	2,998	56.3	1.03
II(a). Hodgkin lymphomas	47	1.0	0.14	289	5.7	0.34	655	12.8	0.50	1,640	30.8	0.76
II(b, c, e). Non-Hodgkin lymphomas	430	8.7	0.42	728	14.4	0.53	824	16.1	0.56	1,253	23.5	0.66
III. CNS[3] & misc. intracranial & intraspinal neoplasms	2,205	44.6	0.95	1,861	36.9	0.86	1,521	29.8	0.76	1,301	24.4	0.68
III(a). Ependymomas and choroids plexus tumor	310	6.3	0.36	110	2.2	0.21	103	2.0	0.20	86	1.6	0.17
III(b). Astrocytomas	979	19.8	0.63	928	18.4	0.60	823	16.1	0.56	759	14.3	0.52
III(c). Intracranial and intraspinal embryonal tumors	586	11.9	0.49	431	8.5	0.41	259	5.1	0.31	155	2.9	0.23
III(d). Other gliomas	254	5.1	0.32	341	6.8	0.37	282	5.5	0.33	240	4.5	0.29
IV. Neuroblastoma and other peripheral nervous cell tumors	1,491	30.2	0.78	217	4.3	0.29	81	1.6	0.18	51	1.0	0.13
IV(a). Neuroblastoma & ganglioneuroblastoma	1,482	30.0	0.78	210	4.2	0.29	59	1.2	0.15	24	0.5	0.09
V. Retinoblastoma	603	12.2	0.50	27	0.5	0.10	6	0.1	0.05	-	-	-
VI. Renal Tumors	908	18.4	0.61	248	4.9	0.31	66	1.3	0.16	97	1.8	0.18
VI(a). Nephroblastoma and other nonepithelial renal tumors	897	18.2	0.61	227	4.5	0.30	31	0.6	0.11	19	0.4	0.08
VII. Hepatic tumors	481	9.7	0.44	39	0.8	0.12	52	1.0	0.14	53	1.0	0.14
VII(a). Hepatoblastoma	460	9.3	0.43	24	0.5	0.10	13	0.3	0.07	-	-	-
VIII. Malignant bone tumors	78	1.6	0.18	300	5.9	0.34	759	14.9	0.54	882	16.6	0.56
VIII(a). Osteosarcomas	16	0.3	0.08	144	2.9	0.24	455	8.9	0.42	511	9.6	0.42
VIII(c). Ewing tumor and related sarcomas of bone	51	1.0	0.14	134	2.7	0.23	237	4.6	0.30	267	5.0	0.31
IX. Soft tissue and other extraosseous sarcomas	721	14.6	0.54	468	9.3	0.43	605	11.8	0.48	858	16.1	0.55
IX(a). Rhabdomyosarcomas	422	8.5	0.42	256	5.1	0.32	194	3.8	0.27	197	3.7	0.26
X. Germ cell & trophoblastic tumors & neoplasms of gonads	317	6.4	0.36	79	1.6	0.18	297	5.8	0.34	2,192	41.2	0.88
X(a). Intracranial & intraspinal germ cell tumors	32	0.6	0.11	67	1.3	0.16	207	4.1	0.28	213	4.0	0.27
X(c). Malignant gonadal germ cell tumors	163	3.3	0.26	-	-	-	65	1.3	0.16	1,834	34.4	0.80
XI. Other malignant epithelial neoplasms and melanomas	127	2.6	0.23	192	3.8	0.27	604	11.8	0.48	1,636	30.7	0.76
XI(b). Thyroid carcinomas	9	0.2	0.06	46	0.9	0.13	204	4.0	0.28	490	9.2	0.42
XI(d). Malignant melanomas	69	1.4	0.17	77	1.5	0.17	130	2.5	0.22	438	8.2	0.39
XII. Other & unspecified malignant neoplasms	68	1.4	0.17	10	0.2	0.06	21	0.4	0.09	27	0.5	0.10

[1] Rates are **per million** population.
[2] See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.
[3] Abbreviations: S.E. - Standard Error; CNS - Central Nervous System.
- Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for All Sites combined.

Total Cancer Cases and Average Annual Age-specific Incidence Rates[1] for Pediatric Cancers
NAACCR United States Combined[2], 2010-2014, All Races, Females

	Ages 0-4			Ages 5-9			Ages 10-14			Ages 15-19		
Cancer Types	*Count*	*Rate*	*S.E.*[3]	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
All Sites	**10,674**	**225.8**	**2.19**	**5,782**	**119.6**	**1.57**	**6,829**	**139.6**	**1.69**	**10,938**	**216.1**	**2.07**
I. Leukemias	3,740	79.1	1.29	1,874	38.8	0.90	1,418	29.0	0.77	1,394	27.5	0.74
I(a). Lymphoid leukemias	2,877	60.9	1.13	1,510	31.2	0.80	879	18.0	0.61	541	10.7	0.46
I(b). Acute myeloid leukemias	545	11.5	0.49	217	4.5	0.30	325	6.6	0.37	511	10.1	0.45
II. Lymphomas & reticuloendothelial neoplasms	622	13.2	0.53	598	12.4	0.51	1,070	21.9	0.67	2,310	45.6	0.95
II(a). Hodgkin lymphomas	21	0.4	0.10	87	1.8	0.19	497	10.2	0.46	1,616	31.9	0.79
II(b, c, e). Non-Hodgkin lymphomas	215	4.5	0.31	323	6.7	0.37	448	9.2	0.43	631	12.5	0.50
III. CNS[3] & misc. intracranial & intraspinal neoplasms	1,966	41.6	0.94	1,588	32.8	0.82	1,319	27.0	0.74	1,067	21.1	0.65
III(a). Ependymomas and choroids plexus tumor	230	4.9	0.32	113	2.3	0.22	68	1.4	0.17	72	1.4	0.17
III(b). Astrocytomas	993	21.0	0.67	848	17.5	0.60	777	15.9	0.57	628	12.4	0.50
III(c). Intracranial and intraspinal embryonal tumors	444	9.4	0.45	232	4.8	0.32	155	3.2	0.25	92	1.8	0.19
III(d). Other gliomas	241	5.1	0.33	344	7.1	0.38	261	5.3	0.33	232	4.6	0.30
IV. Neuroblastoma and other peripheral nervous cell tumors	1,354	28.6	0.78	195	4.0	0.29	71	1.5	0.17	33	0.7	0.11
IV(a). Neuroblastoma & ganglioneuroblastoma	1,349	28.5	0.78	187	3.9	0.28	57	1.2	0.15	19	0.4	0.09
V. Retinoblastoma	642	13.6	0.54	25	0.5	0.10	-	-	-	-	-	-
VI. Renal Tumors	975	20.6	0.66	334	6.9	0.38	81	1.7	0.18	107	2.1	0.20
VI(a). Nephroblastoma and other nonepithelial renal tumors	966	20.4	0.66	320	6.6	0.37	46	0.9	0.14	22	0.4	0.09
VII. Hepatic tumors	257	5.4	0.34	42	0.9	0.13	43	0.9	0.13	68	1.3	0.16
VII(a). Hepatoblastoma	248	5.2	0.33	22	0.5	0.10	11	0.2	0.07	-	-	-
VIII. Malignant bone tumors	90	1.9	0.20	282	5.8	0.35	655	13.4	0.52	551	10.9	0.46
VIII(a). Osteosarcomas	13	0.3	0.08	157	3.2	0.26	411	8.4	0.41	309	6.1	0.35
VIII(c). Ewing tumor and related sarcomas of bone	60	1.3	0.16	96	2.0	0.20	193	3.9	0.28	150	3.0	0.24
IX. Soft tissue and other extraosseous sarcomas	535	11.3	0.49	380	7.9	0.40	596	12.2	0.50	688	13.6	0.52
IX(a). Rhabdomyosarcomas	315	6.7	0.38	189	3.9	0.28	178	3.6	0.27	156	3.1	0.25
X. Germ cell & trophoblastic tumors & neoplasms of gonads	306	6.5	0.37	189	3.9	0.28	454	9.3	0.44	741	14.6	0.54
X(a). Intracranial & intraspinal germ cell tumors	29	0.6	0.11	56	1.2	0.15	74	1.5	0.18	28	0.6	0.10
X(c). Malignant gonadal germ cell tumors	29	0.6	0.11	109	2.3	0.22	310	6.3	0.36	435	8.6	0.41
XI. Other malignant epithelial neoplasms and melanomas	136	2.9	0.25	264	5.5	0.34	1,085	22.2	0.67	3,898	77.0	1.23
XI(b). Thyroid carcinomas	14	0.3	0.08	96	2.0	0.20	625	12.8	0.51	2,391	47.2	0.97
XI(d). Malignant melanomas	88	1.9	0.20	97	2.0	0.20	147	3.0	0.25	653	12.9	0.50
XII. Other & unspecified malignant neoplasms	51	1.1	0.15	11	0.2	0.07	36	0.7	0.12	80	1.6	0.18

[1] Rates are **per million** population.
[2] See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.
[3] Abbreviations: S.E. - Standard Error; CNS - Central Nervous System.
- Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for All Sites combined.

Total Cancer Cases and Average Annual Age-specific Incidence Rates[1] for Pediatric Cancers
NAACCR United States Combined[2], 2010-2014, White, Males

	Ages 0-4			Ages 5-9			Ages 10-14			Ages 15-19		
Cancer Types	*Count*	*Rate*	*S.E.*[3]	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
All Sites	**9,602**	**259.1**	**2.64**	**5,600**	**146.6**	**1.96**	**5,871**	**151.0**	**1.97**	**9,902**	**245.2**	**2.46**
I. Leukemias	3,359	90.6	1.56	1,933	50.6	1.15	1,428	36.7	0.97	1,704	42.2	1.02
I(a). Lymphoid leukemias	2,636	71.1	1.39	1,627	42.6	1.06	987	25.4	0.81	971	24.0	0.77
I(b). Acute myeloid leukemias	426	11.5	0.56	171	4.5	0.34	263	6.8	0.42	420	10.4	0.51
II. Lymphomas & reticuloendothelial neoplasms	804	21.7	0.77	977	25.6	0.82	1,266	32.6	0.92	2,374	58.8	1.21
II(a). Hodgkin lymphomas	33	0.9	0.16	206	5.4	0.38	514	13.2	0.58	1,355	33.6	0.91
II(b, c, e). Non-Hodgkin lymphomas	324	8.7	0.49	566	14.8	0.62	601	15.5	0.63	928	23.0	0.75
III. CNS[3] & misc. intracranial & intraspinal neoplasms	1,754	47.3	1.13	1,474	38.6	1.00	1,231	31.7	0.90	1,040	25.8	0.80
III(a). Ependymomas and choroids plexus tumor	243	6.6	0.42	91	2.4	0.25	85	2.2	0.24	64	1.6	0.20
III(b). Astrocytomas	785	21.2	0.76	749	19.6	0.72	674	17.3	0.67	603	14.9	0.61
III(c). Intracranial and intraspinal embryonal tumors	467	12.6	0.58	345	9.0	0.49	214	5.5	0.38	129	3.2	0.28
III(d). Other gliomas	201	5.4	0.38	255	6.7	0.42	222	5.7	0.38	194	4.8	0.34
IV. Neuroblastoma and other peripheral nervous cell tumors	1,188	32.1	0.93	165	4.3	0.34	60	1.5	0.20	37	0.9	0.15
IV(a). Neuroblastoma & ganglioneuroblastoma	1,181	31.9	0.93	159	4.2	0.33	43	1.1	0.17	19	0.5	0.11
V. Retinoblastoma	447	12.1	0.57	17	0.4	0.11	6	0.2	0.06	-	-	-
VI. Renal Tumors	687	18.5	0.71	186	4.9	0.36	47	1.2	0.18	65	1.6	0.20
VI(a). Nephroblastoma and other nonepithelial renal tumors	682	18.4	0.70	171	4.5	0.34	24	0.6	0.13	11	0.3	0.08
VII. Hepatic tumors	367	9.9	0.52	28	0.7	0.14	43	1.1	0.17	38	0.9	0.15
VII(a). Hepatoblastoma	350	9.4	0.50	16	0.4	0.10	9	0.2	0.08	-	-	-
VIII. Malignant bone tumors	61	1.6	0.21	248	6.5	0.41	597	15.4	0.63	709	17.6	0.66
VIII(a). Osteosarcomas	10	0.3	0.09	105	2.7	0.27	333	8.6	0.47	393	9.7	0.49
VIII(c). Ewing tumor and related sarcomas of bone	46	1.2	0.18	124	3.2	0.29	214	5.5	0.38	239	5.9	0.38
IX. Soft tissue and other extraosseous sarcomas	544	14.7	0.63	346	9.1	0.49	454	11.7	0.55	630	15.6	0.62
IX(a). Rhabdomyosarcomas	321	8.7	0.48	190	5.0	0.36	143	3.7	0.31	139	3.4	0.29
X. Germ cell & trophoblastic tumors & neoplasms of gonads	239	6.4	0.42	63	1.6	0.21	243	6.2	0.40	1,944	48.1	1.09
X(a). Intracranial & intraspinal germ cell tumors	25	0.7	0.13	54	1.4	0.19	163	4.2	0.33	159	3.9	0.31
X(c). Malignant gonadal germ cell tumors	124	3.3	0.30	-	-	-	59	1.5	0.20	1,666	41.3	1.01
XI. Other malignant epithelial neoplasms and melanomas	94	2.5	0.26	156	4.1	0.33	478	12.3	0.56	1,342	33.2	0.91
XI(b). Thyroid carcinomas	6	0.2	0.07	40	1.0	0.17	167	4.3	0.33	412	10.2	0.50
XI(d). Malignant melanomas	58	1.6	0.21	67	1.8	0.21	117	3.0	0.28	408	10.1	0.50
XII. Other & unspecified malignant neoplasms	58	1.6	0.21	7	0.2	0.07	18	0.5	0.11	19	0.5	0.11

[1] Rates are **per million** population.
[2] See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.
[3] Abbreviations: S.E. - Standard Error; CNS - Central Nervous System.
- Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for All Sites combined.

Total Cancer Cases and Average Annual Age-specific Incidence Rates[1] for Pediatric Cancers
NAACCR United States Combined[2], 2010-2014, White, Females

	Ages 0-4			Ages 5-9			Ages 10-14			Ages 15-19		
Cancer Types	*Count*	*Rate*	*S.E.*[3]	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
All Sites	**8,408**	**237.6**	**2.59**	**4,465**	**122.5**	**1.83**	**5,389**	**145.4**	**1.98**	**8,744**	**228.9**	**2.45**
I. Leukemias	3,038	85.9	1.56	1,502	41.2	1.06	1,135	30.6	0.91	1,078	28.2	0.86
I(a). Lymphoid leukemias	2,389	67.5	1.38	1,236	33.9	0.96	731	19.7	0.73	449	11.8	0.55
I(b). Acute myeloid leukemias	410	11.6	0.57	159	4.4	0.35	241	6.5	0.42	380	9.9	0.51
II. Lymphomas & reticuloendothelial neoplasms	499	14.1	0.63	463	12.7	0.59	844	22.8	0.78	1,834	48.0	1.12
II(a). Hodgkin lymphomas	14	0.4	0.11	70	1.9	0.23	388	10.5	0.53	1,319	34.5	0.95
II(b, c, e). Non-Hodgkin lymphomas	165	4.7	0.36	235	6.4	0.42	341	9.2	0.50	457	12.0	0.56
III. CNS[3] & misc. intracranial & intraspinal neoplasms	1,568	44.3	1.12	1,232	33.8	0.96	1,052	28.4	0.88	852	22.3	0.76
III(a). Ependymomas and choroids plexus tumor	184	5.2	0.38	84	2.3	0.25	58	1.6	0.21	55	1.4	0.19
III(b). Astrocytomas	801	22.6	0.80	678	18.6	0.71	628	16.9	0.68	512	13.4	0.59
III(c). Intracranial and intraspinal embryonal tumors	350	9.9	0.53	178	4.9	0.37	121	3.3	0.30	73	1.9	0.22
III(d). Other gliomas	192	5.4	0.39	259	7.1	0.44	207	5.6	0.39	181	4.7	0.35
IV. Neuroblastoma and other peripheral nervous cell tumors	1,055	29.8	0.92	133	3.6	0.32	60	1.6	0.21	22	0.6	0.12
IV(a). Neuroblastoma & ganglioneuroblastoma	1,051	29.7	0.92	128	3.5	0.31	48	1.3	0.19	13	0.3	0.09
V. Retinoblastoma	454	12.8	0.60	17	0.5	0.11	-	-	-	-	-	-
VI. Renal Tumors	729	20.6	0.76	228	6.3	0.41	62	1.7	0.21	84	2.2	0.24
VI(a). Nephroblastoma and other nonepithelial renal tumors	723	20.4	0.76	221	6.1	0.41	38	1.0	0.17	19	0.5	0.11
VII. Hepatic tumors	206	5.8	0.41	28	0.8	0.15	31	0.8	0.15	49	1.3	0.18
VII(a). Hepatoblastoma	198	5.6	0.40	16	0.4	0.11	7	0.2	0.07	-	-	-
VIII. Malignant bone tumors	75	2.1	0.24	216	5.9	0.40	520	14.0	0.62	429	11.2	0.54
VIII(a). Osteosarcomas	10	0.3	0.09	109	3.0	0.29	301	8.1	0.47	224	5.9	0.39
VIII(c). Ewing tumor and related sarcomas of bone	48	1.4	0.20	84	2.3	0.25	179	4.8	0.36	134	3.5	0.30
IX. Soft tissue and other extraosseous sarcomas	416	11.8	0.58	284	7.8	0.46	435	11.7	0.56	507	13.3	0.59
IX(a). Rhabdomyosarcomas	239	6.8	0.44	142	3.9	0.33	131	3.5	0.31	115	3.0	0.28
X. Germ cell & trophoblastic tumors & neoplasms of gonads	224	6.3	0.42	133	3.6	0.32	323	8.7	0.49	532	13.9	0.60
X(a). Intracranial & intraspinal germ cell tumors	24	0.7	0.14	45	1.2	0.18	51	1.4	0.19	20	0.5	0.12
X(c). Malignant gonadal germ cell tumors	22	0.6	0.13	72	2.0	0.23	217	5.9	0.40	312	8.2	0.46
XI. Other malignant epithelial neoplasms and melanomas	111	3.1	0.30	221	6.1	0.41	903	24.4	0.81	3,297	86.3	1.50
XI(b). Thyroid carcinomas	11	0.3	0.09	79	2.2	0.24	533	14.4	0.62	2,041	53.4	1.18
XI(d). Malignant melanomas	74	2.1	0.24	89	2.4	0.26	131	3.5	0.31	611	16.0	0.65
XII. Other & unspecified malignant neoplasms	33	0.9	0.16	8	0.2	0.08	24	0.6	0.13	59	1.5	0.20

[1] Rates are **per million** population.
[2] See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.
[3] Abbreviations: S.E. - Standard Error; CNS - Central Nervous System.
- Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for All Sites combined.

Total Cancer Cases and Average Annual Age-specific Incidence Rates[1] for Pediatric Cancers
NAACCR United States Combined[2], 2010-2014, Black, Males

	Ages 0-4			Ages 5-9			Ages 10-14			Ages 15-19		
Cancer Types	*Count*	*Rate*	*S.E.[3]*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
All Sites	**1,459**	**172.4**	**4.51**	**902**	**107.9**	**3.59**	**1,040**	**122.6**	**3.80**	**1,382**	**153.1**	**4.12**
I. Leukemias	413	48.8	2.40	259	31.0	1.93	251	29.6	1.87	253	28.0	1.76
I(a). Lymphoid leukemias	280	33.1	1.98	194	23.2	1.67	141	16.6	1.40	105	11.6	1.14
I(b). Acute myeloid leukemias	82	9.7	1.07	31	3.7	0.67	60	7.1	0.91	72	8.0	0.94
II. Lymphomas & reticuloendothelial neoplasms	96	11.3	1.16	160	19.1	1.51	258	30.4	1.89	411	45.5	2.25
II(a). Hodgkin lymphomas	9	1.1	0.35	59	7.1	0.92	100	11.8	1.18	201	22.3	1.57
II(b, c, e). Non-Hodgkin lymphomas	51	6.0	0.84	86	10.3	1.11	147	17.3	1.43	206	22.8	1.59
III. CNS[3] & misc. intracranial & intraspinal neoplasms	270	31.9	1.94	239	28.6	1.85	198	23.3	1.66	174	19.3	1.46
III(a). Ependymomas and choroids plexus tumor	39	4.6	0.74	8	1.0	0.34	13	1.5	0.42	14	1.6	0.41
III(b). Astrocytomas	116	13.7	1.27	115	13.8	1.28	102	12.0	1.19	96	10.6	1.09
III(c). Intracranial and intraspinal embryonal tumors	71	8.4	1.00	50	6.0	0.85	30	3.5	0.65	20	2.2	0.50
III(d). Other gliomas	35	4.1	0.70	55	6.6	0.89	40	4.7	0.75	35	3.9	0.66
IV. Neuroblastoma and other peripheral nervous cell tumors	182	21.5	1.59	38	4.5	0.74	17	2.0	0.49	10	1.1	0.35
IV(a). Neuroblastoma & ganglioneuroblastoma	180	21.3	1.59	37	4.4	0.73	12	1.4	0.41	-	-	-
V. Retinoblastoma	87	10.3	1.10	6	0.7	0.29	-	-	-	-	-	-
VI. Renal Tumors	169	20.0	1.54	48	5.7	0.83	15	1.8	0.46	21	2.3	0.51
VI(a). Nephroblastoma and other nonepithelial renal tumors	165	19.5	1.52	44	5.3	0.79	-	-	-	-	-	-
VII. Hepatic tumors	57	6.7	0.89	-	-	-	6	0.7	0.29	9	1.0	0.33
VII(a). Hepatoblastoma	55	6.5	0.88	-	-	-	-	-	-	-	-	-
VIII. Malignant bone tumors	6	0.7	0.29	30	3.6	0.66	106	12.5	1.21	112	12.4	1.17
VIII(a). Osteosarcomas	-	-	-	26	3.1	0.61	87	10.3	1.10	85	9.4	1.02
VIII(c). Ewing tumor and related sarcomas of bone	-	-	-	-	-	-	6	0.7	0.29	9	1.0	0.33
IX. Soft tissue and other extraosseous sarcomas	127	15.0	1.33	86	10.3	1.11	101	11.9	1.18	157	17.4	1.39
IX(a). Rhabdomyosarcomas	79	9.3	1.05	48	5.7	0.83	33	3.9	0.68	43	4.8	0.73
X. Germ cell & trophoblastic tumors & neoplasms of gonads	24	2.8	0.58	-	-	-	20	2.4	0.53	75	8.3	0.96
X(a). Intracranial & intraspinal germ cell tumors	-	-	-	-	-	-	18	2.1	0.50	23	2.5	0.53
X(c). Malignant gonadal germ cell tumors	10	1.2	0.37	-	-	-	-	-	-	42	4.7	0.72
XI. Other malignant epithelial neoplasms and melanomas	21	2.5	0.54	23	2.8	0.57	67	7.9	0.96	152	16.8	1.37
XI(b). Thyroid carcinomas	-	-	-	-	-	-	19	2.2	0.51	31	3.4	0.62
XI(d). Malignant melanomas	7	0.8	0.31	-	-	-	-	-	-	-	-	-
XII. Other & unspecified malignant neoplasms	7	0.8	0.31	-	-	-	-	-	-	8	0.9	0.31

[1] Rates are **per million** population.
[2] See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.
[3] Abbreviations: S.E. - Standard Error; CNS - Central Nervous System.
- Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for All Sites combined.

Total Cancer Cases and Average Annual Age-specific Incidence Rates[1] for Pediatric Cancers
NAACCR United States Combined[2], 2010-2014, Black, Females

	Ages 0-4			Ages 5-9			Ages 10-14			Ages 15-19		
Cancer Types	*Count*	*Rate*	*S.E.[3]*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
All Sites	**1,305**	**159.6**	**4.42**	**806**	**99.6**	**3.51**	**900**	**109.7**	**3.66**	**1,334**	**153.0**	**4.19**
I. Leukemias	342	41.8	2.26	187	23.1	1.69	157	19.1	1.53	184	21.1	1.56
I(a). Lymphoid leukemias	224	27.4	1.83	138	17.1	1.45	78	9.5	1.08	51	5.8	0.82
I(b). Acute myeloid leukemias	76	9.3	1.07	29	3.6	0.67	47	5.7	0.84	77	8.8	1.01
II. Lymphomas & reticuloendothelial neoplasms	52	6.4	0.88	72	8.9	1.05	157	19.1	1.53	321	36.8	2.05
II(a). Hodgkin lymphomas	-	-	-	12	1.5	0.43	85	10.4	1.12	210	24.1	1.66
II(b, c, e). Non-Hodgkin lymphomas	27	3.3	0.64	51	6.3	0.88	68	8.3	1.01	111	12.7	1.21
III. CNS[3] & misc. intracranial & intraspinal neoplasms	253	30.9	1.94	238	29.4	1.91	172	21.0	1.60	136	15.6	1.34
III(a). Ependymomas and choroids plexus tumor	32	3.9	0.69	17	2.1	0.51	8	1.0	0.34	10	1.1	0.36
III(b). Astrocytomas	111	13.6	1.29	115	14.2	1.33	99	12.1	1.21	74	8.5	0.99
III(c). Intracranial and intraspinal embryonal tumors	62	7.6	0.96	26	3.2	0.63	18	2.2	0.52	15	1.7	0.44
III(d). Other gliomas	36	4.4	0.73	67	8.3	1.01	34	4.1	0.71	29	3.3	0.62
IV. Neuroblastoma and other peripheral nervous cell tumors	184	22.5	1.66	47	5.8	0.85	8	1.0	0.34	-	-	-
IV(a). Neuroblastoma & ganglioneuroblastoma	183	22.4	1.65	44	5.4	0.82	7	0.9	0.32	-	-	-
V. Retinoblastoma	116	14.2	1.32	-	-	-	-	-	-	-	-	-
VI. Renal Tumors	183	22.4	1.65	86	10.6	1.15	17	2.1	0.50	20	2.3	0.51
VI(a). Nephroblastoma and other nonepithelial renal tumors	183	22.4	1.65	80	9.9	1.11	7	0.9	0.32	-	-	-
VII. Hepatic tumors	26	3.2	0.62	8	1.0	0.35	9	1.1	0.37	11	1.3	0.38
VII(a). Hepatoblastoma	26	3.2	0.62	-	-	-	-	-	-	-	-	-
VIII. Malignant bone tumors	-	-	-	45	5.6	0.83	92	11.2	1.17	79	9.1	1.02
VIII(a). Osteosarcomas	-	-	-	36	4.4	0.74	77	9.4	1.07	57	6.5	0.87
VIII(c). Ewing tumor and related sarcomas of bone	-	-	-	-	-	-	6	0.7	0.30	7	0.8	0.30
IX. Soft tissue and other extraosseous sarcomas	70	8.6	1.02	61	7.5	0.97	111	13.5	1.28	127	14.6	1.29
IX(a). Rhabdomyosarcomas	44	5.4	0.81	26	3.2	0.63	32	3.9	0.69	30	3.4	0.63
X. Germ cell & trophoblastic tumors & neoplasms of gonads	46	5.6	0.83	34	4.2	0.72	79	9.6	1.08	136	15.6	1.34
X(a). Intracranial & intraspinal germ cell tumors	-	-	-	-	-	-	13	1.6	0.44	-	-	-
X(c). Malignant gonadal germ cell tumors	-	-	-	28	3.5	0.65	55	6.7	0.90	79	9.1	1.02
XI. Other malignant epithelial neoplasms and melanomas	13	1.6	0.44	21	2.6	0.57	89	10.8	1.15	297	34.1	1.98
XI(b). Thyroid carcinomas	-	-	-	8	1.0	0.35	35	4.3	0.72	142	16.3	1.37
XI(d). Malignant melanomas	7	0.9	0.32	-	-	-	8	1.0	0.34	6	0.7	0.28
XII. Other & unspecified malignant neoplasms	15	1.8	0.47	-	-	-	8	1.0	0.34	18	2.1	0.49

[1] Rates are **per million** population.
[2] See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.
[3] Abbreviations: S.E. - Standard Error; CNS - Central Nervous System.
- Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for All Sites combined.

Total Cancer Cases and Average Annual Age-specific Incidence Rates[1] for Pediatric Cancers
NAACCR United States Combined[2], 2010-2014, Asian/Pacific Islander, Males

	Ages 0-4			Ages 5-9			Ages 10-14			Ages 15-19		
Cancer Types	*Count*	*Rate*	*S.E.*[3]	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
All Sites	**615**	**206.8**	**8.34**	**325**	**108.9**	**6.04**	**352**	**123.4**	**6.58**	**520**	**175.9**	**7.71**
I. Leukemias	239	80.4	5.20	115	38.5	3.59	97	34.0	3.45	108	36.5	3.52
I(a). Lymphoid leukemias	177	59.5	4.47	87	29.1	3.12	60	21.0	2.72	55	18.6	2.51
I(b). Acute myeloid leukemias	46	15.5	2.28	17	5.7	1.38	22	7.7	1.64	31	10.5	1.88
II. Lymphomas & reticuloendothelial neoplasms	78	26.2	2.97	60	20.1	2.59	72	25.2	2.97	121	40.9	3.72
II(a). Hodgkin lymphomas	-	-	-	17	5.7	1.38	26	9.1	1.79	48	16.2	2.34
II(b, c, e). Non-Hodgkin lymphomas	36	12.1	2.02	34	11.4	1.95	41	14.4	2.24	69	23.3	2.81
III. CNS[3] & misc. intracranial & intraspinal neoplasms	83	27.9	3.06	83	27.8	3.05	59	20.7	2.69	38	12.9	2.09
III(a). Ependymomas and choroids plexus tumor	21	7.1	1.54	-	-	-	-	-	-	6	2.0	0.83
III(b). Astrocytomas	24	8.1	1.65	35	11.7	1.98	33	11.6	2.01	26	8.8	1.72
III(c). Intracranial and intraspinal embryonal tumors	27	9.1	1.75	23	7.7	1.61	12	4.2	1.21	-	-	-
III(d). Other gliomas	8	2.7	0.95	19	6.4	1.46	10	3.5	1.11	-	-	-
IV. Neuroblastoma and other peripheral nervous cell tumors	72	24.2	2.85	10	3.3	1.06	-	-	-	-	-	-
IV(a). Neuroblastoma & ganglioneuroblastoma	72	24.2	2.85	10	3.3	1.06	-	-	-	-	-	-
V. Retinoblastoma	34	11.4	1.96	-	-	-	-	-	-	-	-	-
VI. Renal Tumors	21	7.1	1.54	-	-	-	-	-	-	8	2.7	0.96
VI(a). Nephroblastoma and other nonepithelial renal tumors	21	7.1	1.54	-	-	-	-	-	-	6	2.0	0.83
VII. Hepatic tumors	25	8.4	1.68	-	-	-	-	-	-	-	-	-
VII(a). Hepatoblastoma	25	8.4	1.68	-	-	-	-	-	-	-	-	-
VIII. Malignant bone tumors	-	-	-	13	4.4	1.21	36	12.6	2.10	41	13.9	2.17
VIII(a). Osteosarcomas	-	-	-	8	2.7	0.95	22	7.7	1.64	21	7.1	1.55
VIII(c). Ewing tumor and related sarcomas of bone	-	-	-	-	-	-	10	3.5	1.11	17	5.8	1.39
IX. Soft tissue and other extraosseous sarcomas	27	9.1	1.75	18	6.0	1.42	22	7.7	1.64	38	12.9	2.09
IX(a). Rhabdomyosarcomas	13	4.4	1.21	11	3.7	1.11	8	2.8	0.99	6	2.0	0.83
X. Germ cell & trophoblastic tumors & neoplasms of gonads	27	9.1	1.75	8	2.7	0.95	26	9.1	1.79	91	30.8	3.23
X(a). Intracranial & intraspinal germ cell tumors	-	-	-	8	2.7	0.95	20	7.0	1.57	30	10.1	1.85
X(c). Malignant gonadal germ cell tumors	17	5.7	1.39	-	-	-	-	-	-	47	15.9	2.32
XI. Other malignant epithelial neoplasms and melanomas	-	-	-	8	2.7	0.95	32	11.2	1.98	70	23.7	2.83
XI(b). Thyroid carcinomas	-	-	-	-	-	-	13	4.6	1.26	26	8.8	1.72
XI(d). Malignant melanomas	-	-	-	-	-	-	-	-	-	-	-	-
XII. Other & unspecified malignant neoplasms	-	-	-	-	-	-	-	-	-	-	-	-

[1] Rates are **per million** population.
[2] See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.
[3] Abbreviations: S.E. - Standard Error; CNS - Central Nervous System.
- Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for All Sites combined.

Total Cancer Cases and Average Annual Age-specific Incidence Rates[1] for Pediatric Cancers
NAACCR United States Combined[2], 2010-2014, Asian/Pacific Islander, Females

	Ages 0-4			Ages 5-9			Ages 10-14			Ages 15-19		
Cancer Types	*Count*	*Rate*	*S.E.*[3]	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
All Sites	**554**	**194.7**	**8.27**	**268**	**91.3**	**5.58**	**302**	**107.5**	**6.19**	**475**	**166.7**	**7.65**
I. Leukemias	209	73.4	5.08	102	34.7	3.44	80	28.5	3.18	76	26.7	3.06
I(a). Lymphoid leukemias	157	55.2	4.40	71	24.2	2.87	49	17.4	2.49	29	10.2	1.89
I(b). Acute myeloid leukemias	35	12.3	2.08	20	6.8	1.52	20	7.1	1.59	32	11.2	1.99
II. Lymphomas & reticuloendothelial neoplasms	42	14.8	2.28	30	10.2	1.87	38	13.5	2.19	92	32.3	3.37
II(a). Hodgkin lymphomas	-	-	-	-	-	-	12	4.3	1.23	45	15.8	2.35
II(b, c, e). Non-Hodgkin lymphomas	19	6.7	1.53	23	7.8	1.63	24	8.5	1.74	44	15.4	2.33
III. CNS[3] & misc. intracranial & intraspinal neoplasms	76	26.7	3.06	57	19.4	2.57	50	17.8	2.52	42	14.7	2.27
III(a). Ependymomas and choroids plexus tumor	6	2.1	0.86	6	2.0	0.83	-	-	-	-	-	-
III(b). Astrocytomas	44	15.5	2.33	19	6.5	1.48	27	9.6	1.85	24	8.4	1.72
III(c). Intracranial and intraspinal embryonal tumors	18	6.3	1.49	18	6.1	1.45	11	3.9	1.18	-	-	-
III(d). Other gliomas	7	2.5	0.93	11	3.7	1.13	8	2.8	1.01	13	4.6	1.27
IV. Neuroblastoma and other peripheral nervous cell tumors	73	25.7	3.00	10	3.4	1.08	-	-	-	-	-	-
IV(a). Neuroblastoma & ganglioneuroblastoma	73	25.7	3.00	10	3.4	1.08	-	-	-	-	-	-
V. Retinoblastoma	46	16.2	2.38	-	-	-	-	-	-	-	-	-
VI. Renal Tumors	27	9.5	1.83	10	3.4	1.08	-	-	-	-	-	-
VI(a). Nephroblastoma and other nonepithelial renal tumors	25	8.8	1.76	9	3.1	1.02	-	-	-	-	-	-
VII. Hepatic tumors	17	6.0	1.45	-	-	-	-	-	-	-	-	-
VII(a). Hepatoblastoma	17	6.0	1.45	-	-	-	-	-	-	-	-	-
VIII. Malignant bone tumors	-	-	-	17	5.8	1.40	22	7.8	1.67	24	8.4	1.72
VIII(a). Osteosarcomas	-	-	-	10	3.4	1.08	15	5.3	1.38	13	4.6	1.27
VIII(c). Ewing tumor and related sarcomas of bone	-	-	-	6	2.0	0.83	7	2.5	0.94	7	2.5	0.93
IX. Soft tissue and other extraosseous sarcomas	28	9.8	1.86	13	4.4	1.23	25	8.9	1.78	27	9.5	1.82
IX(a). Rhabdomyosarcomas	18	6.3	1.49	7	2.4	0.90	7	2.5	0.94	7	2.5	0.93
X. Germ cell & trophoblastic tumors & neoplasms of gonads	24	8.4	1.72	14	4.8	1.27	35	12.5	2.11	41	14.4	2.25
X(a). Intracranial & intraspinal germ cell tumors	-	-	-	-	-	-	9	3.2	1.07	-	-	-
X(c). Malignant gonadal germ cell tumors	-	-	-	7	2.4	0.90	24	8.5	1.74	22	7.7	1.65
XI. Other malignant epithelial neoplasms and melanomas	6	2.1	0.86	9	3.1	1.02	47	16.7	2.44	161	56.5	4.45
XI(b). Thyroid carcinomas	-	-	-	-	-	-	34	12.1	2.08	129	45.3	3.99
XI(d). Malignant melanomas	-	-	-	-	-	-	-	-	-	-	-	-
XII. Other & unspecified malignant neoplasms	-	-	-	-	-	-	-	-	-	-	-	-

[1] Rates are **per million** population.
[2] See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.
[3] Abbreviations: S.E. - Standard Error; CNS - Central Nervous System.
- Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for All Sites combined.

Total Cancer Cases and Average Annual Age-specific Incidence Rates[1] for Pediatric Cancers
NAACCR United States Combined[2], 2010-2014, American Indian/Alaskan Native for CHSDA[3] Counties, Males

	Ages 0-4			Ages 5-9			Ages 10-14			Ages 15-19		
Cancer Types	*Count*	*Rate*	*S.E.[4]*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
All Sites	**101**	**210.4**	**20.93**	**53**	**115.7**	**15.89**	**47**	**106.0**	**15.46**	**75**	**168.9**	**19.50**
I. Leukemias	35	72.9	12.32	19	41.5	9.52	10	22.5	7.13	12	27.0	7.80
I(a). Lymphoid leukemias	25	52.1	10.41	13	28.4	7.87	-	-	-	7	15.8	5.96
I(b). Acute myeloid leukemias	8	16.7	5.89	-	-	-	-	-	-	-	-	-
II. Lymphomas & reticuloendothelial neoplasms	8	16.7	5.89	12	26.2	7.56	12	27.1	7.81	13	29.3	8.12
II(a). Hodgkin lymphomas	-	-	-	-	-	-	-	-	-	-	-	-
II(b, c, e). Non-Hodgkin lymphomas	-	-	-	8	17.5	6.17	8	18.0	6.38	8	18.0	6.37
III. CNS[4] & misc. intracranial & intraspinal neoplasms	16	33.3	8.33	16	34.9	8.73	12	27.1	7.81	10	22.5	7.12
III(a). Ependymomas and choroids plexus tumor	-	-	-	-	-	-	-	-	-	-	-	-
III(b). Astrocytomas	7	14.6	5.51	7	15.3	5.78	-	-	-	7	15.8	5.96
III(c). Intracranial and intraspinal embryonal tumors	-	-	-	-	-	-	-	-	-	-	-	-
III(d). Other gliomas	-	-	-	-	-	-	-	-	-	-	-	-
IV. Neuroblastoma and other peripheral nervous cell tumors	10	20.8	6.59	-	-	-	-	-	-	-	-	-
IV(a). Neuroblastoma & ganglioneuroblastoma	10	20.8	6.59	-	-	-	-	-	-	-	-	-
V. Retinoblastoma	-	-	-	-	-	-	-	-	-	-	-	-
VI. Renal Tumors	6	12.5	5.10	-	-	-	-	-	-	-	-	-
VI(a). Nephroblastoma and other nonepithelial renal tumors	6	12.5	5.10	-	-	-	-	-	-	-	-	-
VII. Hepatic tumors	10	20.8	6.59	-	-	-	-	-	-	-	-	-
VII(a). Hepatoblastoma	10	20.8	6.59	-	-	-	-	-	-	-	-	-
VIII. Malignant bone tumors	-	-	-	-	-	-	-	-	-	-	-	-
VIII(a). Osteosarcomas	-	-	-	-	-	-	-	-	-	-	-	-
VIII(c). Ewing tumor and related sarcomas of bone	-	-	-	-	-	-	-	-	-	-	-	-
IX. Soft tissue and other extraosseous sarcomas	-	-	-	-	-	-	-	-	-	8	18.0	6.37
IX(a). Rhabdomyosarcomas	-	-	-	-	-	-	-	-	-	-	-	-
X. Germ cell & trophoblastic tumors & neoplasms of gonads	-	-	-	-	-	-	-	-	-	13	29.3	8.12
X(a). Intracranial & intraspinal germ cell tumors	-	-	-	-	-	-	-	-	-	-	-	-
X(c). Malignant gonadal germ cell tumors	-	-	-	-	-	-	-	-	-	12	27.0	7.80
XI. Other malignant epithelial neoplasms and melanomas	-	-	-	-	-	-	-	-	-	12	27.0	7.80
XI(b). Thyroid carcinomas	-	-	-	-	-	-	-	-	-	-	-	-
XI(d). Malignant melanomas	-	-	-	-	-	-	-	-	-	-	-	-
XII. Other & unspecified malignant neoplasms	-	-	-	-	-	-	-	-	-	-	-	-

[1] Rates are **per million** population.
[2] See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.
[3.] CHSDA - Indian Health Services Contract Health Service Delivery Areas (See Appendix G)
[4] Abbreviations: S.E. - Standard Error; CNS - Central Nervous System.
- Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for All Sites combined. Statistics exclude cases diagnosed for KS for all years.

Total Cancer Cases and Average Annual Age-specific Incidence Rates[1] for Pediatric Cancers
NAACCR United States Combined[2], 2010-2014, American Indian/Alaskan Native for CHSDA[3] Counties, Females

	Ages 0-4			Ages 5-9			Ages 10-14			Ages 15-19		
Cancer Types	*Count*	*Rate*	*S.E.[4]*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
All Sites	**57**	**122.8**	**16.26**	**52**	**116.9**	**16.21**	**39**	**90.6**	**14.51**	**66**	**154.7**	**19.05**
I. Leukemias	19	40.9	9.39	17	38.2	9.27	10	23.2	7.35	17	39.9	9.67
I(a). Lymphoid leukemias	14	30.2	8.06	13	29.2	8.11	-	-	-	-	-	-
I(b). Acute myeloid leukemias	-	-	-	-	-	-	-	-	-	12	28.1	8.12
II. Lymphomas & reticuloendothelial neoplasms	-	-	-	-	-	-	-	-	-	11	25.8	7.78
II(a). Hodgkin lymphomas	-	-	-	-	-	-	-	-	-	-	-	-
II(b, c, e). Non-Hodgkin lymphomas	-	-	-	-	-	-	-	-	-	6	14.1	5.74
III. CNS[4] & misc. intracranial & intraspinal neoplasms	11	23.7	7.14	14	31.5	8.41	-	-	-	-	-	-
III(a). Ependymomas and choroids plexus tumor	-	-	-	-	-	-	-	-	-	-	-	-
III(b). Astrocytomas	-	-	-	10	22.5	7.11	-	-	-	-	-	-
III(c). Intracranial and intraspinal embryonal tumors	6	12.9	5.28	-	-	-	-	-	-	-	-	-
III(d). Other gliomas	-	-	-	-	-	-	-	-	-	-	-	-
IV. Neuroblastoma and other peripheral nervous cell tumors	6	12.9	5.28	-	-	-	-	-	-	-	-	-
IV(a). Neuroblastoma & ganglioneuroblastoma	6	12.9	5.28	-	-	-	-	-	-	-	-	-
V. Retinoblastoma	-	-	-	-	-	-	-	-	-	-	-	-
VI. Renal Tumors	-	-	-	-	-	-	-	-	-	-	-	-
VI(a). Nephroblastoma and other nonepithelial renal tumors	-	-	-	-	-	-	-	-	-	-	-	-
VII. Hepatic tumors	-	-	-	-	-	-	-	-	-	-	-	-
VII(a). Hepatoblastoma	-	-	-	-	-	-	-	-	-	-	-	-
VIII. Malignant bone tumors	-	-	-	-	-	-	6	13.9	5.69	-	-	-
VIII(a). Osteosarcomas	-	-	-	-	-	-	6	13.9	5.69	-	-	-
VIII(c). Ewing tumor and related sarcomas of bone	-	-	-	-	-	-	-	-	-	-	-	-
IX. Soft tissue and other extraosseous sarcomas	-	-	-	-	-	-	6	13.9	5.69	7	16.4	6.20
IX(a). Rhabdomyosarcomas	-	-	-	-	-	-	-	-	-	-	-	-
X. Germ cell & trophoblastic tumors & neoplasms of gonads	-	-	-	-	-	-	-	-	-	-	-	-
X(a). Intracranial & intraspinal germ cell tumors	-	-	-	-	-	-	-	-	-	-	-	-
X(c). Malignant gonadal germ cell tumors	-	-	-	-	-	-	-	-	-	-	-	-
XI. Other malignant epithelial neoplasms and melanomas	-	-	-	-	-	-	7	16.3	6.15	19	44.5	10.22
XI(b). Thyroid carcinomas	-	-	-	-	-	-	-	-	-	10	23.4	7.41
XI(d). Malignant melanomas	-	-	-	-	-	-	-	-	-	-	-	-
XII. Other & unspecified malignant neoplasms	-	-	-	-	-	-	-	-	-	-	-	-

[1] Rates are **per million** population.
[2] See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.
[3] CHSDA - Indian Health Services Contract Health Service Delivery Areas (See Appendix G)
[4] Abbreviations: S.E. - Standard Error; CNS - Central Nervous System.
- Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for All Sites combined. Statistics exclude cases diagnosed for KS for all years.

Total Cancer Cases and Average Annual Age-specific Incidence Rates[1] for Pediatric Cancers
NAACCR United States Combined[2], 2010-2014, Hispanic/Latino, All Races, All Races, Males

	Ages 0-4			Ages 5-9			Ages 10-14			Ages 15-19		
Cancer Types	*Count*	*Rate*	*S.E.*[3]	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
All Sites	**2,926**	**232.1**	**4.29**	**1,695**	**138.1**	**3.35**	**1,726**	**150.2**	**3.61**	**2,629**	**231.8**	**4.52**
I. Leukemias	1,185	94.0	2.73	708	57.7	2.17	572	49.8	2.08	622	54.8	2.20
I(a). Lymphoid leukemias	919	72.9	2.40	593	48.3	1.98	397	34.5	1.73	399	35.2	1.76
I(b). Acute myeloid leukemias	159	12.6	1.00	64	5.2	0.65	105	9.1	0.89	127	11.2	0.99
II. Lymphomas & reticuloendothelial neoplasms	294	23.3	1.36	324	26.4	1.47	347	30.2	1.62	466	41.1	1.90
II(a). Hodgkin lymphomas	19	1.5	0.35	94	7.7	0.79	147	12.8	1.05	250	22.0	1.39
II(b, c, e). Non-Hodgkin lymphomas	95	7.5	0.77	158	12.9	1.02	155	13.5	1.08	198	17.5	1.24
III. CNS[3] & misc. intracranial & intraspinal neoplasms	423	33.6	1.63	337	27.5	1.50	259	22.5	1.40	205	18.1	1.26
III(a). Ependymomas and choroids plexus tumor	61	4.8	0.62	27	2.2	0.42	23	2.0	0.42	14	1.2	0.33
III(b). Astrocytomas	163	12.9	1.01	158	12.9	1.02	136	11.8	1.01	117	10.3	0.95
III(c). Intracranial and intraspinal embryonal tumors	138	10.9	0.93	81	6.6	0.73	50	4.3	0.62	24	2.1	0.43
III(d). Other gliomas	49	3.9	0.56	57	4.6	0.62	38	3.3	0.54	37	3.3	0.54
IV. Neuroblastoma and other peripheral nervous cell tumors	263	20.9	1.29	25	2.0	0.41	11	1.0	0.29	9	0.8	0.26
IV(a). Neuroblastoma & ganglioneuroblastoma	260	20.6	1.28	21	1.7	0.37	9	0.8	0.26	-	-	-
V. Retinoblastoma	171	13.6	1.04	8	0.7	0.23	-	-	-	-	-	-
VI. Renal Tumors	169	13.4	1.03	50	4.1	0.58	17	1.5	0.36	12	1.1	0.31
VI(a). Nephroblastoma and other nonepithelial renal tumors	166	13.2	1.02	48	3.9	0.56	7	0.6	0.23	-	-	-
VII. Hepatic tumors	136	10.8	0.93	13	1.1	0.29	14	1.2	0.33	12	1.1	0.31
VII(a). Hepatoblastoma	128	10.2	0.90	6	0.5	0.20	-	-	-	-	-	-
VIII. Malignant bone tumors	13	1.0	0.29	62	5.1	0.64	174	15.1	1.15	189	16.7	1.21
VIII(a). Osteosarcomas	-	-	-	36	2.9	0.49	106	9.2	0.90	120	10.6	0.97
VIII(c). Ewing tumor and related sarcomas of bone	9	0.7	0.24	23	1.9	0.39	54	4.7	0.64	42	3.7	0.57
IX. Soft tissue and other extraosseous sarcomas	152	12.1	0.98	105	8.6	0.83	119	10.4	0.95	168	14.8	1.14
IX(a). Rhabdomyosarcomas	84	6.7	0.73	56	4.6	0.61	39	3.4	0.54	38	3.4	0.54
X. Germ cell & trophoblastic tumors & neoplasms of gonads	80	6.3	0.71	25	2.0	0.41	92	8.0	0.83	717	63.2	2.36
X(a). Intracranial & intraspinal germ cell tumors	9	0.7	0.24	17	1.4	0.34	60	5.2	0.67	47	4.1	0.60
X(c). Malignant gonadal germ cell tumors	51	4.0	0.57	-	-	-	24	2.1	0.43	608	53.6	2.17
XI. Other malignant epithelial neoplasms and melanomas	24	1.9	0.39	37	3.0	0.50	112	9.7	0.92	225	19.8	1.32
XI(b). Thyroid carcinomas	-	-	-	8	0.7	0.23	44	3.8	0.58	89	7.8	0.83
XI(d). Malignant melanomas	9	0.7	0.24	15	1.2	0.32	11	1.0	0.29	16	1.4	0.35
XII. Other & unspecified malignant neoplasms	16	1.3	0.32	-	-	-	-	-	-	-	-	-

[1] Rates are **per million** population.
[2] See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.
[3] Abbreviations: S.E. - Standard Error; CNS - Central Nervous System.
- Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for All Sites combined.

N.B. In areas with small Latino populations, methods to indirectly identify Latinos (like NHIAv2) can overestimate the ethnicity-specific counts of cancer cases. Also, even small errors in Latino population estimates can affect the magnitude of the cancer rates. Data Item 190 used in place of NHIA for MA and RI, see technical notes.

Total Cancer Cases and Average Annual Age-specific Incidence Rates[1] for Pediatric Cancers
United States NAACCR Combined[2], 2010-2014, Hispanic/Latina, All Races, All Races, Females

	Ages 0-4			Ages 5-9			Ages 10-14			Ages 15-19		
Cancer Types	*Count*	*Rate*	*S.E.[3]*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
All Sites	**2,601**	**214.6**	**4.21**	**1,394**	**118.2**	**3.17**	**1,559**	**140.9**	**3.57**	**2,079**	**194.3**	**4.26**
I. Leukemias	1,053	86.9	2.68	550	46.6	1.99	433	39.1	1.88	353	33.0	1.76
I(a). Lymphoid leukemias	836	69.0	2.39	463	39.3	1.82	294	26.6	1.55	188	17.6	1.28
I(b). Acute myeloid leukemias	138	11.4	0.97	53	4.5	0.62	86	7.8	0.84	104	9.7	0.95
II. Lymphomas & reticuloendothelial neoplasms	186	15.3	1.13	160	13.6	1.07	243	22.0	1.41	377	35.2	1.81
II(a). Hodgkin lymphomas	10	0.8	0.26	30	2.5	0.46	105	9.5	0.93	253	23.6	1.49
II(b, c, e). Non-Hodgkin lymphomas	54	4.5	0.61	71	6.0	0.71	106	9.6	0.93	106	9.9	0.96
III. CNS[3] & misc. intracranial & intraspinal neoplasms	455	37.5	1.76	333	28.2	1.55	207	18.7	1.30	168	15.7	1.21
III(a). Ependymomas and choroids plexus tumor	63	5.2	0.65	33	2.8	0.49	13	1.2	0.33	15	1.4	0.36
III(b). Astrocytomas	197	16.3	1.16	172	14.6	1.11	122	11.0	1.00	99	9.3	0.93
III(c). Intracranial and intraspinal embryonal tumors	134	11.1	0.95	39	3.3	0.53	26	2.3	0.46	20	1.9	0.42
III(d). Other gliomas	48	4.0	0.57	76	6.4	0.74	37	3.3	0.55	26	2.4	0.48
IV. Neuroblastoma and other peripheral nervous cell tumors	233	19.2	1.26	29	2.5	0.46	7	0.6	0.24	-	-	-
IV(a). Neuroblastoma & ganglioneuroblastoma	232	19.1	1.26	29	2.5	0.46	-	-	-	-	-	-
V. Retinoblastoma	162	13.4	1.05	-	-	-	-	-	-	-	-	-
VI. Renal Tumors	206	17.0	1.18	56	4.7	0.63	14	1.3	0.34	16	1.5	0.37
VI(a). Nephroblastoma and other nonepithelial renal tumors	205	16.9	1.18	53	4.5	0.62	8	0.7	0.26	-	-	-
VII. Hepatic tumors	75	6.2	0.71	7	0.6	0.22	14	1.3	0.34	9	0.8	0.28
VII(a). Hepatoblastoma	74	6.1	0.71	-	-	-	-	-	-	-	-	-
VIII. Malignant bone tumors	22	1.8	0.39	69	5.8	0.70	158	14.3	1.14	92	8.6	0.90
VIII(a). Osteosarcomas	-	-	-	43	3.6	0.56	107	9.7	0.93	50	4.7	0.66
VIII(c). Ewing tumor and related sarcomas of bone	14	1.2	0.31	19	1.6	0.37	37	3.3	0.55	27	2.5	0.49
IX. Soft tissue and other extraosseous sarcomas	121	10.0	0.91	86	7.3	0.79	127	11.5	1.02	148	13.8	1.14
IX(a). Rhabdomyosarcomas	68	5.6	0.68	37	3.1	0.52	39	3.5	0.56	35	3.3	0.55
X. Germ cell & trophoblastic tumors & neoplasms of gonads	60	4.9	0.64	51	4.3	0.61	126	11.4	1.01	213	19.9	1.36
X(a). Intracranial & intraspinal germ cell tumors	7	0.6	0.22	13	1.1	0.31	14	1.3	0.34	7	0.7	0.25
X(c). Malignant gonadal germ cell tumors	10	0.8	0.26	31	2.6	0.47	97	8.8	0.89	137	12.8	1.09
XI. Other malignant epithelial neoplasms and melanomas	20	1.6	0.37	48	4.1	0.59	221	20.0	1.34	684	63.9	2.44
XI(b). Thyroid carcinomas	-	-	-	21	1.8	0.39	129	11.7	1.03	495	46.3	2.08
XI(d). Malignant melanomas	13	1.1	0.30	11	0.9	0.28	26	2.3	0.46	26	2.4	0.48
XII. Other & unspecified malignant neoplasms	8	0.7	0.23	-	-	-	9	0.8	0.27	17	1.6	0.39

[1] Rates are **per million** population.
[2] See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.
[3] Abbreviations: S.E. - Standard Error; CNS - Central Nervous System.
- Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for All Sites combined.

N.B. In areas with small Latino populations, methods to indirectly identify Latinos (like NHIAv2) can overestimate the ethnicity-specific counts of cancer cases. Also, even small errors in Latino population estimates can affect the magnitude of the cancer rates. Data Item 190 used in place of NHIA for MA and RI, see technical notes.

Total Cancer Cases and Average Annual Age-specific Incidence Rates[1] for Pediatric Cancers
NAACCR United States Combined[2], 2010-2014, Non-Hispanic White, All Races, Males

	Ages 0-4			Ages 5-9			Ages 10-14			Ages 15-19		
Cancer Types	*Count*	*Rate*	*S.E.*[3]	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
All Sites	**6,963**	**267.8**	**3.21**	**4,052**	**148.1**	**2.33**	**4,283**	**149.4**	**2.28**	**7,483**	**247.1**	**2.86**
I. Leukemias	2,292	88.2	1.84	1,286	47.0	1.31	911	31.8	1.05	1,119	36.9	1.10
I(a). Lymphoid leukemias	1,811	69.7	1.64	1,079	39.4	1.20	624	21.8	0.87	597	19.7	0.81
I(b). Acute myeloid leukemias	284	10.9	0.65	118	4.3	0.40	174	6.1	0.46	299	9.9	0.57
II. Lymphomas & reticuloendothelial neoplasms	543	20.9	0.90	685	25.0	0.96	943	32.9	1.07	1,951	64.4	1.46
II(a). Hodgkin lymphomas	14	0.5	0.14	122	4.5	0.40	377	13.1	0.68	1,124	37.1	1.11
II(b, c, e). Non-Hodgkin lymphomas	249	9.6	0.61	422	15.4	0.75	458	16.0	0.75	751	24.8	0.90
III. CNS[3] & misc. intracranial & intraspinal neoplasms	1,359	52.3	1.42	1,165	42.6	1.25	993	34.6	1.10	858	28.3	0.97
III(a). Ependymomas and choroids plexus tumor	184	7.1	0.52	66	2.4	0.30	63	2.2	0.28	52	1.7	0.24
III(b). Astrocytomas	634	24.4	0.97	605	22.1	0.90	550	19.2	0.82	500	16.5	0.74
III(c). Intracranial and intraspinal embryonal tumors	337	13.0	0.71	270	9.9	0.60	167	5.8	0.45	107	3.5	0.34
III(d). Other gliomas	156	6.0	0.48	204	7.5	0.52	188	6.6	0.48	162	5.3	0.42
IV. Neuroblastoma and other peripheral nervous cell tumors	952	36.6	1.19	141	5.2	0.43	49	1.7	0.24	30	1.0	0.18
IV(a). Neuroblastoma & ganglioneuroblastoma	948	36.5	1.18	138	5.0	0.43	34	1.2	0.20	15	0.5	0.13
V. Retinoblastoma	296	11.4	0.66	11	0.4	0.12	-	-	-	-	-	-
VI. Renal Tumors	536	20.6	0.89	142	5.2	0.44	32	1.1	0.20	53	1.7	0.24
VI(a). Nephroblastoma and other nonepithelial renal tumors	533	20.5	0.89	129	4.7	0.42	18	0.6	0.15	9	0.3	0.10
VII. Hepatic tumors	241	9.3	0.60	16	0.6	0.15	31	1.1	0.19	26	0.9	0.17
VII(a). Hepatoblastoma	231	8.9	0.58	10	0.4	0.12	-	-	-	-	-	-
VIII. Malignant bone tumors	48	1.8	0.27	189	6.9	0.50	432	15.1	0.72	537	17.7	0.77
VIII(a). Osteosarcomas	7	0.3	0.10	72	2.6	0.31	235	8.2	0.53	283	9.3	0.56
VIII(c). Ewing tumor and related sarcomas of bone	37	1.4	0.23	101	3.7	0.37	162	5.6	0.44	200	6.6	0.47
IX. Soft tissue and other extraosseous sarcomas	409	15.7	0.78	248	9.1	0.58	346	12.1	0.65	478	15.8	0.72
IX(a). Rhabdomyosarcomas	246	9.5	0.60	136	5.0	0.43	106	3.7	0.36	106	3.5	0.34
X. Germ cell & trophoblastic tumors & neoplasms of gonads	170	6.5	0.50	41	1.5	0.23	157	5.5	0.44	1,279	42.2	1.18
X(a). Intracranial & intraspinal germ cell tumors	17	0.7	0.16	38	1.4	0.23	109	3.8	0.36	115	3.8	0.35
X(c). Malignant gonadal germ cell tumors	81	3.1	0.35	-	-	-	35	1.2	0.21	1,104	36.4	1.10
XI. Other malignant epithelial neoplasms and melanomas	75	2.9	0.33	122	4.5	0.40	374	13.0	0.67	1,137	37.5	1.11
XI(b). Thyroid carcinomas	-	-	-	33	1.2	0.21	125	4.4	0.39	328	10.8	0.60
XI(d). Malignant melanomas	51	2.0	0.27	53	1.9	0.27	107	3.7	0.36	395	13.0	0.66
XII. Other & unspecified malignant neoplasms	42	1.6	0.25	6	0.2	0.09	13	0.5	0.13	15	0.5	0.13

[1] Rates are **per million** population.
[2] See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.
[3] Abbreviations: S.E. - Standard Error; CNS - Central Nervous System.
- Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for All Sites combined.

N.B. In areas with small Latino populations, methods to indirectly identify Latinos (like NHIAv2) can overestimate the ethnicity-specific counts of cancer cases. Also, even small errors in Latino population estimates can affect the magnitude of the cancer rates. Data Item 190 used in place of NHIA for MA and RI, see technical notes.

Total Cancer Cases and Average Annual Age-specific Incidence Rates[1] for Pediatric Cancers
United States NAACCR Combined[2], 2010-2014, Non-Hispanic White, All Races, Females

	Ages 0-4			Ages 5-9			Ages 10-14			Ages 15-19		
Cancer Types	*Count*	*Rate*	*S.E.[3]*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
All Sites	**6,043**	**244.2**	**3.14**	**3,206**	**123.2**	**2.18**	**3,953**	**145.2**	**2.31**	**6,850**	**238.8**	**2.89**
I. Leukemias	2,077	83.9	1.84	1,009	38.8	1.22	724	26.6	0.99	757	26.4	0.96
I(a). Lymphoid leukemias	1,619	65.4	1.63	821	31.5	1.10	448	16.5	0.78	270	9.4	0.57
I(b). Acute myeloid leukemias	287	11.6	0.68	113	4.3	0.41	165	6.1	0.47	289	10.1	0.59
II. Lymphomas & reticuloendothelial neoplasms	333	13.5	0.74	318	12.2	0.69	625	23.0	0.92	1,498	52.2	1.35
II(a). Hodgkin lymphomas	-	-	-	44	1.7	0.25	295	10.8	0.63	1,093	38.1	1.15
II(b, c, e). Non-Hodgkin lymphomas	117	4.7	0.44	170	6.5	0.50	246	9.0	0.58	363	12.7	0.66
III. CNS[3] & misc. intracranial & intraspinal neoplasms	1,150	46.5	1.37	933	35.8	1.17	859	31.6	1.08	703	24.5	0.92
III(a). Ependymomas and choroids plexus tumor	127	5.1	0.46	53	2.0	0.28	45	1.7	0.25	41	1.4	0.22
III(b). Astrocytomas	620	25.1	1.01	525	20.2	0.88	516	19.0	0.83	422	14.7	0.72
III(c). Intracranial and intraspinal embryonal tumors	229	9.3	0.61	144	5.5	0.46	96	3.5	0.36	55	1.9	0.26
III(d). Other gliomas	146	5.9	0.49	189	7.3	0.53	173	6.4	0.48	161	5.6	0.44
IV. Neuroblastoma and other peripheral nervous cell tumors	845	34.1	1.17	106	4.1	0.40	54	2.0	0.27	20	0.7	0.16
IV(a). Neuroblastoma & ganglioneuroblastoma	842	34.0	1.17	101	3.9	0.39	44	1.6	0.24	12	0.4	0.12
V. Retinoblastoma	306	12.4	0.71	14	0.5	0.14	-	-	-	-	-	-
VI. Renal Tumors	548	22.1	0.95	179	6.9	0.51	49	1.8	0.26	68	2.4	0.29
VI(a). Nephroblastoma and other nonepithelial renal tumors	543	21.9	0.94	174	6.7	0.51	30	1.1	0.20	15	0.5	0.14
VII. Hepatic tumors	137	5.5	0.47	21	0.8	0.18	19	0.7	0.16	42	1.5	0.23
VII(a). Hepatoblastoma	130	5.3	0.46	11	0.4	0.13	-	-	-	-	-	-
VIII. Malignant bone tumors	54	2.2	0.30	153	5.9	0.48	374	13.7	0.71	348	12.1	0.65
VIII(a). Osteosarcomas	7	0.3	0.11	70	2.7	0.32	204	7.5	0.52	181	6.3	0.47
VIII(c). Ewing tumor and related sarcomas of bone	35	1.4	0.24	67	2.6	0.31	142	5.2	0.44	109	3.8	0.36
IX. Soft tissue and other extraosseous sarcomas	305	12.3	0.71	206	7.9	0.55	319	11.7	0.66	367	12.8	0.67
IX(a). Rhabdomyosarcomas	178	7.2	0.54	108	4.1	0.40	92	3.4	0.35	81	2.8	0.31
X. Germ cell & trophoblastic tumors & neoplasms of gonads	171	6.9	0.53	84	3.2	0.35	210	7.7	0.53	336	11.7	0.64
X(a). Intracranial & intraspinal germ cell tumors	17	0.7	0.17	33	1.3	0.22	39	1.4	0.23	13	0.5	0.13
X(c). Malignant gonadal germ cell tumors	14	0.6	0.15	41	1.6	0.25	130	4.8	0.42	189	6.6	0.48
XI. Other malignant epithelial neoplasms and melanomas	92	3.7	0.39	176	6.8	0.51	704	25.9	0.97	2,667	93.0	1.80
XI(b). Thyroid carcinomas	10	0.4	0.13	59	2.3	0.30	414	15.2	0.75	1,586	55.3	1.39
XI(d). Malignant melanomas	62	2.5	0.32	79	3.0	0.34	111	4.1	0.39	586	20.4	0.84
XII. Other & unspecified malignant neoplasms	25	1.0	0.20	7	0.3	0.10	16	0.6	0.15	43	1.5	0.23

[1] Rates are **per million** population.
[2] See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.
[3] Abbreviations: S.E. - Standard Error; CNS - Central Nervous System.
- Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for All Sites combined.

N.B. In areas with small Latino populations, methods to indirectly identify Latinos (like NHIAv2) can overestimate the ethnicity-specific counts of cancer cases. Also, even small errors in Latino population estimates can affect the magnitude of the cancer rates. Data Item 190 used in place of NHIA for MA and RI, see technical notes.

Total Cancer Cases and Average Annual Age-specific Incidence Rates[1] for Pediatric Cancers
NAACCR United States Combined[2], 2010-2014, Non-Hispanic Black, All Races, Males

	Ages 0-4			Ages 5-9			Ages 10-14			Ages 15-19		
Cancer Types	*Count*	*Rate*	*S.E.*[3]	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
All Sites	**1,369**	**179.5**	**4.85**	**853**	**112.2**	**3.84**	**995**	**127.4**	**4.04**	**1,305**	**155.7**	**4.31**
I. Leukemias	381	50.0	2.56	235	30.9	2.02	233	29.8	1.95	239	28.5	1.84
I(a). Lymphoid leukemias	258	33.8	2.11	177	23.3	1.75	132	16.9	1.47	95	11.3	1.16
I(b). Acute myeloid leukemias	76	10.0	1.14	26	3.4	0.67	52	6.7	0.92	69	8.2	0.99
II. Lymphomas & reticuloendothelial neoplasms	86	11.3	1.22	151	19.9	1.62	252	32.3	2.03	399	47.6	2.38
II(a). Hodgkin lymphomas	9	1.2	0.39	55	7.2	0.98	98	12.5	1.27	194	23.2	1.66
II(b, c, e). Non-Hodgkin lymphomas	45	5.9	0.88	83	10.9	1.20	143	18.3	1.53	202	24.1	1.70
III. CNS[3] & misc. intracranial & intraspinal neoplasms	260	34.1	2.11	227	29.8	1.98	188	24.1	1.76	161	19.2	1.51
III(a). Ependymomas and choroids plexus tumor	39	5.1	0.82	8	1.1	0.37	12	1.5	0.44	13	1.6	0.43
III(b). Astrocytomas	113	14.8	1.39	108	14.2	1.37	95	12.2	1.25	87	10.4	1.11
III(c). Intracranial and intraspinal embryonal tumors	67	8.8	1.07	47	6.2	0.90	29	3.7	0.69	19	2.3	0.52
III(d). Other gliomas	32	4.2	0.74	53	7.0	0.96	39	5.0	0.80	33	3.9	0.69
IV. Neuroblastoma and other peripheral nervous cell tumors	170	22.3	1.71	37	4.9	0.80	17	2.2	0.53	9	1.1	0.36
IV(a). Neuroblastoma & ganglioneuroblastoma	168	22.0	1.70	37	4.9	0.80	12	1.5	0.44	-	-	-
V. Retinoblastoma	83	10.9	1.19	6	0.8	0.32	-	-	-	-	-	-
VI. Renal Tumors	164	21.5	1.68	48	6.3	0.91	14	1.8	0.48	21	2.5	0.55
VI(a). Nephroblastoma and other nonepithelial renal tumors	160	21.0	1.66	44	5.8	0.87	-	-	-	-	-	-
VII. Hepatic tumors	57	7.5	0.99	-	-	-	6	0.8	0.31	9	1.1	0.36
VII(a). Hepatoblastoma	55	7.2	0.97	-	-	-	-	-	-	-	-	-
VIII. Malignant bone tumors	6	0.8	0.32	29	3.8	0.71	103	13.2	1.30	102	12.2	1.21
VIII(a). Osteosarcomas	-	-	-	25	3.3	0.66	85	10.9	1.18	78	9.3	1.05
VIII(c). Ewing tumor and related sarcomas of bone	-	-	-	-	-	-	-	-	-	7	0.8	0.32
IX. Soft tissue and other extraosseous sarcomas	116	15.2	1.41	85	11.2	1.21	99	12.7	1.27	149	17.8	1.46
IX(a). Rhabdomyosarcomas	71	9.3	1.11	48	6.3	0.91	33	4.2	0.74	42	5.0	0.77
X. Germ cell & trophoblastic tumors & neoplasms of gonads	21	2.8	0.60	-	-	-	15	1.9	0.50	63	7.5	0.95
X(a). Intracranial & intraspinal germ cell tumors	-	-	-	-	-	-	13	1.7	0.46	21	2.5	0.55
X(c). Malignant gonadal germ cell tumors	7	0.9	0.35	-	-	-	-	-	-	34	4.1	0.70
XI. Other malignant epithelial neoplasms and melanomas	18	2.4	0.56	23	3.0	0.63	67	8.6	1.05	145	17.3	1.44
XI(b). Thyroid carcinomas	-	-	-	-	-	-	19	2.4	0.56	28	3.3	0.63
XI(d). Malignant melanomas	-	-	-	-	-	-	-	-	-	-	-	-
XII. Other & unspecified malignant neoplasms	7	0.9	0.35	-	-	-	-	-	-	8	1.0	0.34

[1] Rates are **per million** population.
[2] See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.
[3] Abbreviations: S.E. - Standard Error; CNS - Central Nervous System.
- Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for All Sites combined.

N.B. In areas with small Latino populations, methods to indirectly identify Latinos (like NHIAv2) can overestimate the ethnicity-specific counts of cancer cases. Also, even small errors in Latino population estimates can affect the magnitude of the cancer rates. Data Item 190 used in place of NHIA for MA and RI, see technical notes.

Total Cancer Cases and Average Annual Age-specific Incidence Rates[1] for Pediatric Cancers
United States NAACCR Combined[2], 2010-2014, Non-Hispanic Black, All Races, Females

	Ages 0-4			Ages 5-9			Ages 10-14			Ages 15-19		
Cancer Types	*Count*	*Rate*	*S.E.[3]*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
All Sites	**1,232**	**167.0**	**4.76**	**764**	**103.7**	**3.75**	**861**	**113.9**	**3.88**	**1,273**	**157.1**	**4.40**
I. Leukemias	316	42.8	2.41	171	23.2	1.77	151	20.0	1.63	172	21.2	1.62
I(a). Lymphoid leukemias	206	27.9	1.95	125	17.0	1.52	76	10.1	1.15	48	5.9	0.85
I(b). Acute myeloid leukemias	71	9.6	1.14	27	3.7	0.71	43	5.7	0.87	71	8.8	1.04
II. Lymphomas & reticuloendothelial neoplasms	50	6.8	0.96	71	9.6	1.14	148	19.6	1.61	308	38.0	2.17
II(a). Hodgkin lymphomas	-	-	-	11	1.5	0.45	80	10.6	1.18	202	24.9	1.75
II(b, c, e). Non-Hodgkin lymphomas	25	3.4	0.68	51	6.9	0.97	64	8.5	1.06	106	13.1	1.27
III. CNS[3] & misc. intracranial & intraspinal neoplasms	239	32.4	2.10	223	30.3	2.03	168	22.2	1.71	128	15.8	1.40
III(a). Ependymomas and choroids plexus tumor	31	4.2	0.75	16	2.2	0.54	8	1.1	0.37	10	1.2	0.39
III(b). Astrocytomas	105	14.2	1.39	106	14.4	1.40	96	12.7	1.30	70	8.6	1.03
III(c). Intracranial and intraspinal embryonal tumors	56	7.6	1.01	25	3.4	0.68	18	2.4	0.56	14	1.7	0.46
III(d). Other gliomas	35	4.7	0.80	63	8.5	1.08	33	4.4	0.76	26	3.2	0.63
IV. Neuroblastoma and other peripheral nervous cell tumors	173	23.5	1.78	46	6.2	0.92	8	1.1	0.37	-	-	-
IV(a). Neuroblastoma & ganglioneuroblastoma	172	23.3	1.78	43	5.8	0.89	7	0.9	0.35	-	-	-
V. Retinoblastoma	113	15.3	1.44	-	-	-	-	-	-	-	-	-
VI. Renal Tumors	177	24.0	1.80	82	11.1	1.23	17	2.2	0.55	20	2.5	0.55
VI(a). Nephroblastoma and other nonepithelial renal tumors	177	24.0	1.80	77	10.4	1.19	7	0.9	0.35	-	-	-
VII. Hepatic tumors	23	3.1	0.65	8	1.1	0.38	7	0.9	0.35	11	1.4	0.41
VII(a). Hepatoblastoma	23	3.1	0.65	-	-	-	-	-	-	-	-	-
VIII. Malignant bone tumors	-	-	-	42	5.7	0.88	90	11.9	1.25	76	9.4	1.08
VIII(a). Osteosarcomas	-	-	-	34	4.6	0.79	75	9.9	1.15	55	6.8	0.91
VIII(c). Ewing tumor and related sarcomas of bone	-	-	-	-	-	-	6	0.8	0.32	7	0.9	0.33
IX. Soft tissue and other extraosseous sarcomas	66	8.9	1.10	59	8.0	1.04	106	14.0	1.36	125	15.4	1.38
IX(a). Rhabdomyosarcomas	41	5.6	0.87	26	3.5	0.69	32	4.2	0.75	29	3.6	0.66
X. Germ cell & trophoblastic tumors & neoplasms of gonads	43	5.8	0.89	34	4.6	0.79	74	9.8	1.14	131	16.2	1.41
X(a). Intracranial & intraspinal germ cell tumors	-	-	-	-	-	-	11	1.5	0.44	-	-	-
X(c). Malignant gonadal germ cell tumors	-	-	-	28	3.8	0.72	53	7.0	0.96	75	9.3	1.07
XI. Other malignant epithelial neoplasms and melanomas	12	1.6	0.47	21	2.8	0.62	84	11.1	1.21	279	34.4	2.06
XI(b). Thyroid carcinomas	-	-	-	8	1.1	0.38	34	4.5	0.77	133	16.4	1.42
XI(d). Malignant melanomas	6	0.8	0.33	-	-	-	6	0.8	0.32	-	-	-
XII. Other & unspecified malignant neoplasms	15	2.0	0.53	-	-	-	7	0.9	0.35	18	2.2	0.52

[1] Rates are **per million** population.
[2] See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.
[3] Abbreviations: S.E. - Standard Error; CNS - Central Nervous System.
- Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for All Sites combined.

N.B. In areas with small Latino populations, methods to indirectly identify Latinos (like NHIAv2) can overestimate the ethnicity-specific counts of cancer cases. Also, even small errors in Latino population estimates can affect the magnitude of the cancer rates. Data Item 190 used in place of NHIA for MA and RI, see technical notes.

Total Cancer Cases and Average Annual Age-adjusted Incidence Rates[1] for Pediatric Cancers
NAACCR United States Combined[2], 2010-2014, All Races, Males

	Ages 0-14					Ages 0-19				
		2000 U.S.		1996 Cdn			2000 U.S.		1996 Cdn	
Cancer Types	*Count*	*Rate*	*S.E.*[3]	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Rate*	*S.E.*
All Sites	**26,854**	**177.4**	**1.08**	**177.9**	**1.09**	**39,071**	**190.5**	**0.96**	**190.8**	**0.97**
I. Leukemias	8,464	55.8	0.61	56.3	0.61	10,586	51.8	0.50	52.2	0.51
I(a). Lymphoid leukemias	6,466	42.6	0.53	43.1	0.54	7,628	37.4	0.43	37.8	0.43
I(b). Acute myeloid leukemias	1,176	7.8	0.23	7.8	0.23	1,713	8.3	0.20	8.4	0.20
II. Lymphomas & reticuloendothelial neoplasms	3,938	26.1	0.42	26.1	0.42	6,936	33.7	0.40	33.7	0.40
II(a). Hodgkin lymphomas	991	6.6	0.21	6.5	0.21	2,631	12.7	0.25	12.6	0.25
II(b, c, e). Non-Hodgkin lymphomas	1,982	13.2	0.30	13.1	0.30	3,235	15.8	0.28	15.8	0.28
III. CNS[3] & misc. intracranial & intraspinal neoplasms	5,587	37.0	0.49	37.1	0.50	6,888	33.8	0.41	33.9	0.41
III(a). Ependymomas and choroids plexus tumor	523	3.4	0.15	3.5	0.15	609	3.0	0.12	3.0	0.12
III(b). Astrocytomas	2,730	18.1	0.35	18.1	0.35	3,489	17.1	0.29	17.2	0.29
III(c). Intracranial and intraspinal embryonal tumors	1,276	8.4	0.24	8.5	0.24	1,431	7.0	0.19	7.1	0.19
III(d). Other gliomas	877	5.8	0.20	5.8	0.20	1,117	5.5	0.16	5.5	0.16
IV. Neuroblastoma and other peripheral nervous cell tumors	1,789	11.7	0.28	11.8	0.28	1,840	9.0	0.21	9.0	0.21
IV(a). Neuroblastoma & ganglioneuroblastoma	1,751	11.5	0.27	11.5	0.27	1,775	8.7	0.21	8.7	0.21
V. Retinoblastoma	636	4.2	0.17	4.2	0.17	636	3.1	0.12	3.1	0.12
VI. Renal Tumors	1,222	8.0	0.23	8.1	0.23	1,319	6.5	0.18	6.6	0.18
VI(a). Nephroblastoma and other nonepithelial renal tumors	1,155	7.6	0.22	7.7	0.23	1,174	5.8	0.17	5.9	0.17
VII. Hepatic tumors	572	3.8	0.16	3.8	0.16	625	3.1	0.12	3.1	0.12
VII(a). Hepatoblastoma	497	3.3	0.15	3.3	0.15	499	2.4	0.11	2.5	0.11
VIII. Malignant bone tumors	1,137	7.6	0.22	7.5	0.22	2,019	9.8	0.22	9.8	0.22
VIII(a). Osteosarcomas	615	4.1	0.17	4.1	0.16	1,126	5.5	0.16	5.4	0.16
VIII(c). Ewing tumor and related sarcomas of bone	422	2.8	0.14	2.8	0.14	689	3.4	0.13	3.4	0.13
IX. Soft tissue and other extraosseous sarcomas	1,794	11.9	0.28	11.9	0.28	2,652	12.9	0.25	12.9	0.25
IX(a). Rhabdomyosarcomas	872	5.8	0.19	5.8	0.20	1,069	5.2	0.16	5.3	0.16
X. Germ cell & trophoblastic tumors & neoplasms of gonads	693	4.6	0.17	4.5	0.17	2,885	13.8	0.26	13.7	0.26
X(a). Intracranial & intraspinal germ cell tumors	306	2.0	0.12	2.0	0.12	519	2.5	0.11	2.5	0.11
X(c). Malignant gonadal germ cell tumors	230	1.5	0.10	1.5	0.10	2,064	9.8	0.22	9.8	0.22
XI. Other malignant epithelial neoplasms and melanomas	923	6.1	0.20	6.1	0.20	2,559	12.3	0.24	12.3	0.24
XI(b). Thyroid carcinomas	259	1.7	0.11	1.7	0.11	749	3.6	0.13	3.6	0.13
XI(d). Malignant melanomas	276	1.8	0.11	1.8	0.11	714	3.4	0.13	3.4	0.13
XII. Other & unspecified malignant neoplasms	99	0.7	0.07	0.7	0.07	126	0.6	0.05	0.6	0.05

[1] Rates are **per million** population and are age-adjusted by five-year age groups to the 2000 U.S. standard population based on single years of age and to the 1996 Canadian population standard.
[2] See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.
[3] Abbreviations: S.E. - Standard Error; CNS - Central Nervous System.
- Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for All Sites combined.

Total Cancer Cases and Average Annual Age-adjusted Incidence Rates[1] for Pediatric Cancers
NAACCR United States Combined[2], 2010-2014, All Races, Females

	Ages 0-14					Ages 0-19				
		2000 U.S.		1996 Cdn			2000 U.S.		1996 Cdn	
Cancer Types	*Count*	*Rate*	*S.E.[3]*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Rate*	*S.E.*
All Sites	**23,285**	**160.6**	**1.05**	**161.0**	**1.06**	**34,223**	**174.6**	**0.94**	**174.8**	**0.95**
I. Leukemias	7,032	48.4	0.58	48.8	0.58	8,426	43.2	0.47	43.5	0.47
I(a). Lymphoid leukemias	5,266	36.2	0.50	36.7	0.51	5,807	29.8	0.39	30.1	0.40
I(b). Acute myeloid leukemias	1,087	7.5	0.23	7.5	0.23	1,598	8.2	0.20	8.1	0.20
II. Lymphomas & reticuloendothelial neoplasms	2,290	15.9	0.33	15.8	0.33	4,600	23.4	0.34	23.3	0.34
II(a). Hodgkin lymphomas	605	4.2	0.17	4.2	0.17	2,221	11.2	0.24	11.1	0.24
II(b, c, e). Non-Hodgkin lymphomas	986	6.8	0.22	6.8	0.22	1,617	8.3	0.21	8.2	0.20
III. CNS[3] & misc. intracranial & intraspinal neoplasms	4,873	33.7	0.48	33.8	0.48	5,940	30.5	0.40	30.6	0.40
III(a). Ependymomas and choroids plexus tumor	411	2.8	0.14	2.8	0.14	483	2.5	0.11	2.5	0.11
III(b). Astrocytomas	2,618	18.1	0.35	18.2	0.35	3,246	16.7	0.29	16.7	0.29
III(c). Intracranial and intraspinal embryonal tumors	831	5.7	0.20	5.8	0.20	923	4.7	0.16	4.8	0.16
III(d). Other gliomas	846	5.9	0.20	5.9	0.20	1,078	5.5	0.17	5.5	0.17
IV. Neuroblastoma and other peripheral nervous cell tumors	1,620	11.1	0.28	11.1	0.28	1,653	8.5	0.21	8.5	0.21
IV(a). Neuroblastoma & ganglioneuroblastoma	1,593	10.9	0.27	10.9	0.27	1,612	8.3	0.21	8.3	0.21
V. Retinoblastoma	668	4.6	0.18	4.6	0.18	669	3.4	0.13	3.4	0.13
VI. Renal Tumors	1,390	9.5	0.26	9.7	0.26	1,497	7.7	0.20	7.8	0.20
VI(a). Nephroblastoma and other nonepithelial renal tumors	1,332	9.1	0.25	9.3	0.25	1,354	6.9	0.19	7.0	0.19
VII. Hepatic tumors	342	2.3	0.13	2.4	0.13	410	2.1	0.10	2.1	0.10
VII(a). Hepatoblastoma	281	1.9	0.11	1.9	0.12	284	1.5	0.09	1.5	0.09
VIII. Malignant bone tumors	1,027	7.1	0.22	7.1	0.22	1,578	8.1	0.20	8.0	0.20
VIII(a). Osteosarcomas	581	4.0	0.17	4.0	0.17	890	4.6	0.15	4.5	0.15
VIII(c). Ewing tumor and related sarcomas of bone	349	2.4	0.13	2.4	0.13	499	2.6	0.11	2.5	0.11
IX. Soft tissue and other extraosseous sarcomas	1,511	10.4	0.27	10.4	0.27	2,199	11.2	0.24	11.2	0.24
IX(a). Rhabdomyosarcomas	682	4.7	0.18	4.7	0.18	838	4.3	0.15	4.3	0.15
X. Germ cell & trophoblastic tumors & neoplasms of gonads	949	6.6	0.21	6.5	0.21	1,690	8.6	0.21	8.5	0.21
X(a). Intracranial & intraspinal germ cell tumors	159	1.1	0.09	1.1	0.09	187	1.0	0.07	1.0	0.07
X(c). Malignant gonadal germ cell tumors	448	3.1	0.15	3.1	0.15	883	4.5	0.15	4.5	0.15
XI. Other malignant epithelial neoplasms and melanomas	1,485	10.3	0.27	10.2	0.26	5,383	27.1	0.37	27.0	0.37
XI(b). Thyroid carcinomas	735	5.1	0.19	5.1	0.19	3,126	15.7	0.28	15.7	0.28
XI(d). Malignant melanomas	332	2.3	0.13	2.3	0.13	985	5.0	0.16	5.0	0.16
XII. Other & unspecified malignant neoplasms	98	0.7	0.07	0.7	0.07	178	0.9	0.07	0.9	0.07

[1] Rates are **per million** population and are age-adjusted by five-year age groups to the 2000 U.S. standard population based on single years of age and to the 1996 Canadian population standard.
[2] See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.
[3] Abbreviations: S.E. - Standard Error; CNS - Central Nervous System.
- Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for All Sites combined.

Total Cancer Cases and Average Annual Age-adjusted Incidence Rates[1] for Pediatric Cancers
NAACCR United States Combined[2], 2010-2014, White, Males

	Ages 0-14					Ages 0-19				
		2000 U.S.		1996 Cdn			2000 U.S.		1996 Cdn	
Cancer Types	*Count*	*Rate*	*S.E.*[3]	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Rate*	*S.E.*
All Sites	**21,073**	**184.3**	**1.27**	**184.9**	**1.27**	**30,975**	**199.7**	**1.13**	**200.0**	**1.14**
I. Leukemias	6,720	58.7	0.72	59.3	0.72	8,424	54.6	0.59	55.0	0.60
I(a). Lymphoid leukemias	5,250	45.9	0.63	46.4	0.64	6,221	40.4	0.51	40.8	0.52
I(b). Acute myeloid leukemias	860	7.5	0.26	7.5	0.26	1,280	8.2	0.23	8.2	0.23
II. Lymphomas & reticuloendothelial neoplasms	3,047	26.7	0.48	26.6	0.48	5,421	34.8	0.47	34.7	0.47
II(a). Hodgkin lymphomas	753	6.6	0.24	6.5	0.24	2,108	13.4	0.29	13.3	0.29
II(b, c, e). Non-Hodgkin lymphomas	1,491	13.1	0.34	13.1	0.34	2,419	15.6	0.32	15.6	0.32
III. CNS[3] & misc. intracranial & intraspinal neoplasms	4,459	39.0	0.58	39.2	0.59	5,499	35.7	0.48	35.8	0.48
III(a). Ependymomas and choroids plexus tumor	419	3.7	0.18	3.7	0.18	483	3.1	0.14	3.2	0.14
III(b). Astrocytomas	2,208	19.3	0.41	19.4	0.41	2,811	18.2	0.34	18.3	0.34
III(c). Intracranial and intraspinal embryonal tumors	1,026	9.0	0.28	9.0	0.28	1,155	7.5	0.22	7.6	0.22
III(d). Other gliomas	678	5.9	0.23	5.9	0.23	872	5.7	0.19	5.7	0.19
IV. Neuroblastoma and other peripheral nervous cell tumors	1,413	12.3	0.33	12.3	0.33	1,450	9.5	0.25	9.5	0.25
IV(a). Neuroblastoma & ganglioneuroblastoma	1,383	12.1	0.32	12.1	0.32	1,402	9.2	0.24	9.2	0.24
V. Retinoblastoma	470	4.1	0.19	4.1	0.19	470	3.1	0.14	3.1	0.14
VI. Renal Tumors	920	8.0	0.26	8.1	0.27	985	6.4	0.20	6.5	0.21
VI(a). Nephroblastoma and other nonepithelial renal tumors	877	7.6	0.26	7.8	0.26	888	5.8	0.19	5.9	0.20
VII. Hepatic tumors	438	3.8	0.18	3.8	0.18	476	3.1	0.14	3.1	0.14
VII(a). Hepatoblastoma	375	3.3	0.17	3.3	0.17	377	2.5	0.13	2.5	0.13
VIII. Malignant bone tumors	906	7.9	0.26	7.9	0.26	1,615	10.4	0.26	10.3	0.26
VIII(a). Osteosarcomas	448	3.9	0.19	3.9	0.18	841	5.4	0.19	5.4	0.18
VIII(c). Ewing tumor and related sarcomas of bone	384	3.4	0.17	3.4	0.17	623	4.0	0.16	4.0	0.16
IX. Soft tissue and other extraosseous sarcomas	1,344	11.8	0.32	11.8	0.32	1,974	12.7	0.29	12.7	0.29
IX(a). Rhabdomyosarcomas	654	5.7	0.22	5.8	0.23	793	5.1	0.18	5.2	0.18
X. Germ cell & trophoblastic tumors & neoplasms of gonads	545	4.8	0.20	4.7	0.20	2,489	15.7	0.31	15.6	0.31
X(a). Intracranial & intraspinal germ cell tumors	242	2.1	0.14	2.1	0.13	401	2.6	0.13	2.6	0.13
X(c). Malignant gonadal germ cell tumors	185	1.6	0.12	1.6	0.12	1,851	11.6	0.27	11.6	0.27
XI. Other malignant epithelial neoplasms and melanomas	728	6.4	0.24	6.3	0.23	2,070	13.1	0.29	13.1	0.29
XI(b). Thyroid carcinomas	213	1.9	0.13	1.8	0.13	625	4.0	0.16	3.9	0.16
XI(d). Malignant melanomas	242	2.1	0.14	2.1	0.14	650	4.1	0.16	4.1	0.16
XII. Other & unspecified malignant neoplasms	83	0.7	0.08	0.7	0.08	102	0.7	0.07	0.7	0.07

[1] Rates are **per million** population and are age-adjusted by five-year age groups to the 2000 U.S. standard population based on single years of age and to the 1996 Canadian population standard.
[2] See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.
[3] Abbreviations: S.E. - Standard Error; CNS - Central Nervous System.
- Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for All Sites combined.

Total Cancer Cases and Average Annual Age-adjusted Incidence Rates[1] for Pediatric Cancers
NAACCR United States Combined[2], 2010-2014, White, Females

	Ages 0-14					Ages 0-19				
		2000 U.S.		1996 Cdn			2000 U.S.		1996 Cdn	
Cancer Types	*Count*	*Rate*	*S.E.*[3]	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Rate*	*S.E.*
All Sites	**18,262**	**167.4**	**1.24**	**167.8**	**1.24**	**27,006**	**182.9**	**1.11**	**183.2**	**1.12**
I. Leukemias	5,675	52.0	0.69	52.5	0.70	6,753	46.0	0.56	46.4	0.56
I(a). Lymphoid leukemias	4,356	39.9	0.60	40.4	0.61	4,805	32.8	0.47	33.2	0.48
I(b). Acute myeloid leukemias	810	7.4	0.26	7.4	0.26	1,190	8.1	0.23	8.1	0.23
II. Lymphomas & reticuloendothelial neoplasms	1,806	16.6	0.39	16.5	0.39	3,640	24.5	0.41	24.4	0.41
II(a). Hodgkin lymphomas	472	4.3	0.20	4.3	0.20	1,791	11.9	0.28	11.9	0.28
II(b, c, e). Non-Hodgkin lymphomas	741	6.8	0.25	6.8	0.25	1,198	8.1	0.23	8.1	0.23
III. CNS[3] & misc. intracranial & intraspinal neoplasms	3,852	35.3	0.57	35.5	0.57	4,704	32.1	0.47	32.2	0.47
III(a). Ependymomas and choroids plexus tumor	326	3.0	0.17	3.0	0.17	381	2.6	0.13	2.6	0.13
III(b). Astrocytomas	2,107	19.3	0.42	19.4	0.42	2,619	17.8	0.35	17.9	0.35
III(c). Intracranial and intraspinal embryonal tumors	649	5.9	0.23	6.0	0.23	722	4.9	0.18	5.0	0.18
III(d). Other gliomas	658	6.0	0.24	6.1	0.24	839	5.7	0.20	5.7	0.20
IV. Neuroblastoma and other peripheral nervous cell tumors	1,248	11.4	0.32	11.4	0.32	1,270	8.7	0.24	8.7	0.24
IV(a). Neuroblastoma & ganglioneuroblastoma	1,227	11.2	0.32	11.2	0.32	1,240	8.5	0.24	8.5	0.24
V. Retinoblastoma	471	4.3	0.20	4.3	0.20	472	3.2	0.15	3.2	0.15
VI. Renal Tumors	1,019	9.3	0.29	9.4	0.30	1,103	7.5	0.23	7.6	0.23
VI(a). Nephroblastoma and other nonepithelial renal tumors	982	9.0	0.29	9.1	0.29	1,001	6.8	0.22	6.9	0.22
VII. Hepatic tumors	265	2.4	0.15	2.4	0.15	314	2.1	0.12	2.1	0.12
VII(a). Hepatoblastoma	221	2.0	0.14	2.0	0.14	224	1.5	0.10	1.5	0.10
VIII. Malignant bone tumors	811	7.5	0.26	7.4	0.26	1,240	8.4	0.24	8.4	0.24
VIII(a). Osteosarcomas	420	3.9	0.19	3.8	0.19	644	4.4	0.17	4.3	0.17
VIII(c). Ewing tumor and related sarcomas of bone	311	2.9	0.16	2.8	0.16	445	3.0	0.14	3.0	0.14
IX. Soft tissue and other extraosseous sarcomas	1,135	10.4	0.31	10.4	0.31	1,642	11.1	0.27	11.1	0.27
IX(a). Rhabdomyosarcomas	512	4.7	0.21	4.7	0.21	627	4.3	0.17	4.3	0.17
X. Germ cell & trophoblastic tumors & neoplasms of gonads	680	6.3	0.24	6.2	0.24	1,212	8.2	0.24	8.1	0.23
X(a). Intracranial & intraspinal germ cell tumors	120	1.1	0.10	1.1	0.10	140	1.0	0.08	0.9	0.08
X(c). Malignant gonadal germ cell tumors	311	2.9	0.16	2.8	0.16	623	4.2	0.17	4.2	0.17
XI. Other malignant epithelial neoplasms and melanomas	1,235	11.4	0.32	11.2	0.32	4,532	30.2	0.45	30.1	0.45
XI(b). Thyroid carcinomas	623	5.7	0.23	5.7	0.23	2,664	17.7	0.34	17.7	0.34
XI(d). Malignant melanomas	294	2.7	0.16	2.7	0.16	905	6.0	0.20	6.0	0.20
XII. Other & unspecified malignant neoplasms	65	0.6	0.07	0.6	0.07	124	0.8	0.07	0.8	0.07

[1] Rates are **per million** population and are age-adjusted by five-year age groups to the 2000 U.S. standard population based on single years of age and to the 1996 Canadian population standard.
[2] See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.
[3] Abbreviations: S.E. - Standard Error; CNS - Central Nervous System.
- Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for All Sites combined.

Total Cancer Cases and Average Annual Age-adjusted Incidence Rates[1] for Pediatric Cancers
NAACCR United States Combined[2], 2010-2014, Black, Males

	Ages 0-14					Ages 0-19				
		2000 U.S.		1996 Cdn			2000 U.S.		1996 Cdn	
Cancer Types	*Count*	*Rate*	*S.E.*[3]	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Rate*	*S.E.*
All Sites	**3,401**	**133.7**	**2.29**	**134.0**	**2.30**	**4,783**	**138.5**	**2.00**	**138.8**	**2.01**
I. Leukemias	923	36.2	1.19	36.5	1.20	1,176	34.2	1.00	34.3	1.00
I(a). Lymphoid leukemias	615	24.1	0.97	24.4	0.98	720	21.0	0.78	21.2	0.79
I(b). Acute myeloid leukemias	173	6.8	0.52	6.8	0.52	245	7.1	0.45	7.1	0.45
II. Lymphomas & reticuloendothelial neoplasms	514	20.5	0.90	20.4	0.90	925	26.8	0.88	26.7	0.88
II(a). Hodgkin lymphomas	168	6.7	0.52	6.7	0.52	369	10.6	0.55	10.6	0.55
II(b, c, e). Non-Hodgkin lymphomas	284	11.3	0.67	11.3	0.67	490	14.2	0.64	14.2	0.64
III. CNS[3] & misc. intracranial & intraspinal neoplasms	707	27.9	1.05	28.0	1.05	881	25.7	0.87	25.8	0.87
III(a). Ependymomas and choroids plexus tumor	60	2.3	0.30	2.3	0.30	74	2.1	0.25	2.1	0.25
III(b). Astrocytomas	333	13.1	0.72	13.2	0.72	429	12.5	0.60	12.5	0.61
III(c). Intracranial and intraspinal embryonal tumors	151	5.9	0.48	6.0	0.48	171	5.0	0.38	5.0	0.38
III(d). Other gliomas	130	5.2	0.45	5.2	0.45	165	4.8	0.38	4.8	0.38
IV. Neuroblastoma and other peripheral nervous cell tumors	237	9.1	0.59	9.2	0.60	247	7.1	0.45	7.2	0.46
IV(a). Neuroblastoma & ganglioneuroblastoma	229	8.8	0.58	8.9	0.59	232	6.7	0.44	6.7	0.44
V. Retinoblastoma	93	3.6	0.37	3.6	0.37	93	2.7	0.28	2.7	0.28
VI. Renal Tumors	232	9.0	0.59	9.1	0.60	253	7.3	0.46	7.4	0.47
VI(a). Nephroblastoma and other nonepithelial renal tumors	214	8.3	0.56	8.4	0.57	215	6.2	0.42	6.3	0.43
VII. Hepatic tumors	68	2.6	0.32	2.6	0.32	77	2.2	0.25	2.2	0.25
VII(a). Hepatoblastoma	62	2.4	0.30	2.4	0.30	62	1.8	0.23	1.8	0.23
VIII. Malignant bone tumors	142	5.7	0.48	5.6	0.47	254	7.4	0.46	7.3	0.46
VIII(a). Osteosarcomas	117	4.7	0.43	4.6	0.43	202	5.9	0.41	5.8	0.41
VIII(c). Ewing tumor and related sarcomas of bone	10	0.4	0.13	0.4	0.13	19	0.6	0.13	0.5	0.13
IX. Soft tissue and other extraosseous sarcomas	314	12.4	0.70	12.4	0.70	471	13.6	0.63	13.6	0.63
IX(a). Rhabdomyosarcomas	160	6.3	0.50	6.3	0.50	203	5.9	0.41	5.9	0.42
X. Germ cell & trophoblastic tumors & neoplasms of gonads	49	1.9	0.27	1.9	0.27	124	3.5	0.32	3.5	0.31
X(a). Intracranial & intraspinal germ cell tumors	23	0.9	0.19	0.9	0.19	46	1.3	0.20	1.3	0.19
X(c). Malignant gonadal germ cell tumors	10	0.4	0.12	0.4	0.12	52	1.5	0.20	1.5	0.20
XI. Other malignant epithelial neoplasms and melanomas	111	4.4	0.42	4.4	0.42	263	7.5	0.47	7.5	0.46
XI(b). Thyroid carcinomas	23	0.9	0.19	0.9	0.19	54	1.6	0.21	1.5	0.21
XI(d). Malignant melanomas	13	0.5	0.14	0.5	0.14	16	0.5	0.12	0.5	0.12
XII. Other & unspecified malignant neoplasms	11	0.4	0.13	0.4	0.13	19	0.5	0.12	0.5	0.12

[1] Rates are **per million** population and are age-adjusted by five-year age groups to the 2000 U.S. standard population based on single years of age and to the 1996 Canadian population standard.
[2] See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.
[3] Abbreviations: S.E. - Standard Error; CNS - Central Nervous System.
- Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for All Sites combined.

Total Cancer Cases and Average Annual Age-adjusted Incidence Rates[1] for Pediatric Cancers
NAACCR United States Combined[2], 2010-2014, Black, Females

	Ages 0-14					Ages 0-19				
		2000 U.S.		1996 Cdn			2000 U.S.		1996 Cdn	
Cancer Types	*Count*	*Rate*	*S.E.*[3]	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Rate*	*S.E.*
All Sites	**3,011**	**122.4**	**2.23**	**122.5**	**2.23**	**4,345**	**130.1**	**1.97**	**130.2**	**1.98**
I. Leukemias	686	27.8	1.06	28.0	1.07	870	26.1	0.89	26.2	0.89
I(a). Lymphoid leukemias	440	17.8	0.85	18.0	0.86	491	14.8	0.67	14.9	0.67
I(b). Acute myeloid leukemias	152	6.2	0.50	6.2	0.50	229	6.8	0.45	6.8	0.45
II. Lymphomas & reticuloendothelial neoplasms	281	11.6	0.69	11.5	0.69	602	17.9	0.73	17.9	0.73
II(a). Hodgkin lymphomas	100	4.1	0.41	4.1	0.41	310	9.2	0.52	9.1	0.52
II(b, c, e). Non-Hodgkin lymphomas	146	6.0	0.50	6.0	0.50	257	7.7	0.48	7.7	0.48
III. CNS[3] & misc. intracranial & intraspinal neoplasms	663	27.0	1.05	27.1	1.05	799	24.2	0.85	24.2	0.86
III(a). Ependymomas and choroids plexus tumor	57	2.3	0.31	2.3	0.31	67	2.0	0.25	2.0	0.25
III(b). Astrocytomas	325	13.3	0.74	13.3	0.74	399	12.1	0.60	12.1	0.61
III(c). Intracranial and intraspinal embryonal tumors	106	4.3	0.42	4.3	0.42	121	3.6	0.33	3.7	0.33
III(d). Other gliomas	137	5.6	0.48	5.6	0.48	166	5.0	0.39	5.1	0.39
IV. Neuroblastoma and other peripheral nervous cell tumors	239	9.5	0.62	9.6	0.62	244	7.3	0.47	7.3	0.47
IV(a). Neuroblastoma & ganglioneuroblastoma	234	9.3	0.61	9.4	0.61	237	7.1	0.46	7.1	0.46
V. Retinoblastoma	121	4.8	0.44	4.8	0.44	121	3.6	0.33	3.6	0.33
VI. Renal Tumors	286	11.5	0.68	11.7	0.69	306	9.2	0.53	9.3	0.53
VI(a). Nephroblastoma and other nonepithelial renal tumors	270	10.8	0.66	11.0	0.67	272	8.2	0.50	8.3	0.50
VII. Hepatic tumors	43	1.7	0.26	1.7	0.26	54	1.6	0.22	1.6	0.22
VII(a). Hepatoblastoma	32	1.3	0.23	1.3	0.23	32	1.0	0.17	1.0	0.17
VIII. Malignant bone tumors	142	5.9	0.49	5.8	0.49	221	6.7	0.45	6.6	0.45
VIII(a). Osteosarcomas	115	4.8	0.45	4.7	0.44	172	5.2	0.40	5.2	0.40
VIII(c). Ewing tumor and related sarcomas of bone	14	0.6	0.15	0.6	0.15	21	0.6	0.14	0.6	0.14
IX. Soft tissue and other extraosseous sarcomas	242	9.9	0.64	9.9	0.64	369	11.1	0.58	11.1	0.58
IX(a). Rhabdomyosarcomas	102	4.1	0.41	4.2	0.41	132	4.0	0.35	4.0	0.35
X. Germ cell & trophoblastic tumors & neoplasms of gonads	159	6.5	0.52	6.4	0.51	295	8.8	0.51	8.7	0.51
X(a). Intracranial & intraspinal germ cell tumors	19	0.8	0.18	0.8	0.18	23	0.7	0.15	0.7	0.15
X(c). Malignant gonadal germ cell tumors	87	3.6	0.39	3.6	0.38	166	5.0	0.39	4.9	0.38
XI. Other malignant epithelial neoplasms and melanomas	123	5.1	0.46	5.0	0.45	420	12.4	0.60	12.3	0.60
XI(b). Thyroid carcinomas	44	1.8	0.28	1.8	0.27	186	5.5	0.40	5.4	0.40
XI(d). Malignant melanomas	17	0.7	0.17	0.7	0.17	23	0.7	0.14	0.7	0.14
XII. Other & unspecified malignant neoplasms	26	1.0	0.21	1.1	0.21	44	1.3	0.20	1.3	0.20

[1] Rates are **per million** population and are age-adjusted by five-year age groups to the 2000 U.S. standard population based on single years of age and to the 1996 Canadian population standard.
[2] See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.
[3] Abbreviations: S.E. - Standard Error; CNS - Central Nervous System.
- Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for All Sites combined.

Total Cancer Cases and Average Annual Age-adjusted Incidence Rates[1] for Pediatric Cancers
NAACCR United States Combined[2], 2010-2014, Asian/Pacific Islander, Males

	Ages 0-14					Ages 0-19				
		2000 U.S.		1996 Cdn			2000 U.S.		1996 Cdn	
Cancer Types	*Count*	*Rate*	*S.E.*[3]	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Rate*	*S.E.*
All Sites	**1,292**	**145.4**	**4.05**	**145.7**	**4.06**	**1,812**	**153.1**	**3.60**	**153.3**	**3.60**
I. Leukemias	451	50.4	2.38	50.8	2.39	559	46.9	1.99	47.2	2.00
I(a). Lymphoid leukemias	324	36.1	2.01	36.6	2.03	379	31.7	1.63	32.0	1.65
I(b). Acute myeloid leukemias	85	9.5	1.04	9.5	1.04	116	9.8	0.91	9.8	0.91
II. Lymphomas & reticuloendothelial neoplasms	210	23.8	1.65	23.8	1.64	331	28.1	1.55	28.1	1.55
II(a). Hodgkin lymphomas	46	5.3	0.79	5.3	0.78	94	8.1	0.83	8.1	0.83
II(b, c, e). Non-Hodgkin lymphomas	111	12.6	1.20	12.7	1.20	180	15.3	1.14	15.3	1.14
III. CNS[3] & misc. intracranial & intraspinal neoplasms	225	25.4	1.70	25.4	1.70	263	22.3	1.37	22.3	1.37
III(a). Ependymomas and choroids plexus tumor	28	3.1	0.58	3.1	0.59	34	2.8	0.48	2.8	0.49
III(b). Astrocytomas	92	10.5	1.09	10.5	1.09	118	10.1	0.93	10.1	0.93
III(c). Intracranial and intraspinal embryonal tumors	62	7.0	0.88	7.0	0.89	63	5.3	0.67	5.3	0.67
III(d). Other gliomas	37	4.2	0.69	4.2	0.69	41	3.5	0.54	3.5	0.55
IV. Neuroblastoma and other peripheral nervous cell tumors	84	9.2	1.00	9.3	1.01	84	6.9	0.75	6.9	0.76
IV(a). Neuroblastoma & ganglioneuroblastoma	84	9.2	1.00	9.3	1.01	84	6.9	0.75	6.9	0.76
V. Retinoblastoma	37	4.0	0.66	4.0	0.66	37	3.0	0.50	3.0	0.50
VI. Renal Tumors	27	3.0	0.57	3.0	0.58	35	2.9	0.49	2.9	0.50
VI(a). Nephroblastoma and other nonepithelial renal tumors	24	2.6	0.53	2.7	0.54	30	2.5	0.45	2.5	0.46
VII. Hepatic tumors	30	3.3	0.60	3.3	0.61	35	2.9	0.49	2.9	0.49
VII(a). Hepatoblastoma	28	3.0	0.58	3.1	0.58	28	2.3	0.43	2.3	0.44
VIII. Malignant bone tumors	54	6.3	0.86	6.2	0.85	95	8.2	0.84	8.2	0.84
VIII(a). Osteosarcomas	32	3.7	0.66	3.7	0.66	53	4.6	0.63	4.6	0.63
VIII(c). Ewing tumor and related sarcomas of bone	14	1.6	0.44	1.6	0.43	31	2.7	0.48	2.7	0.48
IX. Soft tissue and other extraosseous sarcomas	67	7.6	0.93	7.6	0.92	105	8.9	0.87	8.9	0.87
IX(a). Rhabdomyosarcomas	32	3.6	0.64	3.6	0.64	38	3.2	0.52	3.2	0.52
X. Germ cell & trophoblastic tumors & neoplasms of gonads	61	6.9	0.89	6.9	0.88	152	12.9	1.05	12.9	1.05
X(a). Intracranial & intraspinal germ cell tumors	31	3.6	0.65	3.6	0.64	61	5.3	0.67	5.2	0.67
X(c). Malignant gonadal germ cell tumors	21	2.3	0.51	2.3	0.51	68	5.7	0.70	5.7	0.70
XI. Other malignant epithelial neoplasms and melanomas	43	5.0	0.77	5.0	0.76	113	9.7	0.92	9.7	0.91
XI(b). Thyroid carcinomas	15	1.8	0.46	1.8	0.45	41	3.5	0.55	3.5	0.55
XI(d). Malignant melanomas	7	0.8	0.30	0.8	0.30	11	0.9	0.28	0.9	0.28
XII. Other & unspecified malignant neoplasms	-	-	-	-	-	-	-	-	-	-

[1] Rates are **per million** population and are age-adjusted by five-year age groups to the 2000 U.S. standard population based on single years of age and to the 1996 Canadian population standard.
[2] See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.
[3] Abbreviations: S.E. - Standard Error; CNS - Central Nervous System.
- Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for All Sites combined.

Total Cancer Cases and Average Annual Age-adjusted Incidence Rates[1] for Pediatric Cancers
NAACCR United States Combined[2], 2010-2014, Asian/Pacific Islander, Females

	Ages 0-14					Ages 0-19				
		2000 U.S.		1996 Cdn			2000 U.S.		1996 Cdn	
Cancer Types	*Count*	*Rate*	*S.E.*[3]	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Rate*	*S.E.*
All Sites	**1,124**	**130.3**	**3.89**	**130.4**	**3.89**	**1,599**	**139.4**	**3.49**	**139.6**	**3.49**
I. Leukemias	391	45.0	2.28	45.4	2.30	467	40.4	1.87	40.7	1.88
I(a). Lymphoid leukemias	277	31.8	1.91	32.2	1.93	306	26.3	1.51	26.7	1.52
I(b). Acute myeloid leukemias	75	8.7	1.00	8.7	1.00	107	9.3	0.90	9.3	0.90
II. Lymphomas & reticuloendothelial neoplasms	110	12.8	1.22	12.8	1.22	202	17.7	1.25	17.7	1.25
II(a). Hodgkin lymphomas	14	1.7	0.45	1.7	0.44	59	5.2	0.68	5.2	0.68
II(b, c, e). Non-Hodgkin lymphomas	66	7.7	0.95	7.7	0.95	110	9.6	0.92	9.7	0.92
III. CNS[3] & misc. intracranial & intraspinal neoplasms	183	21.2	1.57	21.3	1.57	225	19.6	1.31	19.6	1.31
III(a). Ependymomas and choroids plexus tumor	13	1.5	0.41	1.5	0.42	16	1.4	0.35	1.4	0.35
III(b). Astrocytomas	90	10.4	1.10	10.5	1.10	114	9.9	0.93	10.0	0.93
III(c). Intracranial and intraspinal embryonal tumors	47	5.4	0.79	5.4	0.79	49	4.2	0.61	4.3	0.61
III(d). Other gliomas	26	3.0	0.59	3.0	0.59	39	3.4	0.55	3.4	0.55
IV. Neuroblastoma and other peripheral nervous cell tumors	84	9.6	1.05	9.6	1.05	89	7.6	0.81	7.6	0.81
IV(a). Neuroblastoma & ganglioneuroblastoma	83	9.5	1.04	9.5	1.04	85	7.3	0.79	7.3	0.79
V. Retinoblastoma	47	5.4	0.78	5.4	0.78	47	4.0	0.58	4.0	0.59
VI. Renal Tumors	38	4.3	0.70	4.4	0.71	40	3.4	0.54	3.4	0.54
VI(a). Nephroblastoma and other nonepithelial renal tumors	35	4.0	0.68	4.0	0.68	35	3.0	0.51	3.0	0.51
VII. Hepatic tumors	23	2.6	0.55	2.6	0.55	26	2.2	0.44	2.2	0.44
VII(a). Hepatoblastoma	20	2.3	0.51	2.3	0.51	20	1.7	0.38	1.7	0.38
VIII. Malignant bone tumors	44	5.2	0.78	5.2	0.78	68	6.0	0.73	6.0	0.73
VIII(a). Osteosarcomas	26	3.1	0.60	3.1	0.60	39	3.5	0.55	3.4	0.55
VIII(c). Ewing tumor and related sarcomas of bone	17	2.0	0.48	2.0	0.48	24	2.1	0.43	2.1	0.43
IX. Soft tissue and other extraosseous sarcomas	66	7.7	0.95	7.7	0.95	93	8.1	0.85	8.1	0.84
IX(a). Rhabdomyosarcomas	32	3.7	0.65	3.7	0.66	39	3.4	0.54	3.4	0.54
X. Germ cell & trophoblastic tumors & neoplasms of gonads	73	8.6	1.01	8.5	1.00	114	10.1	0.94	10.0	0.94
X(a). Intracranial & intraspinal germ cell tumors	16	1.9	0.47	1.9	0.47	20	1.8	0.40	1.8	0.39
X(c). Malignant gonadal germ cell tumors	33	3.9	0.69	3.9	0.68	55	4.9	0.66	4.9	0.66
XI. Other malignant epithelial neoplasms and melanomas	62	7.4	0.94	7.3	0.93	223	19.8	1.32	19.7	1.32
XI(b). Thyroid carcinomas	41	4.9	0.77	4.9	0.76	170	15.1	1.16	15.0	1.15
XI(d). Malignant melanomas	-	-	-	-	-	7	0.6	0.23	0.6	0.23
XII. Other & unspecified malignant neoplasms	-	-	-	-	-	-	-	-	-	-

[1] Rates are **per million** population and are age-adjusted by five-year age groups to the 2000 U.S. standard population based on single years of age and to the 1996 Canadian population standard.
[2] See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.
[3] Abbreviations: S.E. - Standard Error; CNS - Central Nervous System.
- Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for All Sites combined.

Total Cancer Cases and Average Annual Age-adjusted Incidence Rates[1] for Pediatric Cancers
NAACCR United States Combined[2], 2010-2014, American Indian/Alaskan Native for CHSDA[3] Counties, Males

	Ages 0-14					Ages 0-19				
		2000 U.S.		1996 Cdn			2000 U.S.		1996 Cdn	
Cancer Types	*Count*	*Rate*	*S.E.*[4]	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Rate*	*S.E.*
All Sites	**201**	**142.9**	**10.09**	**143.4**	**10.13**	**276**	**149.4**	**9.01**	**149.8**	**9.03**
I. Leukemias	64	45.2	5.66	45.7	5.72	76	40.6	4.67	41.0	4.71
I(a). Lymphoid leukemias	43	30.3	4.62	30.8	4.69	50	26.6	3.77	27.0	3.82
I(b). Acute myeloid leukemias	12	8.4	2.42	8.4	2.43	17	9.1	2.21	9.1	2.22
II. Lymphomas & reticuloendothelial neoplasms	32	23.4	4.15	23.5	4.15	45	24.9	3.72	24.9	3.72
II(a). Hodgkin lymphomas	-	-	-	-	-	8	4.5	1.58	4.5	1.58
II(b, c, e). Non-Hodgkin lymphomas	18	13.4	3.16	13.3	3.14	26	14.5	2.85	14.5	2.84
III. CNS[4] & misc. intracranial & intraspinal neoplasms	44	31.7	4.79	31.8	4.80	54	29.4	4.01	29.4	4.01
III(a). Ependymomas and choroids plexus tumor	7	5.1	1.92	5.0	1.91	8	4.4	1.55	4.3	1.54
III(b). Astrocytomas	19	13.7	3.15	13.7	3.15	26	14.2	2.79	14.2	2.80
III(c). Intracranial and intraspinal embryonal tumors	6	4.2	1.72	4.2	1.73	6	3.2	1.29	3.2	1.29
III(d). Other gliomas	7	5.1	1.94	5.1	1.94	8	4.4	1.56	4.4	1.56
IV. Neuroblastoma and other peripheral nervous cell tumors	11	7.5	2.25	7.5	2.27	13	6.7	1.87	6.8	1.88
IV(a). Neuroblastoma & ganglioneuroblastoma	11	7.5	2.25	7.5	2.27	12	6.2	1.78	6.2	1.79
V. Retinoblastoma	-	-	-	-	-	-	-	-	-	-
VI. Renal Tumors	7	4.8	1.80	4.8	1.83	7	3.6	1.35	3.6	1.37
VI(a). Nephroblastoma and other nonepithelial renal tumors	7	4.8	1.80	4.8	1.83	7	3.6	1.35	3.6	1.37
VII. Hepatic tumors	10	6.7	2.10	6.6	2.07	10	5.0	1.58	4.9	1.55
VII(a). Hepatoblastoma	10	6.7	2.10	6.6	2.07	10	5.0	1.58	4.9	1.55
VIII. Malignant bone tumors	11	8.0	2.43	8.1	2.43	16	8.8	2.21	8.9	2.22
VIII(a). Osteosarcomas	-	-	-	-	-	7	4.0	1.49	3.9	1.49
VIII(c). Ewing tumor and related sarcomas of bone	7	5.0	1.90	5.1	1.92	8	4.3	1.53	4.4	1.54
IX. Soft tissue and other extraosseous sarcomas	7	5.0	1.91	5.0	1.90	15	8.3	2.15	8.3	2.14
IX(a). Rhabdomyosarcomas	-	-	-	-	-	7	3.9	1.46	3.9	1.47
X. Germ cell & trophoblastic tumors & neoplasms of gonads	7	4.9	1.85	4.9	1.84	20	11.0	2.47	11.0	2.46
X(a). Intracranial & intraspinal germ cell tumors	-	-	-	-	-	-	-	-	-	-
X(c). Malignant gonadal germ cell tumors	-	-	-	-	-	14	7.8	2.09	7.8	2.10
XI. Other malignant epithelial neoplasms and melanomas	-	-	-	-	-	15	8.5	2.20	8.5	2.19
XI(b). Thyroid carcinomas	-	-	-	-	-	-	-	-	-	-
XI(d). Malignant melanomas	-	-	-	-	-	-	-	-	-	-
XII. Other & unspecified malignant neoplasms	-	-	-	-	-	-	-	-	-	-

[1] Rates are **per million** population and are age-adjusted by five-year age groups to the 2000 U.S. standard population based on single years of age and to the 1996 Canadian population standard.
[2] See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.
[3] CHSDA - Indian Health Services Contract Health Service Delivery Areas (See Appendix G)
[4] Abbreviations: S.E. - Standard Error; CNS - Central Nervous System.
- Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for All Sites combined. Statistics exclude cases diagnosed for KS for all years.

Total Cancer Cases and Average Annual Age-adjusted Incidence Rates[1] for Pediatric Cancers
NAACCR United States Combined[2], 2010-2014, American Indian/Alaskan Native for CHSDA[3] Counties, Females

	Ages 0-14					Ages 0-19				
		2000 U.S.		1996 Cdn			2000 U.S.		1996 Cdn	
Cancer Types	*Count*	*Rate*	*S.E.*[4]	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Rate*	*S.E.*
All Sites	**148**	**109.7**	**9.03**	**109.4**	**9.01**	**214**	**121.1**	**8.29**	**120.8**	**8.27**
I. Leukemias	46	34.0	5.02	34.1	5.04	63	35.5	4.48	35.6	4.49
I(a). Lymphoid leukemias	31	22.8	4.10	23.1	4.14	33	18.2	3.18	18.4	3.21
I(b). Acute myeloid leukemias	8	6.1	2.16	6.1	2.16	20	11.6	2.61	11.6	2.60
II. Lymphomas & reticuloendothelial neoplasms	11	8.4	2.52	8.3	2.52	22	12.7	2.72	12.7	2.72
II(a). Hodgkin lymphomas	-	-	-	-	-	6	3.5	1.42	3.5	1.42
II(b, c, e). Non-Hodgkin lymphomas	-	-	-	-	-	10	5.9	1.85	5.8	1.85
III. CNS[4] & misc. intracranial & intraspinal neoplasms	29	21.4	3.98	21.4	3.98	34	19.0	3.26	19.0	3.26
III(a). Ependymomas and choroids plexus tumor	-	-	-	-	-	-	-	-	-	-
III(b). Astrocytomas	12	9.0	2.60	9.0	2.60	15	8.5	2.20	8.5	2.20
III(c). Intracranial and intraspinal embryonal tumors	6	4.2	1.69	4.2	1.71	6	3.1	1.27	3.1	1.28
III(d). Other gliomas	7	5.3	2.02	5.3	2.01	9	5.2	1.73	5.2	1.72
IV. Neuroblastoma and other peripheral nervous cell tumors	6	4.2	1.69	4.2	1.71	6	3.1	1.27	3.1	1.28
IV(a). Neuroblastoma & ganglioneuroblastoma	6	4.2	1.69	4.2	1.71	6	3.1	1.27	3.1	1.28
V. Retinoblastoma	-	-	-	-	-	-	-	-	-	-
VI. Renal Tumors	6	4.3	1.77	4.2	1.73	6	3.2	1.32	3.2	1.29
VI(a). Nephroblastoma and other nonepithelial renal tumors	6	4.3	1.77	4.2	1.73	6	3.2	1.32	3.2	1.29
VII. Hepatic tumors	-	-	-	-	-	6	3.2	1.29	3.1	1.26
VII(a). Hepatoblastoma	-	-	-	-	-	-	-	-	-	-
VIII. Malignant bone tumors	9	7.0	2.32	6.9	2.30	12	7.0	2.02	6.9	2.00
VIII(a). Osteosarcomas	7	5.5	2.08	5.4	2.05	10	5.9	1.86	5.8	1.84
VIII(c). Ewing tumor and related sarcomas of bone	-	-	-	-	-	-	-	-	-	-
IX. Soft tissue and other extraosseous sarcomas	15	11.3	2.92	11.3	2.91	22	12.6	2.69	12.6	2.68
IX(a). Rhabdomyosarcomas	7	5.2	1.98	5.2	1.98	9	5.1	1.70	5.1	1.70
X. Germ cell & trophoblastic tumors & neoplasms of gonads	-	-	-	-	-	7	4.0	1.53	4.0	1.51
X(a). Intracranial & intraspinal germ cell tumors	-	-	-	-	-	-	-	-	-	-
X(c). Malignant gonadal germ cell tumors	-	-	-	-	-	6	3.5	1.44	3.5	1.43
XI. Other malignant epithelial neoplasms and melanomas	11	8.6	2.58	8.5	2.56	30	17.6	3.22	17.5	3.20
XI(b). Thyroid carcinomas	-	-	-	-	-	15	8.8	2.28	8.8	2.27
XI(d). Malignant melanomas	-	-	-	-	-	-	-	-	-	-
XII. Other & unspecified malignant neoplasms	-	-	-	-	-	-	-	-	-	-

[1] Rates are **per million** population and are age-adjusted by five-year age groups to the 2000 U.S. standard population based on single years of age and to the 1996 Canadian population standard.
[2] See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.
[3] CHSDA - Indian Health Services Contract Health Service Delivery Areas (See Appendix G)
[4] Abbreviations: S.E. - Standard Error; CNS - Central Nervous System.
- Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for All Sites combined. Statistics exclude cases diagnosed for KS for all years.

Total Cancer Cases and Average Annual Age-adjusted Incidence Rates[1] for Pediatric Cancers
NAACCR United States Combined[2], 2010-2014, Hispanic/Latino, All Races, Males

	Ages 0-14					Ages 0-19				
		2000 U.S.		1996 Cdn			2000 U.S.		1996 Cdn	
Cancer Types	*Count*	*Rate*	*S.E.*[3]	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Rate*	*S.E.*
All Sites	**6,347**	**172.5**	**2.17**	**172.9**	**2.17**	**8,976**	**187.4**	**1.98**	**187.7**	**1.98**
I. Leukemias	2,465	66.7	1.35	67.1	1.35	3,087	63.7	1.15	64.0	1.15
I(a). Lymphoid leukemias	1,909	51.5	1.18	52.0	1.19	2,308	47.4	0.99	47.8	1.00
I(b). Acute myeloid leukemias	328	8.9	0.49	8.9	0.49	455	9.5	0.45	9.5	0.45
II. Lymphomas & reticuloendothelial neoplasms	965	26.7	0.86	26.7	0.86	1,431	30.3	0.80	30.3	0.80
II(a). Hodgkin lymphomas	260	7.4	0.46	7.4	0.46	510	11.1	0.49	11.1	0.49
II(b, c, e). Non-Hodgkin lymphomas	408	11.4	0.56	11.3	0.56	606	12.9	0.52	12.9	0.52
III. CNS[3] & misc. intracranial & intraspinal neoplasms	1,019	27.7	0.87	27.8	0.87	1,224	25.3	0.72	25.4	0.73
III(a). Ependymomas and choroids plexus tumor	111	3.0	0.28	3.0	0.29	125	2.5	0.23	2.6	0.23
III(b). Astrocytomas	457	12.5	0.59	12.6	0.59	574	12.0	0.50	12.0	0.50
III(c). Intracranial and intraspinal embryonal tumors	269	7.2	0.44	7.3	0.44	293	5.9	0.35	6.0	0.35
III(d). Other gliomas	144	3.9	0.33	4.0	0.33	181	3.8	0.28	3.8	0.28
IV. Neuroblastoma and other peripheral nervous cell tumors	299	7.7	0.45	7.7	0.45	308	6.0	0.34	6.0	0.34
IV(a). Neuroblastoma & ganglioneuroblastoma	290	7.5	0.44	7.5	0.44	295	5.7	0.33	5.7	0.33
V. Retinoblastoma	183	4.7	0.35	4.7	0.35	183	3.5	0.26	3.5	0.26
VI. Renal Tumors	236	6.2	0.40	6.3	0.41	248	4.9	0.31	5.0	0.32
VI(a). Nephroblastoma and other nonepithelial renal tumors	221	5.8	0.39	5.9	0.39	223	4.4	0.29	4.4	0.30
VII. Hepatic tumors	163	4.2	0.33	4.3	0.33	175	3.4	0.26	3.5	0.26
VII(a). Hepatoblastoma	139	3.6	0.30	3.6	0.31	140	2.7	0.23	2.7	0.23
VIII. Malignant bone tumors	249	7.2	0.46	7.1	0.45	438	9.6	0.46	9.5	0.46
VIII(a). Osteosarcomas	145	4.2	0.35	4.2	0.35	265	5.8	0.36	5.8	0.35
VIII(c). Ewing tumor and related sarcomas of bone	86	2.5	0.27	2.4	0.26	128	2.8	0.25	2.8	0.24
IX. Soft tissue and other extraosseous sarcomas	376	10.3	0.53	10.3	0.53	544	11.4	0.49	11.4	0.49
IX(a). Rhabdomyosarcomas	179	4.8	0.36	4.9	0.36	217	4.5	0.30	4.5	0.30
X. Germ cell & trophoblastic tumors & neoplasms of gonads	197	5.5	0.39	5.4	0.38	914	20.0	0.66	19.9	0.66
X(a). Intracranial & intraspinal germ cell tumors	86	2.5	0.27	2.4	0.26	133	2.9	0.25	2.9	0.25
X(c). Malignant gonadal germ cell tumors	77	2.1	0.24	2.1	0.23	685	15.0	0.57	15.0	0.57
XI. Other malignant epithelial neoplasms and melanomas	173	4.9	0.38	4.9	0.37	398	8.7	0.44	8.7	0.43
XI(b). Thyroid carcinomas	53	1.5	0.21	1.5	0.21	142	3.1	0.26	3.1	0.26
XI(d). Malignant melanomas	35	1.0	0.16	1.0	0.16	51	1.1	0.15	1.1	0.15
XII. Other & unspecified malignant neoplasms	22	0.6	0.12	0.6	0.13	26	0.5	0.10	0.5	0.10

[1] Rates are **per million** population and are age-adjusted by five-year age groups to the 2000 U.S. standard population based on single years of age and to the 1996 Canadian population standard.
[2] See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.
[3] Abbreviations: S.E. - Standard Error; CNS - Central Nervous System.
- Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for All Sites combined.

N.B. In areas with small Latino populations, methods to indirectly identify Latinos (like NHIAv2) can overestimate the ethnicity-specific counts of cancer cases. Also, even small errors in Latino population estimates can affect the magnitude of the cancer rates. Data Item 190 used in place of NHIA for MA and RI, see technical notes.

Total Cancer Cases and Average Annual Age-adjusted Incidence Rates[1] for Pediatric Cancers
NAACCR United States Combined[2], 2010-2014, Hispanic/Latina, All Races, Females

	Ages 0-14					Ages 0-19				
		2000 U.S.		1996 Cdn			2000 U.S.		1996 Cdn	
Cancer Types	*Count*	*Rate*	*S.E.*[3]	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Rate*	*S.E.*
All Sites	**5,554**	**156.9**	**2.11**	**157.4**	**2.11**	**7,633**	**166.3**	**1.91**	**166.7**	**1.91**
I. Leukemias	2,036	57.0	1.27	57.5	1.28	2,389	51.0	1.05	51.3	1.05
I(a). Lymphoid leukemias	1,593	44.5	1.12	45.0	1.13	1,781	37.7	0.90	38.1	0.90
I(b). Acute myeloid leukemias	277	7.8	0.47	7.8	0.47	381	8.3	0.43	8.3	0.43
II. Lymphomas & reticuloendothelial neoplasms	589	17.0	0.70	17.0	0.70	966	21.6	0.70	21.5	0.69
II(a). Hodgkin lymphomas	145	4.4	0.36	4.3	0.36	398	9.2	0.46	9.2	0.46
II(b, c, e). Non-Hodgkin lymphomas	231	6.7	0.44	6.7	0.44	337	7.5	0.41	7.5	0.41
III. CNS[3] & misc. intracranial & intraspinal neoplasms	995	28.0	0.89	28.1	0.89	1,163	24.9	0.73	25.0	0.73
III(a). Ependymomas and choroids plexus tumor	109	3.0	0.29	3.0	0.29	124	2.6	0.23	2.6	0.24
III(b). Astrocytomas	491	13.9	0.63	14.0	0.63	590	12.7	0.53	12.8	0.53
III(c). Intracranial and intraspinal embryonal tumors	199	5.5	0.39	5.5	0.39	219	4.6	0.31	4.6	0.31
III(d). Other gliomas	161	4.6	0.36	4.6	0.36	187	4.0	0.30	4.1	0.30
IV. Neuroblastoma and other peripheral nervous cell tumors	269	7.2	0.44	7.3	0.44	271	5.5	0.33	5.5	0.33
IV(a). Neuroblastoma & ganglioneuroblastoma	266	7.2	0.44	7.2	0.44	267	5.4	0.33	5.4	0.33
V. Retinoblastoma	166	4.4	0.34	4.4	0.34	166	3.3	0.26	3.3	0.26
VI. Renal Tumors	276	7.5	0.45	7.6	0.46	292	6.0	0.35	6.1	0.36
VI(a). Nephroblastoma and other nonepithelial renal tumors	266	7.2	0.44	7.3	0.45	270	5.5	0.33	5.6	0.34
VII. Hepatic tumors	96	2.6	0.27	2.6	0.27	105	2.2	0.21	2.2	0.21
VII(a). Hepatoblastoma	82	2.2	0.24	2.2	0.24	84	1.7	0.19	1.7	0.19
VIII. Malignant bone tumors	249	7.4	0.47	7.4	0.47	341	7.7	0.42	7.7	0.42
VIII(a). Osteosarcomas	153	4.6	0.37	4.6	0.37	203	4.6	0.32	4.6	0.32
VIII(c). Ewing tumor and related sarcomas of bone	70	2.1	0.25	2.0	0.24	97	2.2	0.22	2.2	0.22
IX. Soft tissue and other extraosseous sarcomas	334	9.6	0.53	9.6	0.52	482	10.7	0.49	10.6	0.49
IX(a). Rhabdomyosarcomas	144	4.1	0.34	4.1	0.34	179	3.9	0.29	3.9	0.29
X. Germ cell & trophoblastic tumors & neoplasms of gonads	237	6.9	0.45	6.9	0.45	450	10.2	0.48	10.1	0.48
X(a). Intracranial & intraspinal germ cell tumors	34	1.0	0.17	1.0	0.17	41	0.9	0.14	0.9	0.14
X(c). Malignant gonadal germ cell tumors	138	4.1	0.35	4.1	0.35	275	6.3	0.38	6.3	0.38
XI. Other malignant epithelial neoplasms and melanomas	289	8.7	0.51	8.6	0.51	973	22.6	0.72	22.5	0.72
XI(b). Thyroid carcinomas	151	4.6	0.37	4.5	0.37	646	15.1	0.59	15.0	0.59
XI(d). Malignant melanomas	50	1.5	0.21	1.5	0.21	76	1.7	0.20	1.7	0.20
XII. Other & unspecified malignant neoplasms	18	0.5	0.12	0.5	0.12	35	0.8	0.13	0.8	0.13

[1] Rates are **per million** population and are age-adjusted by five-year age groups to the 2000 U.S. standard population based on single years of age and to the 1996 Canadian population standard.
[2] See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.
[3] Abbreviations: S.E. - Standard Error; CNS - Central Nervous System.
- Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for All Sites combined.

N.B. In areas with small Latino populations, methods to indirectly identify Latinos (like NHIAv2) can overestimate the ethnicity-specific counts of cancer cases. Also, even small errors in Latino population estimates can affect the magnitude of the cancer rates. Data Item 190 used in place of NHIA for MA and RI, see technical notes.

Total Cancer Cases and Average Annual Age-adjusted Incidence Rates[1] for Pediatric Cancers
NAACCR United States Combined[2], 2010-2014, Non-Hispanic White, Males

	Ages 0-14					Ages 0-19				
		2000 U.S.		1996 Cdn			2000 U.S.		1996 Cdn	
Cancer Types	*Count*	*Rate*	*S.E.*[3]	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Rate*	*S.E.*
All Sites	**15,298**	**187.1**	**1.51**	**187.6**	**1.52**	**22,781**	**202.2**	**1.34**	**202.6**	**1.34**
I. Leukemias	4,489	55.0	0.82	55.6	0.83	5,608	50.5	0.67	50.9	0.68
I(a). Lymphoid leukemias	3,514	43.1	0.73	43.7	0.74	4,111	37.2	0.58	37.6	0.59
I(b). Acute myeloid leukemias	576	7.0	0.29	7.1	0.29	875	7.8	0.26	7.8	0.26
II. Lymphomas & reticuloendothelial neoplasms	2,171	26.4	0.57	26.3	0.56	4,122	35.9	0.56	35.9	0.56
II(a). Hodgkin lymphomas	513	6.2	0.27	6.1	0.27	1,637	13.9	0.34	13.9	0.34
II(b, c, e). Non-Hodgkin lymphomas	1,129	13.7	0.41	13.7	0.41	1,880	16.5	0.38	16.5	0.38
III. CNS[3] & misc. intracranial & intraspinal neoplasms	3,517	43.0	0.72	43.1	0.73	4,375	39.3	0.59	39.4	0.60
III(a). Ependymomas and choroids plexus tumor	313	3.8	0.22	3.9	0.22	365	3.3	0.17	3.3	0.17
III(b). Astrocytomas	1,789	21.8	0.52	21.9	0.52	2,289	20.5	0.43	20.6	0.43
III(c). Intracranial and intraspinal embryonal tumors	774	9.5	0.34	9.5	0.34	881	8.0	0.27	8.0	0.27
III(d). Other gliomas	548	6.7	0.29	6.7	0.29	710	6.3	0.24	6.3	0.24
IV. Neuroblastoma and other peripheral nervous cell tumors	1,142	14.2	0.42	14.2	0.42	1,172	10.8	0.32	10.8	0.32
IV(a). Neuroblastoma & ganglioneuroblastoma	1,120	13.9	0.42	13.9	0.42	1,135	10.5	0.31	10.5	0.31
V. Retinoblastoma	309	3.8	0.22	3.8	0.22	309	2.9	0.16	2.9	0.16
VI. Renal Tumors	710	8.8	0.33	8.9	0.33	763	7.0	0.25	7.1	0.26
VI(a). Nephroblastoma and other nonepithelial renal tumors	680	8.4	0.32	8.5	0.33	689	6.4	0.24	6.5	0.25
VII. Hepatic tumors	288	3.6	0.21	3.6	0.21	314	2.9	0.16	2.9	0.16
VII(a). Hepatoblastoma	246	3.1	0.19	3.1	0.20	247	2.3	0.15	2.3	0.15
VIII. Malignant bone tumors	669	8.0	0.31	8.0	0.31	1,206	10.5	0.30	10.4	0.30
VIII(a). Osteosarcomas	314	3.8	0.21	3.7	0.21	597	5.2	0.21	5.1	0.21
VIII(c). Ewing tumor and related sarcomas of bone	300	3.6	0.21	3.6	0.21	500	4.4	0.20	4.4	0.20
IX. Soft tissue and other extraosseous sarcomas	1,003	12.2	0.39	12.2	0.39	1,481	13.1	0.34	13.1	0.34
IX(a). Rhabdomyosarcomas	488	6.0	0.27	6.0	0.27	594	5.4	0.22	5.4	0.22
X. Germ cell & trophoblastic tumors & neoplasms of gonads	368	4.5	0.23	4.4	0.23	1,647	14.0	0.34	13.9	0.34
X(a). Intracranial & intraspinal germ cell tumors	164	2.0	0.15	2.0	0.15	279	2.4	0.15	2.4	0.14
X(c). Malignant gonadal germ cell tumors	116	1.4	0.13	1.4	0.13	1,220	10.2	0.29	10.2	0.29
XI. Other malignant epithelial neoplasms and melanomas	571	6.9	0.29	6.8	0.28	1,708	14.6	0.35	14.5	0.35
XI(b). Thyroid carcinomas	163	2.0	0.15	1.9	0.15	491	4.2	0.19	4.2	0.19
XI(d). Malignant melanomas	211	2.6	0.18	2.5	0.17	606	5.2	0.21	5.2	0.21
XII. Other & unspecified malignant neoplasms	61	0.7	0.10	0.8	0.10	76	0.7	0.08	0.7	0.08

[1] Rates are **per million** population and are age-adjusted by five-year age groups to the 2000 U.S. standard population based on single years of age and to the 1996 Canadian population standard.
[2] See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.
[3] Abbreviations: S.E. - Standard Error; CNS - Central Nervous System.
- Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for All Sites combined.

N.B. In areas with small Latino populations, methods to indirectly identify Latinos (like NHIAv2) can overestimate the ethnicity-specific counts of cancer cases. Also, even small errors in Latino population estimates can affect the magnitude of the cancer rates. Data Item 190 used in place of NHIA for MA and RI, see technical notes.

Total Cancer Cases and Average Annual Age-adjusted Incidence Rates[1] for Pediatric Cancers
NAACCR United States Combined[2], 2010-2014, Non-Hispanic White, Females

	Ages 0-14					Ages 0-19				
		2000 U.S.		1996 Cdn			2000 U.S.		1996 Cdn	
Cancer Types	*Count*	*Rate*	*S.E.*[3]	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Rate*	*S.E.*
All Sites	**13,202**	**169.7**	**1.48**	**170.0**	**1.48**	**20,052**	**187.1**	**1.32**	**187.3**	**1.32**
I. Leukemias	3,810	49.1	0.80	49.6	0.80	4,567	43.4	0.64	43.8	0.65
I(a). Lymphoid leukemias	2,888	37.3	0.69	37.8	0.70	3,158	30.3	0.54	30.6	0.55
I(b). Acute myeloid leukemias	565	7.3	0.31	7.3	0.31	854	8.0	0.27	8.0	0.27
II. Lymphomas & reticuloendothelial neoplasms	1,276	16.3	0.46	16.2	0.45	2,774	25.3	0.48	25.3	0.48
II(a). Hodgkin lymphomas	344	4.3	0.23	4.3	0.23	1,437	12.8	0.34	12.8	0.34
II(b, c, e). Non-Hodgkin lymphomas	533	6.8	0.29	6.8	0.29	896	8.3	0.28	8.3	0.28
III. CNS[3] & misc. intracranial & intraspinal neoplasms	2,942	37.8	0.70	37.9	0.70	3,645	34.4	0.57	34.6	0.57
III(a). Ependymomas and choroids plexus tumor	225	2.9	0.19	2.9	0.20	266	2.5	0.16	2.5	0.16
III(b). Astrocytomas	1,661	21.3	0.52	21.4	0.53	2,083	19.7	0.43	19.7	0.43
III(c). Intracranial and intraspinal embryonal tumors	469	6.1	0.28	6.1	0.28	524	5.0	0.22	5.0	0.22
III(d). Other gliomas	508	6.5	0.29	6.5	0.29	669	6.3	0.24	6.3	0.24
IV. Neuroblastoma and other peripheral nervous cell tumors	1,005	13.1	0.41	13.1	0.41	1,025	10.0	0.31	10.0	0.31
IV(a). Neuroblastoma & ganglioneuroblastoma	987	12.9	0.41	12.9	0.41	999	9.7	0.31	9.7	0.31
V. Retinoblastoma	320	4.2	0.23	4.2	0.23	321	3.1	0.18	3.1	0.17
VI. Renal Tumors	776	10.1	0.36	10.2	0.37	844	8.1	0.28	8.2	0.28
VI(a). Nephroblastoma and other nonepithelial renal tumors	747	9.7	0.35	9.8	0.36	762	7.4	0.27	7.5	0.27
VII. Hepatic tumors	177	2.3	0.17	2.3	0.17	219	2.1	0.14	2.1	0.14
VII(a). Hepatoblastoma	146	1.9	0.16	1.9	0.16	147	1.4	0.12	1.4	0.12
VIII. Malignant bone tumors	581	7.4	0.31	7.3	0.30	929	8.6	0.28	8.5	0.28
VIII(a). Osteosarcomas	281	3.5	0.21	3.5	0.21	462	4.2	0.20	4.2	0.20
VIII(c). Ewing tumor and related sarcomas of bone	244	3.1	0.20	3.1	0.20	353	3.3	0.17	3.3	0.17
IX. Soft tissue and other extraosseous sarcomas	830	10.6	0.37	10.6	0.37	1,197	11.2	0.32	11.2	0.32
IX(a). Rhabdomyosarcomas	378	4.9	0.25	4.9	0.25	459	4.4	0.20	4.4	0.20
X. Germ cell & trophoblastic tumors & neoplasms of gonads	465	6.0	0.28	5.9	0.27	801	7.4	0.26	7.3	0.26
X(a). Intracranial & intraspinal germ cell tumors	89	1.1	0.12	1.1	0.12	102	1.0	0.10	1.0	0.09
X(c). Malignant gonadal germ cell tumors	185	2.3	0.17	2.3	0.17	374	3.4	0.18	3.4	0.18
XI. Other malignant epithelial neoplasms and melanomas	972	12.3	0.39	12.1	0.39	3,639	32.6	0.54	32.5	0.54
XI(b). Thyroid carcinomas	483	6.1	0.28	6.0	0.27	2,069	18.5	0.41	18.4	0.40
XI(d). Malignant melanomas	252	3.2	0.20	3.2	0.20	838	7.6	0.26	7.5	0.26
XII. Other & unspecified malignant neoplasms	48	0.6	0.09	0.6	0.09	91	0.8	0.09	0.8	0.09

[1] Rates are **per million** population and are age-adjusted by five-year age groups to the 2000 U.S. standard population based on single years of age and to the 1996 Canadian population standard.
[2] See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.
[3] Abbreviations: S.E. - Standard Error; CNS - Central Nervous System.
- Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for All Sites combined.

N.B. In areas with small Latino populations, methods to indirectly identify Latinos (like NHIAv2) can overestimate the ethnicity-specific counts of cancer cases. Also, even small errors in Latino population estimates can affect the magnitude of the cancer rates. Data Item 190 used in place of NHIA for MA and RI, see technical notes.

Total Cancer Cases and Average Annual Age-adjusted Incidence Rates[1] for Pediatric Cancers
NAACCR United States Combined[2], 2010-2014, Non-Hispanic Black, Males

	Ages 0-14					Ages 0-19				
		2000 U.S.		1996 Cdn			2000 U.S.		1996 Cdn	
Cancer Types	*Count*	*Rate*	*S.E.*[3]	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Rate*	*S.E.*
All Sites	**3,217**	**139.0**	**2.45**	**139.4**	**2.46**	**4,522**	**143.2**	**2.13**	**143.5**	**2.14**
I. Leukemias	849	36.7	1.26	36.9	1.27	1,088	34.6	1.05	34.8	1.06
I(a). Lymphoid leukemias	567	24.5	1.03	24.7	1.04	662	21.2	0.82	21.4	0.83
I(b). Acute myeloid leukemias	154	6.6	0.53	6.7	0.54	223	7.0	0.47	7.1	0.47
II. Lymphomas & reticuloendothelial neoplasms	489	21.3	0.96	21.2	0.96	888	27.9	0.94	27.8	0.94
II(a). Hodgkin lymphomas	162	7.1	0.56	7.0	0.55	356	11.1	0.59	11.1	0.59
II(b, c, e). Non-Hodgkin lymphomas	271	11.8	0.72	11.8	0.71	473	14.9	0.69	14.9	0.68
III. CNS[3] & misc. intracranial & intraspinal neoplasms	675	29.2	1.13	29.4	1.13	836	26.7	0.92	26.8	0.93
III(a). Ependymomas and choroids plexus tumor	59	2.5	0.33	2.5	0.33	72	2.3	0.27	2.3	0.27
III(b). Astrocytomas	316	13.7	0.77	13.7	0.77	403	12.9	0.64	12.9	0.64
III(c). Intracranial and intraspinal embryonal tumors	143	6.2	0.52	6.2	0.52	162	5.2	0.41	5.2	0.41
III(d). Other gliomas	124	5.4	0.49	5.4	0.49	157	5.0	0.40	5.0	0.40
IV. Neuroblastoma and other peripheral nervous cell tumors	224	9.6	0.64	9.6	0.64	233	7.4	0.49	7.5	0.49
IV(a). Neuroblastoma & ganglioneuroblastoma	217	9.3	0.63	9.3	0.63	220	7.0	0.47	7.1	0.48
V. Retinoblastoma	89	3.8	0.40	3.8	0.40	89	2.8	0.30	2.8	0.30
VI. Renal Tumors	226	9.7	0.64	9.8	0.65	247	7.9	0.50	8.0	0.51
VI(a). Nephroblastoma and other nonepithelial renal tumors	209	8.9	0.62	9.1	0.63	210	6.7	0.46	6.8	0.47
VII. Hepatic tumors	68	2.9	0.35	2.9	0.35	77	2.4	0.28	2.5	0.28
VII(a). Hepatoblastoma	62	2.6	0.33	2.7	0.34	62	2.0	0.25	2.0	0.25
VIII. Malignant bone tumors	138	6.0	0.51	6.0	0.51	240	7.6	0.49	7.5	0.49
VIII(a). Osteosarcomas	114	5.0	0.47	4.9	0.46	192	6.1	0.44	6.0	0.44
VIII(c). Ewing tumor and related sarcomas of bone	9	0.4	0.13	0.4	0.13	16	0.5	0.13	0.5	0.13
IX. Soft tissue and other extraosseous sarcomas	300	13.0	0.75	13.0	0.75	449	14.2	0.67	14.2	0.67
IX(a). Rhabdomyosarcomas	152	6.6	0.53	6.6	0.54	194	6.2	0.44	6.2	0.45
X. Germ cell & trophoblastic tumors & neoplasms of gonads	40	1.7	0.27	1.7	0.27	103	3.2	0.31	3.2	0.31
X(a). Intracranial & intraspinal germ cell tumors	18	0.8	0.18	0.8	0.18	39	1.2	0.20	1.2	0.19
X(c). Malignant gonadal germ cell tumors	7	0.3	0.11	0.3	0.11	41	1.2	0.19	1.2	0.19
XI. Other malignant epithelial neoplasms and melanomas	108	4.7	0.45	4.7	0.45	253	7.9	0.50	7.8	0.49
XI(b). Thyroid carcinomas	23	1.0	0.21	1.0	0.21	51	1.6	0.22	1.6	0.22
XI(d). Malignant melanomas	11	0.5	0.14	0.5	0.14	14	0.4	0.12	0.4	0.12
XII. Other & unspecified malignant neoplasms	11	0.5	0.14	0.5	0.14	19	0.6	0.14	0.6	0.14

[1] Rates are **per million** population and are age-adjusted by five-year age groups to the 2000 U.S. standard population based on single years of age and to the 1996 Canadian population standard.
[2] See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.
[3] Abbreviations: S.E. - Standard Error; CNS - Central Nervous System.
- Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for All Sites combined.

N.B. In areas with small Latino populations, methods to indirectly identify Latinos (like NHIAv2) can overestimate the ethnicity-specific counts of cancer cases. Also, even small errors in Latino population estimates can affect the magnitude of the cancer rates. Data Item 190 used in place of NHIA for MA and RI, see technical notes.

Total Cancer Cases and Average Annual Age-adjusted Incidence Rates[1] for Pediatric Cancers
NAACCR United States Combined[2], 2010-2014, Non-Hispanic Black, Females

	Ages 0-14					Ages 0-19				
		2000 U.S.		1996 Cdn			2000 U.S.		1996 Cdn	
Cancer Types	*Count*	*Rate*	*S.E.*[3]	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Rate*	*S.E.*
All Sites	**2,857**	**127.6**	**2.39**	**127.7**	**2.39**	**4,130**	**135.0**	**2.10**	**135.1**	**2.10**
I. Leukemias	638	28.4	1.13	28.6	1.13	810	26.6	0.94	26.7	0.94
I(a). Lymphoid leukemias	407	18.1	0.90	18.3	0.91	455	15.1	0.71	15.2	0.71
I(b). Acute myeloid leukemias	141	6.3	0.53	6.3	0.53	212	6.9	0.47	6.9	0.47
II. Lymphomas & reticuloendothelial neoplasms	269	12.1	0.74	12.0	0.73	577	18.6	0.78	18.6	0.77
II(a). Hodgkin lymphomas	94	4.2	0.44	4.2	0.43	296	9.4	0.55	9.4	0.55
II(b, c, e). Non-Hodgkin lymphomas	140	6.3	0.53	6.3	0.53	246	8.0	0.51	8.0	0.51
III. CNS[3] & misc. intracranial & intraspinal neoplasms	630	28.2	1.12	28.3	1.13	758	25.1	0.91	25.1	0.91
III(a). Ependymomas and choroids plexus tumor	55	2.4	0.33	2.5	0.33	65	2.1	0.27	2.2	0.27
III(b). Astrocytomas	307	13.8	0.79	13.8	0.79	377	12.5	0.64	12.5	0.64
III(c). Intracranial and intraspinal embryonal tumors	99	4.4	0.44	4.4	0.44	113	3.7	0.35	3.7	0.35
III(d). Other gliomas	131	5.9	0.52	5.9	0.52	157	5.2	0.42	5.2	0.42
IV. Neuroblastoma and other peripheral nervous cell tumors	227	10.0	0.67	10.1	0.67	232	7.7	0.50	7.7	0.51
IV(a). Neuroblastoma & ganglioneuroblastoma	222	9.8	0.66	9.8	0.66	225	7.4	0.50	7.5	0.50
V. Retinoblastoma	118	5.2	0.48	5.2	0.48	118	3.9	0.36	3.9	0.36
VI. Renal Tumors	276	12.2	0.74	12.4	0.75	296	9.8	0.57	9.9	0.58
VI(a). Nephroblastoma and other nonepithelial renal tumors	261	11.6	0.72	11.7	0.73	263	8.7	0.54	8.9	0.55
VII. Hepatic tumors	38	1.7	0.27	1.7	0.28	49	1.6	0.23	1.6	0.23
VII(a). Hepatoblastoma	28	1.2	0.23	1.2	0.24	28	0.9	0.17	0.9	0.18
VIII. Malignant bone tumors	137	6.2	0.53	6.1	0.52	213	7.0	0.48	7.0	0.48
VIII(a). Osteosarcomas	111	5.0	0.48	5.0	0.47	166	5.5	0.42	5.4	0.42
VIII(c). Ewing tumor and related sarcomas of bone	13	0.6	0.16	0.6	0.16	20	0.7	0.15	0.7	0.15
IX. Soft tissue and other extraosseous sarcomas	231	10.4	0.68	10.3	0.68	356	11.6	0.62	11.6	0.62
IX(a). Rhabdomyosarcomas	99	4.4	0.44	4.4	0.45	128	4.2	0.37	4.2	0.37
X. Germ cell & trophoblastic tumors & neoplasms of gonads	151	6.8	0.55	6.7	0.54	282	9.1	0.54	9.1	0.54
X(a). Intracranial & intraspinal germ cell tumors	17	0.8	0.19	0.8	0.18	21	0.7	0.15	0.7	0.15
X(c). Malignant gonadal germ cell tumors	84	3.8	0.41	3.8	0.41	159	5.2	0.41	5.1	0.41
XI. Other malignant epithelial neoplasms and melanomas	117	5.3	0.49	5.2	0.48	396	12.6	0.63	12.6	0.63
XI(b). Thyroid carcinomas	43	1.9	0.30	1.9	0.29	176	5.6	0.42	5.6	0.42
XI(d). Malignant melanomas	14	0.6	0.17	0.6	0.17	19	0.6	0.14	0.6	0.14
XII. Other & unspecified malignant neoplasms	25	1.1	0.22	1.1	0.22	43	1.4	0.21	1.4	0.21

[1] Rates are **per million** population and are age-adjusted by five-year age groups to the 2000 U.S. standard population based on single years of age and to the 1996 Canadian population standard.
[2] See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.
[3] Abbreviations: S.E. - Standard Error; CNS - Central Nervous System.
- Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for All Sites combined.

N.B. In areas with small Latino populations, methods to indirectly identify Latinos (like NHIAv2) can overestimate the ethnicity-specific counts of cancer cases. Also, even small errors in Latino population estimates can affect the magnitude of the cancer rates. Data Item 190 used in place of NHIA for MA and RI, see technical notes.

NAACCR Canadian Combined Pediatric Cancer Incidence
2010-2014

Total Cancer Cases and Average Annual Age-specific Incidence Rates[1] for Pediatric Cancers
NAACCR Canadian Combined[2], 2010-2014, All Races, Males

	Ages 0-4			Ages 5-9			Ages 10-14			Ages 15-19		
Cancer Types	*Count*	*Rate*	*S.E.*[3]	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
All Sites	**436**	**228.1**	**10.93**	**249**	**135.6**	**8.59**	**244**	**129.3**	**8.28**	**480**	**222.5**	**10.15**
I. Leukemias	162	84.8	6.66	79	43.0	4.84	53	28.1	3.86	75	34.8	4.01
I(a). Lymphoid leukemias	130	68.0	5.97	71	38.7	4.59	37	19.6	3.22	42	19.5	3.00
I(b). Acute myeloid leukemias	22	11.5	2.45	-	-	-	10	5.3	1.68	18	8.3	1.97
II. Lymphomas & reticuloendothelial neoplasms	31	16.2	2.91	44	24.0	3.61	63	33.4	4.21	119	55.2	5.06
II(a). Hodgkin lymphomas	-	-	-	10	5.4	1.72	22	11.7	2.49	63	29.2	3.68
II(b, c, e). Non-Hodgkin lymphomas	21	11.0	2.40	26	14.2	2.78	34	18.0	3.09	53	24.6	3.37
III. CNS[3] & misc. intracranial & intraspinal neoplasms	68	35.6	4.31	62	33.8	4.29	46	24.4	3.59	41	19.0	2.97
III(a). Ependymomas and choroids plexus tumor	15	7.8	2.03	-	-	-	-	-	-	9	4.2	1.39
III(b). Astrocytomas	18	9.4	2.22	31	16.9	3.03	23	12.2	2.54	17	7.9	1.91
III(c). Intracranial and intraspinal embryonal tumors	26	13.6	2.67	17	9.3	2.25	9	4.8	1.59	9	4.2	1.39
III(d). Other gliomas	8	4.2	1.48	10	5.4	1.72	9	4.8	1.59	6	2.8	1.14
IV. Neuroblastoma and other peripheral nervous cell tumors	66	34.5	4.25	8	4.4	1.54	-	-	-	-	-	-
IV(a). Neuroblastoma & ganglioneuroblastoma	66	34.5	4.25	8	4.4	1.54	-	-	-	-	-	-
V. Retinoblastoma	21	11.0	2.40	-	-	-	-	-	-	-	-	-
VI. Renal Tumors	33	17.3	3.01	16	8.7	2.18	-	-	-	-	-	-
VI(a). Nephroblastoma and other nonepithelial renal tumors	33	17.3	3.01	15	8.2	2.11	-	-	-	-	-	-
VII. Hepatic tumors	14	7.3	1.96	-	-	-	-	-	-	-	-	-
VII(a). Hepatoblastoma	13	6.8	1.89	-	-	-	-	-	-	-	-	-
VIII. Malignant bone tumors	-	-	-	8	4.4	1.54	20	10.6	2.37	31	14.4	2.58
VIII(a). Osteosarcomas	-	-	-	6	3.3	1.33	10	5.3	1.68	19	8.8	2.02
VIII(c). Ewing tumor and related sarcomas of bone	-	-	-	-	-	-	10	5.3	1.68	10	4.6	1.47
IX. Soft tissue and other extraosseous sarcomas	24	12.6	2.56	18	9.8	2.31	20	10.6	2.37	40	18.5	2.93
IX(a). Rhabdomyosarcomas	16	8.4	2.09	12	6.5	1.89	10	5.3	1.68	10	4.6	1.47
X. Germ cell & trophoblastic tumors & neoplasms of gonads	8	4.2	1.48	6	3.3	1.33	14	7.4	1.98	98	45.4	4.59
X(a). Intracranial & intraspinal germ cell tumors	-	-	-	-	-	-	11	5.8	1.76	12	5.6	1.61
X(c). Malignant gonadal germ cell tumors	-	-	-	-	-	-	-	-	-	84	38.9	4.25
XI. Other malignant epithelial neoplasms and melanomas	-	-	-	-	-	-	24	12.7	2.60	67	31.1	3.79
XI(b). Thyroid carcinomas	-	-	-	-	-	-	-	-	-	21	9.7	2.12
XI(d). Malignant melanomas	-	-	-	-	-	-	-	-	-	10	4.6	1.47
XII. Other & unspecified malignant neoplasms	-	-	-	-	-	-	-	-	-	-	-	-

[1] Rates are **per million** population.
[2] See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.
[3] Abbreviations: S.E. - Standard Error; CNS - Central Nervous System.
- Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for All Sites combined.

Total Cancer Cases and Average Annual Age-specific Incidence Rates[1] for Pediatric Cancers
NAACCR Canadian Combined[2], 2010-2014, All Races, Females

	Ages 0-4			Ages 5-9			Ages 10-14			Ages 15-19		
Cancer Types	*Count*	*Rate*	*S.E.*[3]	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
All Sites	**427**	**235.9**	**11.42**	**205**	**118.0**	**8.24**	**208**	**117.0**	**8.12**	**346**	**172.0**	**9.25**
I. Leukemias	167	92.3	7.14	69	39.7	4.78	45	25.3	3.77	41	20.4	3.18
I(a). Lymphoid leukemias	131	72.4	6.32	53	30.5	4.19	29	16.3	3.03	15	7.5	1.93
I(b). Acute myeloid leukemias	25	13.8	2.76	10	5.8	1.82	12	6.8	1.95	21	10.4	2.28
II. Lymphomas & reticuloendothelial neoplasms	32	17.7	3.13	22	12.7	2.70	42	23.6	3.65	81	40.3	4.47
II(a). Hodgkin lymphomas	-	-	-	-	-	-	19	10.7	2.45	60	29.8	3.85
II(b, c, e). Non-Hodgkin lymphomas	14	7.7	2.07	13	7.5	2.08	19	10.7	2.45	19	9.4	2.17
III. CNS[3] & misc. intracranial & intraspinal neoplasms	67	37.0	4.52	58	33.4	4.38	34	19.1	3.28	33	16.4	2.86
III(a). Ependymomas and choroids plexus tumor	9	5.0	1.66	-	-	-	-	-	-	6	3.0	1.22
III(b). Astrocytomas	31	17.1	3.08	28	16.1	3.05	22	12.4	2.64	17	8.5	2.05
III(c). Intracranial and intraspinal embryonal tumors	17	9.4	2.28	10	5.8	1.82	6	3.4	1.38	-	-	-
III(d). Other gliomas	9	5.0	1.66	12	6.9	1.99	-	-	-	7	3.5	1.32
IV. Neuroblastoma and other peripheral nervous cell tumors	62	34.3	4.35	6	3.5	1.41	-	-	-	-	-	-
IV(a). Neuroblastoma & ganglioneuroblastoma	62	34.3	4.35	6	3.5	1.41	-	-	-	-	-	-
V. Retinoblastoma	13	7.2	1.99	-	-	-	-	-	-	-	-	-
VI. Renal Tumors	37	20.4	3.36	11	6.3	1.91	-	-	-	-	-	-
VI(a). Nephroblastoma and other nonepithelial renal tumors	37	20.4	3.36	10	5.8	1.82	-	-	-	-	-	-
VII. Hepatic tumors	9	5.0	1.66	-	-	-	-	-	-	-	-	-
VII(a). Hepatoblastoma	9	5.0	1.66	-	-	-	-	-	-	-	-	-
VIII. Malignant bone tumors	-	-	-	14	8.1	2.15	25	14.1	2.81	30	14.9	2.72
VIII(a). Osteosarcomas	-	-	-	6	3.5	1.41	16	9.0	2.25	18	8.9	2.11
VIII(c). Ewing tumor and related sarcomas of bone	-	-	-	7	4.0	1.52	9	5.1	1.69	8	4.0	1.41
IX. Soft tissue and other extraosseous sarcomas	22	12.2	2.59	10	5.8	1.82	20	11.3	2.52	23	11.4	2.38
IX(a). Rhabdomyosarcomas	10	5.5	1.75	7	4.0	1.52	-	-	-	-	-	-
X. Germ cell & trophoblastic tumors & neoplasms of gonads	12	6.6	1.91	-	-	-	12	6.8	1.95	20	9.9	2.22
X(a). Intracranial & intraspinal germ cell tumors	-	-	-	-	-	-	-	-	-	-	-	-
X(c). Malignant gonadal germ cell tumors	-	-	-	-	-	-	11	6.2	1.87	13	6.5	1.79
XI. Other malignant epithelial neoplasms and melanomas	-	-	-	12	6.9	1.99	27	15.2	2.92	112	55.7	5.26
XI(b). Thyroid carcinomas	-	-	-	-	-	-	10	5.6	1.78	58	28.8	3.79
XI(d). Malignant melanomas	-	-	-	7	4.0	1.52	-	-	-	16	8.0	1.99
XII. Other & unspecified malignant neoplasms	-	-	-	-	-	-	-	-	-	-	-	-

[1] Rates are **per million** population.
[2] See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.
[3] Abbreviations: S.E. - Standard Error; CNS - Central Nervous System.
- Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for All Sites combined.

Total Cancer Cases and Average Annual Age-adjusted Incidence Rates[1] for Pediatric Cancers
NAACCR Canadian Combined[2], 2010-2014, All Races, Males

	Ages 0-14					Ages 0-19				
		2000 U.S.		1996 Cdn			2000 U.S.		1996 Cdn	
Cancer Types	*Count*	*Rate*	*S.E.*[3]	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Rate*	*S.E.*
All Sites	**929**	**163.3**	**5.36**	**163.8**	**5.38**	**1,409**	**178.1**	**4.76**	**178.6**	**4.77**
I. Leukemias	294	51.4	3.00	52.0	3.03	369	47.2	2.46	47.7	2.48
I(a). Lymphoid leukemias	238	41.7	2.70	42.2	2.74	280	36.1	2.16	36.5	2.18
I(b). Acute myeloid leukemias	35	6.1	1.03	6.1	1.03	53	6.6	0.91	6.7	0.92
II. Lymphomas & reticuloendothelial neoplasms	138	24.7	2.10	24.6	2.10	257	32.3	2.02	32.3	2.02
II(a). Hodgkin lymphomas	33	6.0	1.04	5.9	1.03	96	11.8	1.21	11.8	1.20
II(b, c, e). Non-Hodgkin lymphomas	81	14.5	1.61	14.5	1.61	134	17.0	1.47	17.0	1.47
III. CNS[3] & misc. intracranial & intraspinal neoplasms	176	31.2	2.35	31.3	2.36	217	28.1	1.91	28.2	1.92
III(a). Ependymomas and choroids plexus tumor	18	3.1	0.72	3.1	0.73	27	3.3	0.65	3.4	0.65
III(b). Astrocytomas	72	12.9	1.52	12.9	1.52	89	11.6	1.23	11.6	1.23
III(c). Intracranial and intraspinal embryonal tumors	52	9.1	1.27	9.2	1.28	61	7.9	1.01	8.0	1.02
III(d). Other gliomas	27	4.8	0.93	4.8	0.93	33	4.3	0.75	4.3	0.75
IV. Neuroblastoma and other peripheral nervous cell tumors	76	12.9	1.48	12.9	1.48	77	9.8	1.12	9.8	1.11
IV(a). Neuroblastoma & ganglioneuroblastoma	75	12.8	1.47	12.7	1.47	75	9.5	1.10	9.5	1.10
V. Retinoblastoma	22	3.7	0.79	3.7	0.78	22	2.8	0.59	2.7	0.58
VI. Renal Tumors	51	8.9	1.24	9.0	1.26	53	6.9	0.94	7.0	0.96
VI(a). Nephroblastoma and other nonepithelial renal tumors	50	8.7	1.23	8.8	1.25	51	6.6	0.93	6.7	0.94
VII. Hepatic tumors	16	2.7	0.68	2.8	0.69	20	2.5	0.56	2.5	0.57
VII(a). Hepatoblastoma	14	2.4	0.63	2.4	0.64	14	1.8	0.48	1.8	0.48
VIII. Malignant bone tumors	31	5.6	1.00	5.5	1.00	62	7.8	0.99	7.8	0.99
VIII(a). Osteosarcomas	16	2.9	0.73	2.9	0.72	35	4.4	0.74	4.4	0.74
VIII(c). Ewing tumor and related sarcomas of bone	15	2.7	0.69	2.7	0.69	25	3.2	0.64	3.2	0.63
IX. Soft tissue and other extraosseous sarcomas	62	11.0	1.39	10.9	1.39	102	12.9	1.28	12.9	1.28
IX(a). Rhabdomyosarcomas	38	6.7	1.09	6.7	1.09	48	6.2	0.89	6.2	0.90
X. Germ cell & trophoblastic tumors & neoplasms of gonads	28	5.0	0.94	5.0	0.94	126	15.2	1.35	15.1	1.35
X(a). Intracranial & intraspinal germ cell tumors	15	2.7	0.70	2.7	0.69	27	3.4	0.66	3.4	0.66
X(c). Malignant gonadal germ cell tumors	8	1.4	0.50	1.4	0.50	92	10.8	1.13	10.8	1.13
XI. Other malignant epithelial neoplasms and melanomas	30	5.4	0.98	5.3	0.97	97	11.8	1.20	11.8	1.20
XI(b). Thyroid carcinomas	6	1.1	0.44	1.1	0.44	27	3.3	0.63	3.3	0.63
XI(d). Malignant melanomas	9	1.6	0.53	1.6	0.52	19	2.4	0.54	2.3	0.54
XII. Other & unspecified malignant neoplasms	-	-	-	-	-	7	0.9	0.34	0.9	0.34

[1] Rates are **per million** population and are age-adjusted by five-year age groups to the 2000 U.S. standard population based on single years of age and to the 1996 Canadian population standard.
[2] See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.
[3] Abbreviations: S.E. - Standard Error; CNS - Central Nervous System.
- Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for All Sites combined.

Total Cancer Cases and Average Annual Age-adjusted Incidence Rates[1] for Pediatric Cancers
NAACCR Canadian Combined[2], 2010-2014, All Races, Females

	Ages 0-14					Ages 0-19				
		2000 U.S.		1996 Cdn			2000 U.S.		1996 Cdn	
Cancer Types	*Count*	*Rate*	*S.E.[3]*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Rate*	*S.E.*
All Sites	**840**	**155.6**	**5.37**	**156.1**	**5.39**	**1,186**	**159.7**	**4.65**	**160.1**	**4.66**
I. Leukemias	281	51.8	3.09	52.3	3.12	322	43.9	2.45	44.3	2.47
I(a). Lymphoid leukemias	213	39.2	2.69	39.8	2.73	228	31.2	2.07	31.6	2.10
I(b). Acute myeloid leukemias	47	8.7	1.27	8.7	1.27	68	9.1	1.11	9.1	1.11
II. Lymphomas & reticuloendothelial neoplasms	96	18.0	1.84	18.0	1.84	177	23.6	1.78	23.6	1.78
II(a). Hodgkin lymphomas	22	4.2	0.90	4.2	0.89	82	10.7	1.18	10.6	1.17
II(b, c, e). Non-Hodgkin lymphomas	46	8.7	1.28	8.7	1.28	65	8.9	1.10	8.9	1.10
III. CNS[3] & misc. intracranial & intraspinal neoplasms	159	29.7	2.36	29.9	2.37	192	26.4	1.91	26.5	1.91
III(a). Ependymomas and choroids plexus tumor	16	3.0	0.74	3.0	0.74	22	3.0	0.63	3.0	0.64
III(b). Astrocytomas	81	15.2	1.69	15.2	1.69	98	13.5	1.36	13.5	1.37
III(c). Intracranial and intraspinal embryonal tumors	33	6.1	1.07	6.1	1.07	36	5.0	0.83	5.0	0.83
III(d). Other gliomas	25	4.7	0.94	4.7	0.95	32	4.4	0.78	4.4	0.78
IV. Neuroblastoma and other peripheral nervous cell tumors	69	12.4	1.49	12.3	1.48	73	9.8	1.14	9.7	1.14
IV(a). Neuroblastoma & ganglioneuroblastoma	68	12.2	1.48	12.1	1.47	70	9.4	1.12	9.3	1.11
V. Retinoblastoma	13	2.3	0.64	2.3	0.65	13	1.7	0.48	1.7	0.48
VI. Renal Tumors	48	8.7	1.26	8.8	1.27	49	6.7	0.95	6.7	0.96
VI(a). Nephroblastoma and other nonepithelial renal tumors	47	8.5	1.24	8.6	1.26	47	6.4	0.93	6.4	0.94
VII. Hepatic tumors	11	2.0	0.60	2.0	0.60	11	1.5	0.45	1.5	0.45
VII(a). Hepatoblastoma	10	1.8	0.57	1.8	0.56	10	1.3	0.42	1.3	0.42
VIII. Malignant bone tumors	41	7.9	1.23	7.8	1.22	71	9.6	1.15	9.6	1.14
VIII(a). Osteosarcomas	22	4.2	0.90	4.2	0.89	40	5.4	0.86	5.4	0.85
VIII(c). Ewing tumor and related sarcomas of bone	18	3.4	0.81	3.4	0.81	26	3.6	0.70	3.6	0.70
IX. Soft tissue and other extraosseous sarcomas	52	9.7	1.34	9.7	1.34	75	10.1	1.17	10.1	1.17
IX(a). Rhabdomyosarcomas	21	3.9	0.85	4.0	0.86	25	3.4	0.69	3.5	0.69
X. Germ cell & trophoblastic tumors & neoplasms of gonads	27	5.0	0.97	5.0	0.96	47	6.3	0.91	6.2	0.91
X(a). Intracranial & intraspinal germ cell tumors	-	-	-	-	-	-	-	-	-	-
X(c). Malignant gonadal germ cell tumors	16	3.0	0.76	3.0	0.76	29	3.9	0.73	3.9	0.72
XI. Other malignant epithelial neoplasms and melanomas	42	8.0	1.24	7.9	1.23	154	20.0	1.62	19.9	1.61
XI(b). Thyroid carcinomas	14	2.7	0.72	2.7	0.71	72	9.3	1.09	9.2	1.09
XI(d). Malignant melanomas	12	2.3	0.66	2.3	0.66	28	3.7	0.70	3.7	0.70
XII. Other & unspecified malignant neoplasms	-	-	-	-	-	-	-	-	-	-

[1] Rates are **per million** population and are age-adjusted by five-year age groups to the 2000 U.S. standard population based on single years of age and to the 1996 Canadian population standard.
[2] See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.
[3] Abbreviations: S.E. - Standard Error; CNS - Central Nervous System.
- Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for All Sites combined.

NAACCR North American Combined Pediatric Cancer Incidence 2010-2014

Total Cancer Cases and Average Annual Age-specific Incidence Rates[1] for Pediatric Cancers
NAACCR North American Combined[2], 2010-2014, All Races, Males

	Ages 0-4			Ages 5-9			Ages 10-14			Ages 15-19		
Cancer Types	*Count*	*Rate*	*S.E.*[3]	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
All Sites	**12,681**	**247.2**	**2.20**	**7,344**	**140.5**	**1.64**	**7,758**	**146.4**	**1.66**	**12,697**	**229.1**	**2.03**
I. Leukemias	4,385	85.5	1.29	2,472	47.3	0.95	1,901	35.9	0.82	2,197	39.6	0.85
I(a). Lymphoid leukemias	3,389	66.1	1.13	2,041	39.0	0.86	1,274	24.0	0.67	1,204	21.7	0.63
I(b). Acute myeloid leukemias	607	11.8	0.48	238	4.6	0.30	366	6.9	0.36	555	10.0	0.43
II. Lymphomas & reticuloendothelial neoplasms	1,054	20.5	0.63	1,305	25.0	0.69	1,717	32.4	0.78	3,117	56.2	1.01
II(a). Hodgkin lymphomas	48	0.9	0.14	299	5.7	0.33	677	12.8	0.49	1,703	30.7	0.74
II(b, c, e). Non-Hodgkin lymphomas	451	8.8	0.41	754	14.4	0.53	858	16.2	0.55	1,306	23.6	0.65
III. CNS[3] & misc. intracranial & intraspinal neoplasms	2,273	44.3	0.93	1,923	36.8	0.84	1,567	29.6	0.75	1,342	24.2	0.66
III(a). Ependymomas and choroids plexus tumor	325	6.3	0.35	111	2.1	0.20	105	2.0	0.19	95	1.7	0.18
III(b). Astrocytomas	997	19.4	0.62	959	18.3	0.59	846	16.0	0.55	776	14.0	0.50
III(c). Intracranial and intraspinal embryonal tumors	612	11.9	0.48	448	8.6	0.40	268	5.1	0.31	164	3.0	0.23
III(d). Other gliomas	262	5.1	0.32	351	6.7	0.36	291	5.5	0.32	246	4.4	0.28
IV. Neuroblastoma and other peripheral nervous cell tumors	1,557	30.4	0.77	225	4.3	0.29	83	1.6	0.17	52	0.9	0.13
IV(a). Neuroblastoma & ganglioneuroblastoma	1,548	30.2	0.77	218	4.2	0.28	60	1.1	0.15	24	0.4	0.09
V. Retinoblastoma	624	12.2	0.49	28	0.5	0.10	6	0.1	0.05	-	-	-
VI. Renal Tumors	941	18.3	0.60	264	5.0	0.31	68	1.3	0.16	99	1.8	0.18
VI(a). Nephroblastoma and other nonepithelial renal tumors	930	18.1	0.59	242	4.6	0.30	33	0.6	0.11	20	0.4	0.08
VII. Hepatic tumors	495	9.6	0.43	41	0.8	0.12	52	1.0	0.14	57	1.0	0.14
VII(a). Hepatoblastoma	473	9.2	0.42	25	0.5	0.10	13	0.2	0.07	-	-	-
VIII. Malignant bone tumors	81	1.6	0.18	308	5.9	0.34	779	14.7	0.53	913	16.5	0.55
VIII(a). Osteosarcomas	16	0.3	0.08	150	2.9	0.23	465	8.8	0.41	530	9.6	0.42
VIII(c). Ewing tumor and related sarcomas of bone	54	1.1	0.14	136	2.6	0.22	247	4.7	0.30	277	5.0	0.30
IX. Soft tissue and other extraosseous sarcomas	745	14.5	0.53	486	9.3	0.42	625	11.8	0.47	898	16.2	0.54
IX(a). Rhabdomyosarcomas	438	8.5	0.41	268	5.1	0.31	204	3.9	0.27	207	3.7	0.26
X. Germ cell & trophoblastic tumors & neoplasms of gonads	325	6.3	0.35	85	1.6	0.18	311	5.9	0.33	2,290	41.3	0.86
X(a). Intracranial & intraspinal germ cell tumors	32	0.6	0.11	71	1.4	0.16	218	4.1	0.28	225	4.1	0.27
X(c). Malignant gonadal germ cell tumors	167	3.3	0.25	-	-	-	68	1.3	0.16	1,918	34.6	0.79
XI. Other malignant epithelial neoplasms and melanomas	130	2.5	0.22	195	3.7	0.27	628	11.9	0.47	1,703	30.7	0.74
XI(b). Thyroid carcinomas	9	0.2	0.06	48	0.9	0.13	208	3.9	0.27	511	9.2	0.41
XI(d). Malignant melanomas	72	1.4	0.17	78	1.5	0.17	135	2.5	0.22	448	8.1	0.38
XII. Other & unspecified malignant neoplasms	71	1.4	0.16	12	0.2	0.07	21	0.4	0.09	29	0.5	0.10

[1] Rates are **per million** population.
[2] See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.
[3] Abbreviations: S.E. - Standard Error; CNS - Central Nervous System.
- Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for All Sites combined.

Total Cancer Cases and Average Annual Age-specific Incidence Rates[1] for Pediatric Cancers
NAACCR North American Combined[2], 2010-2014, All Races, Females

	Ages 0-4			Ages 5-9			Ages 10-14			Ages 15-19		
Cancer Types	*Count*	*Rate*	*S.E.*[3]	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
All Sites	**11,101**	**226.2**	**2.15**	**5,987**	**119.5**	**1.54**	**7,037**	**138.8**	**1.65**	**11,284**	**214.4**	**2.02**
I. Leukemias	3,907	79.6	1.27	1,943	38.8	0.88	1,463	28.9	0.75	1,435	27.3	0.72
I(a). Lymphoid leukemias	3,008	61.3	1.12	1,563	31.2	0.79	908	17.9	0.59	556	10.6	0.45
I(b). Acute myeloid leukemias	570	11.6	0.49	227	4.5	0.30	337	6.6	0.36	532	10.1	0.44
II. Lymphomas & reticuloendothelial neoplasms	654	13.3	0.52	620	12.4	0.50	1,112	21.9	0.66	2,391	45.4	0.93
II(a). Hodgkin lymphomas	21	0.4	0.09	90	1.8	0.19	516	10.2	0.45	1,676	31.8	0.78
II(b, c, e). Non-Hodgkin lymphomas	229	4.7	0.31	336	6.7	0.37	467	9.2	0.43	650	12.4	0.48
III. CNS[3] & misc. intracranial & intraspinal neoplasms	2,033	41.4	0.92	1,646	32.9	0.81	1,353	26.7	0.73	1,100	20.9	0.63
III(a). Ependymomas and choroids plexus tumor	239	4.9	0.31	118	2.4	0.22	70	1.4	0.17	78	1.5	0.17
III(b). Astrocytomas	1,024	20.9	0.65	876	17.5	0.59	799	15.8	0.56	645	12.3	0.48
III(c). Intracranial and intraspinal embryonal tumors	461	9.4	0.44	242	4.8	0.31	161	3.2	0.25	95	1.8	0.19
III(d). Other gliomas	250	5.1	0.32	356	7.1	0.38	265	5.2	0.32	239	4.5	0.29
IV. Neuroblastoma and other peripheral nervous cell tumors	1,416	28.8	0.77	201	4.0	0.28	72	1.4	0.17	37	0.7	0.12
IV(a). Neuroblastoma & ganglioneuroblastoma	1,411	28.7	0.77	193	3.9	0.28	57	1.1	0.15	21	0.4	0.09
V. Retinoblastoma	655	13.3	0.52	25	0.5	0.10	-	-	-	-	-	-
VI. Renal Tumors	1,012	20.6	0.65	345	6.9	0.37	81	1.6	0.18	108	2.1	0.20
VI(a). Nephroblastoma and other nonepithelial renal tumors	1,003	20.4	0.65	330	6.6	0.36	46	0.9	0.13	22	0.4	0.09
VII. Hepatic tumors	266	5.4	0.33	42	0.8	0.13	45	0.9	0.13	68	1.3	0.16
VII(a). Hepatoblastoma	257	5.2	0.33	22	0.4	0.09	12	0.2	0.07	-	-	-
VIII. Malignant bone tumors	92	1.9	0.20	296	5.9	0.34	680	13.4	0.51	581	11.0	0.46
VIII(a). Osteosarcomas	13	0.3	0.07	163	3.3	0.25	427	8.4	0.41	327	6.2	0.34
VIII(c). Ewing tumor and related sarcomas of bone	62	1.3	0.16	103	2.1	0.20	202	4.0	0.28	158	3.0	0.24
IX. Soft tissue and other extraosseous sarcomas	557	11.3	0.48	390	7.8	0.39	616	12.2	0.49	711	13.5	0.51
IX(a). Rhabdomyosarcomas	325	6.6	0.37	196	3.9	0.28	182	3.6	0.27	160	3.0	0.24
X. Germ cell & trophoblastic tumors & neoplasms of gonads	318	6.5	0.36	192	3.8	0.28	466	9.2	0.43	761	14.5	0.52
X(a). Intracranial & intraspinal germ cell tumors	29	0.6	0.11	56	1.1	0.15	75	1.5	0.17	28	0.5	0.10
X(c). Malignant gonadal germ cell tumors	31	0.6	0.11	112	2.2	0.21	321	6.3	0.35	448	8.5	0.40
XI. Other malignant epithelial neoplasms and melanomas	139	2.8	0.24	276	5.5	0.33	1,112	21.9	0.66	4,010	76.2	1.20
XI(b). Thyroid carcinomas	14	0.3	0.08	100	2.0	0.20	635	12.5	0.50	2,449	46.5	0.94
XI(d). Malignant melanomas	90	1.8	0.19	104	2.1	0.20	150	3.0	0.24	669	12.7	0.49
XII. Other & unspecified malignant neoplasms	52	1.1	0.15	11	0.2	0.07	36	0.7	0.12	81	1.5	0.17

[1] Rates are **per million** population.
[2] See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.
[3] Abbreviations: S.E. - Standard Error; CNS - Central Nervous System.
- Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for All Sites combined.

Total Cancer Cases and Average Annual Age-adjusted Incidence Rates[1] for Pediatric Cancers
NAACCR North American Combined[2], 2010-2014, All Races, Males

	Ages 0-14					Ages 0-19				
		2000 U.S.		1996 Cdn			2000 U.S.		1996 Cdn	
Cancer Types	*Count*	*Rate*	*S.E.*[3]	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Rate*	*S.E.*
All Sites	**27,783**	**176.9**	**1.06**	**177.4**	**1.06**	**40,480**	**190.0**	**0.94**	**190.4**	**0.95**
I. Leukemias	8,758	55.7	0.59	56.2	0.60	10,955	51.6	0.49	52.0	0.50
I(a). Lymphoid leukemias	6,704	42.6	0.52	43.1	0.53	7,908	37.3	0.42	37.7	0.42
I(b). Acute myeloid leukemias	1,211	7.7	0.22	7.7	0.22	1,766	8.3	0.20	8.3	0.20
II. Lymphomas & reticuloendothelial neoplasms	4,076	26.1	0.41	26.0	0.41	7,193	33.7	0.40	33.6	0.40
II(a). Hodgkin lymphomas	1,024	6.6	0.21	6.5	0.20	2,727	12.7	0.24	12.6	0.24
II(b, c, e). Non-Hodgkin lymphomas	2,063	13.2	0.29	13.2	0.29	3,369	15.8	0.27	15.8	0.27
III. CNS[3] & misc. intracranial & intraspinal neoplasms	5,763	36.7	0.48	36.9	0.49	7,105	33.6	0.40	33.7	0.40
III(a). Ependymomas and choroids plexus tumor	541	3.4	0.15	3.5	0.15	636	3.0	0.12	3.0	0.12
III(b). Astrocytomas	2,802	17.9	0.34	17.9	0.34	3,578	16.9	0.28	16.9	0.28
III(c). Intracranial and intraspinal embryonal tumors	1,328	8.5	0.23	8.5	0.23	1,492	7.1	0.18	7.1	0.18
III(d). Other gliomas	904	5.8	0.19	5.8	0.19	1,150	5.4	0.16	5.4	0.16
IV. Neuroblastoma and other peripheral nervous cell tumors	1,865	11.8	0.27	11.8	0.27	1,917	9.1	0.21	9.1	0.21
IV(a). Neuroblastoma & ganglioneuroblastoma	1,826	11.5	0.27	11.5	0.27	1,850	8.7	0.20	8.8	0.20
V. Retinoblastoma	658	4.2	0.16	4.1	0.16	658	3.1	0.12	3.1	0.12
VI. Renal Tumors	1,273	8.0	0.23	8.2	0.23	1,372	6.5	0.17	6.6	0.18
VI(a). Nephroblastoma and other nonepithelial renal tumors	1,205	7.6	0.22	7.7	0.22	1,225	5.8	0.17	5.9	0.17
VII. Hepatic tumors	588	3.7	0.15	3.7	0.15	645	3.0	0.12	3.0	0.12
VII(a). Hepatoblastoma	511	3.2	0.14	3.2	0.14	513	2.4	0.11	2.4	0.11
VIII. Malignant bone tumors	1,168	7.5	0.22	7.4	0.22	2,081	9.8	0.21	9.7	0.21
VIII(a). Osteosarcomas	631	4.1	0.16	4.0	0.16	1,161	5.4	0.16	5.4	0.16
VIII(c). Ewing tumor and related sarcomas of bone	437	2.8	0.13	2.8	0.13	714	3.4	0.13	3.3	0.13
IX. Soft tissue and other extraosseous sarcomas	1,856	11.8	0.27	11.8	0.27	2,754	12.9	0.25	12.9	0.25
IX(a). Rhabdomyosarcomas	910	5.8	0.19	5.8	0.19	1,117	5.3	0.16	5.3	0.16
X. Germ cell & trophoblastic tumors & neoplasms of gonads	721	4.6	0.17	4.5	0.17	3,011	13.8	0.25	13.8	0.25
X(a). Intracranial & intraspinal germ cell tumors	321	2.1	0.12	2.0	0.11	546	2.6	0.11	2.5	0.11
X(c). Malignant gonadal germ cell tumors	238	1.5	0.10	1.5	0.10	2,156	9.8	0.21	9.8	0.21
XI. Other malignant epithelial neoplasms and melanomas	953	6.1	0.20	6.0	0.20	2,656	12.3	0.24	12.3	0.24
XI(b). Thyroid carcinomas	265	1.7	0.10	1.7	0.10	776	3.6	0.13	3.6	0.13
XI(d). Malignant melanomas	285	1.8	0.11	1.8	0.11	733	3.4	0.13	3.4	0.13
XII. Other & unspecified malignant neoplasms	104	0.7	0.06	0.7	0.06	133	0.6	0.05	0.6	0.05

[1] Rates are **per million** population and are age-adjusted by five-year age groups to the 2000 U.S. standard population based on single years of age and to the 1996 Canadian population standard.
[2] See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.
[3] Abbreviations: S.E. - Standard Error; CNS - Central Nervous System.
- Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for All Sites combined.

Total Cancer Cases and Average Annual Age-adjusted Incidence Rates[1] for Pediatric Cancers
NAACCR North American Combined[2], 2010-2014, All Races, Females

	Ages 0-14					Ages 0-19				
		2000 U.S.		1996 Cdn			2000 U.S.		1996 Cdn	
Cancer Types	*Count*	*Rate*	*S.E.[3]*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Rate*	*S.E.*
All Sites	**24,125**	**160.5**	**1.03**	**160.8**	**1.04**	**35,409**	**174.0**	**0.93**	**174.3**	**0.93**
I. Leukemias	7,313	48.5	0.57	49.0	0.57	8,748	43.2	0.46	43.5	0.47
I(a). Lymphoid leukemias	5,479	36.3	0.49	36.8	0.50	6,035	29.8	0.38	30.2	0.39
I(b). Acute myeloid leukemias	1,134	7.5	0.22	7.5	0.22	1,666	8.2	0.20	8.2	0.20
II. Lymphomas & reticuloendothelial neoplasms	2,386	15.9	0.33	15.9	0.32	4,777	23.4	0.34	23.3	0.34
II(a). Hodgkin lymphomas	627	4.2	0.17	4.2	0.17	2,303	11.2	0.23	11.1	0.23
II(b, c, e). Non-Hodgkin lymphomas	1,032	6.9	0.21	6.9	0.21	1,682	8.3	0.20	8.3	0.20
III. CNS[3] & misc. intracranial & intraspinal neoplasms	5,032	33.5	0.47	33.6	0.47	6,132	30.3	0.39	30.4	0.39
III(a). Ependymomas and choroids plexus tumor	427	2.8	0.14	2.9	0.14	505	2.5	0.11	2.5	0.11
III(b). Astrocytomas	2,699	18.0	0.35	18.1	0.35	3,344	16.5	0.29	16.6	0.29
III(c). Intracranial and intraspinal embryonal tumors	864	5.7	0.20	5.8	0.20	959	4.7	0.15	4.8	0.15
III(d). Other gliomas	871	5.8	0.20	5.8	0.20	1,110	5.5	0.17	5.5	0.17
IV. Neuroblastoma and other peripheral nervous cell tumors	1,689	11.2	0.27	11.2	0.27	1,726	8.5	0.21	8.5	0.21
IV(a). Neuroblastoma & ganglioneuroblastoma	1,661	11.0	0.27	11.0	0.27	1,682	8.3	0.20	8.3	0.20
V. Retinoblastoma	681	4.5	0.17	4.5	0.17	682	3.4	0.13	3.4	0.13
VI. Renal Tumors	1,438	9.5	0.25	9.6	0.25	1,546	7.6	0.19	7.7	0.20
VI(a). Nephroblastoma and other nonepithelial renal tumors	1,379	9.1	0.25	9.2	0.25	1,401	6.9	0.19	7.0	0.19
VII. Hepatic tumors	353	2.3	0.12	2.3	0.12	421	2.1	0.10	2.1	0.10
VII(a). Hepatoblastoma	291	1.9	0.11	1.9	0.11	294	1.5	0.08	1.5	0.08
VIII. Malignant bone tumors	1,068	7.2	0.22	7.1	0.22	1,649	8.1	0.20	8.1	0.20
VIII(a). Osteosarcomas	603	4.0	0.16	4.0	0.16	930	4.6	0.15	4.6	0.15
VIII(c). Ewing tumor and related sarcomas of bone	367	2.5	0.13	2.4	0.13	525	2.6	0.11	2.6	0.11
IX. Soft tissue and other extraosseous sarcomas	1,563	10.4	0.26	10.4	0.26	2,274	11.2	0.23	11.2	0.23
IX(a). Rhabdomyosarcomas	703	4.7	0.18	4.7	0.18	863	4.3	0.15	4.3	0.15
X. Germ cell & trophoblastic tumors & neoplasms of gonads	976	6.5	0.21	6.4	0.21	1,737	8.5	0.20	8.5	0.20
X(a). Intracranial & intraspinal germ cell tumors	160	1.1	0.08	1.1	0.08	188	0.9	0.07	0.9	0.07
X(c). Malignant gonadal germ cell tumors	464	3.1	0.14	3.1	0.14	912	4.5	0.15	4.4	0.15
XI. Other malignant epithelial neoplasms and melanomas	1,527	10.2	0.26	10.1	0.26	5,537	26.8	0.36	26.7	0.36
XI(b). Thyroid carcinomas	749	5.0	0.18	5.0	0.18	3,198	15.5	0.27	15.4	0.27
XI(d). Malignant melanomas	344	2.3	0.12	2.3	0.12	1,013	4.9	0.15	4.9	0.15
XII. Other & unspecified malignant neoplasms	99	0.7	0.07	0.7	0.07	180	0.9	0.07	0.9	0.07

[1] Rates are **per million** population and are age-adjusted by five-year age groups to the 2000 U.S. standard population based on single years of age and to the 1996 Canadian population standard.
[2] See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.
[3] Abbreviations: S.E. - Standard Error; CNS - Central Nervous System.
- Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for All Sites combined.

Section IV

NAACCR Combined Stage at Diagnosis by Age for Select Cancer Sites in the United States, Canada and North America

Stage at Diagnosis by Age and Race and Hispanic Ethnicity for Select Cancer Sites
NAACCR United States Combined, 2010-2014

Cervix, 2010-2014

	All Races			White			Black			AI/AN CHSDA[1]			API[1]			Hispanic, All Races		
Stage at Diagnosis	Count	%	Rate[2]	Count	%	Rate[2]	Count	%	Rate[2]	Count	%	Rate[2]	Count	%	Rate[2]	Count	%	Rate[2]
All Ages																		
Localized	25,568	42.4	3.4	20,128	43.7	3.4	3,500	35.7	3.3	152	40.6	3.6	1,215	42.8	2.5	4,347	43.4	4.0
Regional	21,449	35.6	2.6	16,076	34.9	2.5	3,846	39.3	3.6	138	36.9	3.4	1,095	38.6	2.3	3,649	36.5	3.6
Distant	8,891	14.7	1.0	6,692	14.5	1.0	1,682	17.2	1.6	52	13.9	1.2	383	13.5	0.8	1,245	12.4	1.3
Unknown	4,426	7.3	0.5	3,158	6.9	0.5	764	7.8	0.7	32	8.6	0.8	146	5.1	0.3	768	7.7	0.8

Colon and Rectum, 2010-2014

	All Races			White			Black			AI/AN CHSDA[1]			API[1]			Hispanic, All Races		
Stage at Diagnosis	Count	%	Rate[2]	Count	%	Rate[2]	Count	%	Rate[2]	Count	%	Rate[2]	Count	%	Rate[2]	Count	%	Rate[2]
Male																		
All Ages																		
In Situ	18,902	5.1	2.4	14,972	4.9	2.2	2,626	5.9	3.4	43	2.8	1.3	643	4.8	1.8	1,347	4.5	2.0
Localized	132,276	35.8	17.3	109,883	36.1	17.0	14,895	33.3	19.4	501	32.2	16.5	4,908	36.5	14.1	10,170	33.9	15.1
Regional	119,205	32.2	15.6	99,993	32.8	15.5	13,188	29.5	17.2	513	33.0	16.5	4,499	33.5	13.0	9,922	33.1	14.3
Distant	73,749	19.9	9.5	59,740	19.6	9.1	10,631	23.8	13.6	331	21.3	10.8	2,514	18.7	7.1	6,268	20.9	8.8
Unknown	25,602	6.9	3.5	20,130	6.6	3.2	3,394	7.6	5.1	168	10.8	6.3	882	6.6	2.8	2,252	7.5	3.7
Age < 50																		
In Situ	1,268	3.3	0.3	953	3.2	0.2	199	3.6	0.3	-	-	-	38	2.2	0.1	121	2.5	0.1
Localized	11,193	29.1	2.3	8,740	29.1	2.3	1,530	27.5	2.4	72	29.1	2.4	549	31.1	1.8	1,379	28.0	1.6
Regional	14,292	37.2	2.9	11,422	38.1	3.0	1,914	34.4	3.0	89	36.0	3.0	671	38.0	2.2	1,872	38.0	2.2
Distant	9,842	25.6	2.0	7,610	25.4	2.0	1,647	29.6	2.6	61	24.7	2.1	429	24.3	1.4	1,294	26.2	1.5
Unknown	1,816	4.7	0.4	1,281	4.3	0.3	270	4.9	0.4	24	9.7	0.8	79	4.5	0.3	266	5.4	0.3
Age ≥ 50																		
In Situ	17,634	5.3	8.0	14,019	5.1	7.5	2,427	6.2	11.4	-	-	-	605	5.2	6.2	1,226	4.9	6.9
Localized	121,083	36.5	56.6	101,143	36.8	55.5	13,365	34.1	64.1	429	32.8	53.5	4,359	37.3	46.3	8,791	35.1	50.6
Regional	104,913	31.7	48.7	88,571	32.2	48.2	11,274	28.8	54.3	424	32.4	51.7	3,828	32.8	41.2	8,050	32.2	45.9
Distant	63,907	19.3	29.1	52,130	19.0	27.9	8,984	22.9	42.5	270	20.6	33.6	2,085	17.9	22.1	4,974	19.9	27.8
Unknown	23,786	7.2	11.7	18,849	6.9	10.8	3,124	8.0	17.2	144	11.0	20.6	803	6.9	9.4	1,986	7.9	12.4
Female																		
All Ages																		
In Situ	14,906	4.4	1.6	11,446	4.2	1.5	2,498	5.6	2.4	27	1.8	0.7	486	4.0	1.1	1,080	4.3	1.3
Localized	122,142	36.1	13.2	99,993	36.3	12.9	15,596	34.8	15.0	479	32.1	13.3	4,386	36.3	10.2	8,773	34.7	10.6
Regional	110,288	32.6	11.9	91,368	33.2	11.8	13,415	29.9	13.0	536	36.0	15.4	4,136	34.2	9.6	8,444	33.4	10.2
Distant	65,552	19.4	7.1	52,588	19.1	6.9	9,970	22.3	9.6	322	21.6	8.7	2,262	18.7	5.2	5,002	19.8	5.9
Unknown	25,290	7.5	2.6	20,093	7.3	2.4	3,320	7.4	3.3	126	8.5	3.9	811	6.7	2.0	1,986	7.9	2.6
Age < 50																		
In Situ	1,289	3.7	0.3	970	3.7	0.3	212	3.6	0.3	-	-	-	52	2.9	0.2	147	3.4	0.2
Localized	11,147	32.0	2.2	8,414	32.0	2.2	1,854	31.8	2.6	67	27.7	2.3	529	29.6	1.6	1,336	30.6	1.6
Regional	12,018	34.5	2.4	9,167	34.9	2.4	1,939	33.2	2.7	84	34.7	2.9	668	37.3	2.0	1,527	34.9	1.9
Distant	8,747	25.1	1.8	6,623	25.2	1.7	1,518	26.0	2.1	62	25.6	2.1	463	25.9	1.4	1,107	25.3	1.3
Unknown	1,665	4.8	0.3	1,096	4.2	0.3	311	5.3	0.4	28	11.6	1.0	77	4.3	0.2	253	5.8	0.3
Age ≥ 50																		
In Situ	13,617	4.5	5.1	10,476	4.2	4.7	2,286	5.9	7.8	-	-	-	434	4.2	3.7	933	4.5	4.2
Localized	110,995	36.6	41.9	91,579	36.7	40.9	13,742	35.3	47.7	412	33.0	42.0	3,857	37.5	32.7	7,437	35.6	34.2
Regional	98,270	32.4	36.8	82,201	33.0	36.4	11,476	29.5	40.0	452	36.2	48.3	3,468	33.7	29.6	6,917	33.1	32.1
Distant	56,805	18.7	21.1	45,965	18.4	20.3	8,452	21.7	29.1	260	20.8	26.1	1,799	17.5	15.2	3,895	18.6	17.7
Unknown	23,625	7.8	8.4	18,997	7.6	7.8	3,009	7.7	11.0	98	7.9	11.6	734	7.1	6.6	1,733	8.3	8.4

[1] Abbreviations: AI/AN CHSDA - American Indian/Alaskan Native Indian Health Services Contract Health Service Delivery Areas (See Appendix G); API - Asian/Pacific Islander
[2] Rates are per 100,000 population and are age-adjusted by five-year age groups to the 2000 U.S. standard population based on single years of age.
- Counts and percents are suppressed when fewer than 6 cases were reported for the specific stage.
Note: Percentages based upon small cell sizes should be used with caution.

Stage at Diagnosis by Age and Race and Hispanic Ethnicity for Select Cancer Sites
NAACCR United States Combined, 2010-2014

Female Breast, 2010-2014

	All Races			White			Black			AI/AN CHSDA[1]			API[1]			Hispanic, All Races		
Stage at Diagnosis	Count	%	Rate[2]	Count	%	Rate[2]	Count	%	Rate[2]	Count	%	Rate[2]	Count	%	Rate[2]	Count	%	Rate[2]
All Ages																		
In Situ	271,007	19.7	30.6	219,011	19.2	30.3	34,037	20.7	31.7	737	16.1	18.7	13,329	24.1	28.5	20,446	19.6	21.9
Localized	699,170	50.7	77.4	591,897	52.0	79.4	71,410	43.4	67.5	2,222	48.6	58.4	26,780	48.4	58.1	47,698	45.7	53.3
Regional	309,943	22.5	35.4	249,574	21.9	34.9	43,810	26.6	41.0	1,195	26.1	30.6	12,227	22.1	26.0	27,925	26.8	29.4
Distant	64,478	4.7	7.1	50,521	4.4	6.8	11,197	6.8	10.5	224	4.9	6.0	1,977	3.6	4.2	5,106	4.9	5.6
Unknown	34,058	2.5	3.6	27,071	2.4	3.4	4,137	2.5	4.1	198	4.3	5.6	1,032	1.9	2.3	3,144	3.0	3.6
Age < 50																		
In Situ	60,028	22.2	12.2	46,761	22.4	12.2	7,570	19.2	10.7	178	16.5	6.3	4,291	24.8	12.9	6,516	20.0	8.3
Localized	115,583	42.7	23.5	90,670	43.5	23.8	15,212	38.5	21.4	444	41.1	15.5	7,664	44.3	22.8	12,930	39.7	16.2
Regional	78,779	29.1	16.2	59,661	28.6	15.8	13,150	33.3	18.5	359	33.2	12.6	4,519	26.1	13.4	10,691	32.8	13.2
Distant	11,702	4.3	2.4	8,183	3.9	2.2	2,759	7.0	3.9	62	5.7	2.2	542	3.1	1.6	1,598	4.9	2.0
Unknown	4,671	1.7	1.0	3,202	1.5	0.8	788	2.0	1.1	37	3.4	1.3	274	1.6	0.8	850	2.6	1.0
Age ≥ 50																		
In Situ	210,979	19.0	78.9	172,250	18.5	77.8	26,467	21.2	86.6	559	16.0	50.9	9,038	23.7	69.1	13,930	19.4	57.5
Localized	583,587	52.7	218.7	501,227	53.9	225.1	56,198	44.9	188.2	1,778	50.9	170.8	19,116	50.2	150.4	34,768	48.5	150.3
Regional	231,164	20.9	85.9	189,913	20.4	84.9	30,660	24.5	99.9	836	23.9	77.8	7,708	20.3	59.2	17,234	24.0	71.7
Distant	52,776	4.8	19.6	42,338	4.6	18.8	8,438	6.7	27.9	162	4.6	16.2	1,435	3.8	11.2	3,508	4.9	15.0
Unknown	29,387	2.7	10.6	23,869	2.6	10.1	3,349	2.7	11.8	161	4.6	16.8	758	2.0	6.2	2,294	3.2	10.4

Lung and Bronchus, 2010-2014

	All Races			White			Black			AI/AN CHSDA[1]			API[1]			Hispanic, All Races		
Stage at Diagnosis	Count	%	Rate[2]	Count	%	Rate[2]	Count	%	Rate[2]	Count	%	Rate[2]	Count	%	Rate[2]	Count	%	Rate[2]
Male																		
In Situ	876	0.2	0.1	782	0.2	0.1	61	0.1	0.1	-	-	-	23	0.2	0.1	33	0.1	0.1
Localized	94,808	17.1	12.8	82,642	17.5	12.9	9,179	14.6	13.2	272	14.2	10.6	2,141	15.1	7.1	3,368	14.4	6.1
Regional	124,099	22.3	16.2	106,641	22.5	16.3	13,495	21.5	18.3	411	21.5	15.2	2,856	20.1	9.0	4,864	20.7	8.5
Distant	287,622	51.7	37.4	242,022	51.1	36.8	34,891	55.5	46.2	982	51.3	35.7	8,180	57.7	25.6	12,869	54.9	21.6
Unknown	48,597	8.7	6.7	41,454	8.8	6.6	5,282	8.4	8.1	246	12.9	10.5	979	6.9	3.5	2,323	9.9	4.4
Female																		
In Situ	1,169	0.2	0.1	1,007	0.2	0.1	112	0.2	0.1	-	-	-	44	0.4	0.1	59	0.3	0.1
Localized	105,163	21.3	11.4	92,742	21.7	11.9	9,314	18.6	9.3	317	16.9	9.5	2,187	18.9	5.3	3,966	20.4	5.2
Regional	110,954	22.5	12.0	96,087	22.5	12.4	11,711	23.3	11.4	435	23.2	12.8	2,211	19.1	5.4	3,937	20.3	5.1
Distant	237,495	48.1	25.4	203,592	47.6	26.0	25,569	50.9	24.8	922	49.2	26.8	6,301	54.5	15.1	9,667	49.7	12.4
Unknown	39,417	8.0	4.1	34,436	8.0	4.2	3,504	7.0	3.6	201	10.7	6.8	810	7.0	2.1	1,805	9.3	2.5

Prostate, 2010-2014

	All Races			White			Black			AI/AN CHSDA[1]			API[1]			Hispanic, All Races		
Stage at Diagnosis	Count	%	Rate[2]	Count	%	Rate[2]	Count	%	Rate[2]	Count	%	Rate[2]	Count	%	Rate[2]	Count	%	Rate[2]
All Ages																		
Localized	708,911	75.1	85.6	555,296	75.5	79.1	114,610	76.2	136.5	1,515	65.6	48.6	14,287	71.5	41.1	45,979	72.4	68.7
Regional	110,328	11.7	12.6	91,120	12.4	12.3	14,829	9.9	16.0	285	12.3	7.9	2,696	13.5	7.2	7,220	11.4	9.7
Distant	50,019	5.3	6.7	38,676	5.3	6.1	9,419	6.3	13.3	181	7.8	7.5	1,226	6.1	4.1	3,957	6.2	6.9
Unknown	74,597	7.9	10.1	50,150	6.8	7.9	11,495	7.6	17.7	330	14.3	14.3	1,763	8.8	6.0	6,355	10.0	11.8
Age < 65																		
Localized	312,456	77.0	36.8	232,959	77.3	33.0	63,611	77.9	66.6	672	69.1	17.3	5,304	73.2	13.1	20,588	74.8	24.2
Regional	59,814	14.7	7.0	47,489	15.8	6.7	10,136	12.4	10.6	171	17.6	4.4	1,246	17.2	3.1	3,936	14.3	4.6
Distant	14,628	3.6	1.7	10,076	3.3	1.4	4,025	4.9	4.2	56	5.8	1.4	302	4.2	0.7	1,410	5.1	1.7
Unknown	19,000	4.7	2.2	10,828	3.6	1.5	3,839	4.7	4.0	73	7.5	1.9	395	5.5	1.0	1,602	5.8	1.9
Age ≥ 65																		
Localized	396,455	73.7	422.9	322,337	74.3	398.0	50,999	74.2	619.7	843	63.0	265.1	8,983	70.6	234.9	25,391	70.6	376.4
Regional	50,514	9.4	51.1	43,631	10.1	51.2	4,693	6.8	53.4	114	8.5	31.9	1,450	11.4	35.6	3,284	9.1	45.2
Distant	35,391	6.6	41.3	28,600	6.6	38.2	5,394	7.8	76.5	125	9.3	49.0	924	7.3	27.4	2,547	7.1	42.9
Unknown	55,597	10.3	64.4	39,322	9.1	52.5	7,656	11.1	112.6	257	19.2	100.2	1,368	10.8	40.9	4,753	13.2	79.9

[1] Abbreviations: AI/AN CHSDA - American Indian/Alaskan Native Indian Health Services Contract Health Service Delivery Areas (See Appendix G) ; API - Asian/Pacific Islander
[2] Rates are per 100,000 population and are age-adjusted by five-year age groups to the 2000 U.S. standard population based on single years of age.
- Counts and percents are suppressed when fewer than 6 cases were reported for the specific stage.
Note: Percentages based upon small cell sizes should be used with caution.

Stage at Diagnosis by Age for Select Cancer Sites
NAACCR Canada Combined, 2010-2014

Cervix, 2010-2014

	All Races		
Stage at Diagnosis	Count	%	Rate[1]
All Ages			
Localized	1,308	48.8	3.9
Regional	875	32.7	2.4
Distant	307	11.5	0.8
Unknown	189	7.1	0.5

Colon and Rectum, 2010-2014

	All Races		
Stage at Diagnosis	Count	%	Rate[1]
Male			
All Ages			
In Situ	3,362	12.0	9.0
Localized	8,706	31.0	24.3
Regional	9,580	34.1	26.5
Distant	5,333	19.0	14.8
Unknown	1,089	3.9	3.4
Age < 50			
In Situ	184	9.8	0.8
Localized	495	26.4	2.2
Regional	727	38.7	3.2
Distant	434	23.1	1.9
Unknown	38	2.0	0.2
Age ≥ 50			
In Situ	3,178	12.1	30.6
Localized	8,211	31.3	82.4
Regional	8,853	33.8	87.5
Distant	4,899	18.7	48.5
Unknown	1,051	4.0	11.8
Female			
All Ages			
In Situ	1,974	9.1	4.8
Localized	7,052	32.3	17.3
Regional	7,500	34.4	18.2
Distant	4,082	18.7	9.9
Unknown	1,195	5.5	2.7
Age < 50			
In Situ	129	8.3	0.6
Localized	490	31.5	2.2
Regional	517	33.2	2.3
Distant	372	23.9	1.7
Unknown	49	3.1	0.2
Age ≥ 50			
In Situ	1,845	9.1	16.0
Localized	6,562	32.4	56.7
Regional	6,983	34.5	60.0
Distant	3,710	18.3	31.5
Unknown	1,146	5.7	9.0

Female Breast, 2010-2014

	All Races		
Stage at Diagnosis	Count	%	Rate[1]
All Ages			
In Situ	6,878	13.0	17.4
Localized	28,057	52.9	69.7
Regional	14,880	28.0	38.0
Distant	2,720	5.1	6.8
Unknown	515	1.0	1.1
Age < 50			
In Situ	1,456	15.0	6.5
Localized	4,246	43.8	19.0
Regional	3,504	36.2	15.8
Distant	444	4.6	2.0
Unknown	34	0.4	0.2
Age ≥ 50			
In Situ	5,422	12.5	46.0
Localized	23,811	54.9	202.5
Regional	11,376	26.2	96.1
Distant	2,276	5.2	19.4
Unknown	481	1.1	3.7

Lung and Bronchus, 2010-2014

	All Races		
Stage at Diagnosis	Count	%	Rate[1]
Male			
In Situ	44	0.2	0.1
Localized	3,312	14.4	9.5
Regional	5,706	24.8	16.2
Distant	13,429	58.3	37.6
Unknown	552	2.4	1.7
Female			
In Situ	85	0.4	0.2
Localized	3,849	17.4	9.4
Regional	5,593	25.3	13.7
Distant	12,118	54.8	29.4
Unknown	462	2.1	1.1

Prostate, 2010-2014

	All Races		
Stage at Diagnosis	Count	%	Rate[1]
All Ages			
Localized	30,224	70.9	79.1
Regional	7,161	16.8	17.7
Distant	3,712	8.7	11.1
Unknown	1,503	3.5	4.7
Age < 65			
Localized	11,775	72.8	29.0
Regional	3,465	21.4	8.5
Distant	708	4.4	1.7
Unknown	217	1.3	0.5
Age ≥ 65			
Localized	18,449	69.8	424.9
Regional	3,696	14.0	81.1
Distant	3,004	11.4	75.7
Unknown	1,286	4.9	33.6

[1] Rates are per 100,000 population and are age-adjusted by five-year age groups to the 2000 U.S. standard population based on single years of age.
- Counts and percents are suppressed when fewer than 6 cases were reported for the specific stage.
Note: Percentages based upon small cell sizes should be used with caution.

Stage at Diagnosis by Age for Select Cancer Sites
NAACCR North America Combined, 2010-2014

Cervix, 2010-2014

Stage at Diagnosis	All Races Count	%	Rate[1]
All Ages			
Localized	26,876	42.7	3.4
Regional	22,324	35.4	2.6
Distant	9,198	14.6	1.0
Unknown	4,615	7.3	0.5

Colon and Rectum, 2010-2014

Stage at Diagnosis	All Races Count	%	Rate[1]
Male			
All Ages			
In Situ	22,264	5.6	2.7
Localized	140,982	35.4	17.6
Regional	128,785	32.4	16.1
Distant	79,082	19.9	9.7
Unknown	26,691	6.7	3.5
Age < 50			
In Situ	1,452	3.6	0.3
Localized	11,688	29.0	2.3
Regional	15,019	37.3	2.9
Distant	10,276	25.5	2.0
Unknown	1,854	4.6	0.4
Age ≥ 50			
In Situ	20,812	5.8	9.0
Localized	129,294	36.2	57.7
Regional	113,766	31.8	50.5
Distant	68,806	19.2	30.0
Unknown	24,837	6.9	11.7
Female			
All Ages			
In Situ	16,880	4.7	1.7
Localized	129,194	35.9	13.4
Regional	117,788	32.7	12.2
Distant	69,634	19.3	7.2
Unknown	26,485	7.4	2.6
Age < 50			
In Situ	1,418	3.9	0.3
Localized	11,637	31.9	2.2
Regional	12,535	34.4	2.4
Distant	9,119	25.0	1.8
Unknown	1,714	4.7	0.3
Age ≥ 50			
In Situ	15,462	4.8	5.6
Localized	117,557	36.3	42.5
Regional	105,253	32.5	37.8
Distant	60,515	18.7	21.6
Unknown	24,771	7.7	8.4

Female Breast, 2010-2014

Stage at Diagnosis	All Races Count	%	Rate[1]
All Ages			
In Situ	277,885	19.4	30.0
Localized	727,227	50.8	77.1
Regional	324,823	22.7	35.5
Distant	67,198	4.7	7.1
Unknown	34,573	2.4	3.5
Age < 50			
In Situ	61,484	21.9	11.9
Localized	119,829	42.7	23.3
Regional	82,283	29.3	16.1
Distant	12,146	4.3	2.4
Unknown	4,705	1.7	0.9
Age ≥ 50			
In Situ	216,401	18.8	77.5
Localized	607,398	52.8	218.0
Regional	242,540	21.1	86.3
Distant	55,052	4.8	19.6
Unknown	29,868	2.6	10.3

Lung and Bronchus, 2010-2014

Stage at Diagnosis	All Races Count	%	Rate[1]
Male			
In Situ	920	0.2	0.1
Localized	98,120	16.9	12.6
Regional	129,805	22.4	16.2
Distant	301,051	52.0	37.4
Unknown	49,149	8.5	6.5
Female			
In Situ	1,254	0.2	0.1
Localized	109,012	21.1	11.3
Regional	116,547	22.6	12.0
Distant	249,613	48.3	25.5
Unknown	39,879	7.7	4.0

Prostate, 2010-2014

Stage at Diagnosis	All Races Count	%	Rate[1]
All Ages			
Localized	739,135	74.9	85.3
Regional	117,489	11.9	12.8
Distant	53,731	5.4	6.9
Unknown	76,100	7.7	9.8
Age < 65			
Localized	324,231	76.8	36.5
Regional	63,279	15.0	7.1
Distant	15,336	3.6	1.7
Unknown	19,217	4.6	2.1
Age ≥ 65			
Localized	414,904	73.5	423.0
Regional	54,210	9.6	52.5
Distant	38,395	6.8	42.8
Unknown	56,883	10.1	63.0

[1] Rates are per 100,000 population and are age-adjusted by five-year age groups to the 2000 U.S. standard population based on single years of age.
- Counts and percents are suppressed when fewer than 6 cases were reported for the specific stage.
Note: Percentages based upon small cell sizes should be used with caution.

Section V

Average-annual Age-adjusted Cancer Incidence Observed and Delay Adjusted Counts and Rates for Select Cancer Sites in the United States, Canada and North America

NAACCR U.S. Combined Delay Adjusted Incidence 2010-2014, All Races Combined

Average Annual Age-adjusted Cancer Incidence Delay Adjusted Counts and Rates*, Selected Sites
NAACCR United States Combined[1], 2010-2014, All Races

	MALES (ALL RACES)						FEMALES (ALL RACES)					
	Observed			Delay Adjusted			Observed			Delay Adjusted		
SITES	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
All Sites (invasive & in situ bladder)	3,822,183	492.0	0.26	3,904,031	502.5	0.26	3,719,881	411.4	0.22	3,800,087	420.3	0.22
Oral Cavity & Pharynx	141,503	17.4	0.05	144,665	17.8	0.05	58,235	6.4	0.03	59,484	6.5	0.03
Digestive												
Esophagus	62,815	8.0	0.03	63,620	8.1	0.03	16,881	1.8	0.01	17,094	1.8	0.01
Stomach	69,817	9.2	0.04	70,690	9.3	0.04	42,821	4.7	0.02	43,354	4.7	0.02
Colon & Rectum	350,832	45.9	0.08	355,894	46.5	0.08	323,272	34.8	0.06	327,845	35.3	0.06
Liver & IBD[2]	100,892	12.1	0.04	103,113	12.4	0.04	39,348	4.2	0.02	40,213	4.3	0.02
Pancreas	109,099	14.2	0.04	110,601	14.4	0.04	104,962	11.1	0.03	106,375	11.2	0.04
Respiratory												
Larynx	48,658	6.0	0.03	49,388	6.1	0.03	12,639	1.4	0.01	12,826	1.4	0.01
Lung & bronchus	555,092	73.0	0.10	562,396	74.0	0.10	492,990	52.8	0.08	499,516	53.5	0.08
Skin												
Melanomas of the skin	201,442	26.6	0.06	207,176	27.3	0.06	141,306	16.3	0.04	145,162	16.8	0.05
Breast (female cases only)	-	-	-	-	-	-	1,107,649	123.6	0.12	1,125,695	125.6	0.12
Female Genital												
Cervix uteri	-	-	-	-	-	-	60,334	7.6	0.03	61,235	7.7	0.03
Corpus & uterus, NOS[2]	-	-	-	-	-	-	241,508	25.9	0.05	244,146	26.1	0.05
Ovary	-	-	-	-	-	-	103,334	11.5	0.04	105,904	11.7	0.04
Male Genital												
Prostate	943,855	115.0	0.12	972,131	118.5	0.12	-	-	-	-	-	-
Testis	40,376	5.5	0.03	40,947	5.6	0.03	-	-	-	-	-	-
Urinary												
Bladder (incl. *in situ*)	262,379	35.8	0.07	267,825	36.6	0.07	83,602	8.8	0.03	85,318	9.0	0.03
Kidney & renal pelvis	171,900	21.8	0.05	175,943	22.4	0.06	102,450	11.3	0.04	104,836	11.6	0.04
Brain & Other Nervous System	59,106	7.7	0.03	60,522	7.9	0.03	47,305	5.5	0.03	48,417	5.7	0.03
Endocrine												
Thyroid gland	55,668	7.2	0.03	56,543	7.3	0.03	169,974	21.1	0.05	172,649	21.5	0.05
Lymphomas												
Hodgkin Lymphoma	22,955	3.1	0.02	23,362	3.1	0.02	18,446	2.4	0.02	18,768	2.4	0.02
Non-Hodgkin Lymphoma	173,091	22.9	0.06	178,116	23.6	0.06	144,038	15.8	0.04	148,195	16.2	0.04
Multiple Myeloma	61,561	8.1	0.03	66,194	8.7	0.04	49,548	5.3	0.02	53,265	5.7	0.03
Leukemias	129,047	17.3	0.05	141,055	18.9	0.05	94,068	10.6	0.04	102,674	11.5	0.04

* Rates are per 100,000 population and are age-adjusted by five-year age groups to the 2000 U.S. standard population based on single years of age and to standard.

- Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for All Sites combined.

[1] NAACCR United States Combined includes registries for which a U.S. average delay adjustment factor was calculated. See technical notes.

[2] Abbreviations: IBD – Intrahepatic Bile Duct; NOS - Not Otherwise Specified

Note: Delay adjusted counts are rounded to the nearest whole number. A small registry and/or rare cancer site may have the same delay adjusted count and observed count, but this is just due to rounding to whole counts. For more information about cancer incidence rates adjusted for reporting delay see https://surveillance.cancer.gov/delay/.

Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Delay Adjusted Counts and Rates[1] for the Five Most Commonly Diagnosed Cancers
Selected Areas in the NAACCR United States Combined[2], 2010-2014, All Races, Males

	All Sites			Prostate			Lung & Bronchus			Colon & Rectum			Bladder			Melanoma of the Skin		
	Count	*Rate*	*S.E.*[3]	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
NAACCR U.S. Combined																		
Observed	3,822,183	492.0	0.26	943,855	115.0	0.12	555,092	73.0	0.10	350,832	45.9	0.08	262,379	35.8	0.07	201,442	26.6	0.06
Delay Adjusted	3,904,031	502.5	0.26	972,131	118.5	0.12	562,396	74.0	0.10	355,894	46.5	0.08	267,825	36.6	0.07	207,176	27.3	0.06
Alabama																		
Observed	66,882	527.2	2.10	17,371	129.3	1.01	11,673	92.5	0.88	6,498	52.2	0.67	4,058	34.0	0.55	3,304	27.2	0.49
Delay Adjusted	67,601	532.9	2.12	17,628	131.2	1.02	11,736	93.0	0.89	6,546	52.6	0.67	4,094	34.3	0.56	3,343	27.5	0.49
Alaska																		
Observed	6,860	441.3	6.03	1,548	91.4	2.63	937	67.5	2.49	707	46.6	1.99	451	33.8	1.78	248	15.2	1.07
Delay Adjusted	7,102	457.0	6.25	1,625	96.1	2.77	959	69.1	2.55	724	47.7	2.04	461	34.6	1.82	272	16.7	1.18
Arizona																		
Observed	71,408	405.9	1.54	14,985	80.8	0.67	9,840	56.0	0.57	6,778	39.2	0.48	5,534	32.2	0.44	4,429	25.7	0.39
Delay Adjusted	74,823	425.3	1.61	15,940	86.0	0.71	10,066	57.3	0.59	7,109	41.1	0.51	5,776	33.6	0.46	4,752	27.5	0.42
Arkansas																		
Observed	41,512	519.2	2.60	10,204	120.7	1.22	7,946	99.0	1.13	3,937	50.1	0.82	2,745	36.0	0.70	1,921	25.1	0.59
Delay Adjusted	42,037	525.8	2.64	10,353	122.4	1.23	8,009	99.8	1.14	3,966	50.4	0.82	2,770	36.3	0.71	1,968	25.7	0.60
California																		
Observed	399,638	450.0	0.73	100,652	109.3	0.35	42,874	51.0	0.25	37,576	42.5	0.22	25,904	31.4	0.20	25,399	28.9	0.19
Delay Adjusted	405,980	457.2	0.74	102,908	111.8	0.36	43,375	51.6	0.25	37,933	42.9	0.23	26,262	31.8	0.20	25,794	29.4	0.19
Greater California																		
Observed	226,606	458.4	0.98	55,682	108.4	0.47	25,439	54.0	0.35	20,841	42.4	0.30	15,128	32.9	0.27	15,910	32.6	0.26
Delay Adjusted	230,455	466.2	1.00	57,037	111.1	0.48	25,741	54.7	0.35	21,058	42.8	0.30	15,367	33.4	0.28	16,169	33.1	0.27
Greater Bay																		
Observed	78,970	456.0	1.66	20,916	115.8	0.82	7,967	49.0	0.56	7,006	40.6	0.50	4,915	30.4	0.44	5,443	31.8	0.44
Delay Adjusted	79,846	461.1	1.68	21,154	117.2	0.83	8,032	49.4	0.57	7,039	40.8	0.50	4,957	30.7	0.45	5,503	32.2	0.45
Los Angeles																		
Observed	93,893	425.7	1.42	23,993	105.9	0.70	9,448	45.6	0.48	9,717	44.2	0.46	5,855	28.7	0.38	4,043	18.5	0.30
Delay Adjusted	95,502	433.0	1.45	24,652	108.8	0.72	9,582	46.3	0.49	9,824	44.7	0.46	5,932	29.1	0.39	4,119	18.9	0.30
Colorado																		
Observed	54,154	436.3	1.95	14,649	110.9	0.96	5,581	48.3	0.67	4,577	37.9	0.58	3,696	32.7	0.56	3,422	27.8	0.49
Delay Adjusted	55,274	445.3	1.99	15,421	116.7	1.01	5,624	48.7	0.68	4,609	38.1	0.59	3,827	33.9	0.58	3,520	28.6	0.51
Connecticut																		
Observed	49,666	515.9	2.37	12,169	118.8	1.10	6,578	69.7	0.88	4,205	44.4	0.70	4,342	46.9	0.73	2,566	27.4	0.55
Delay Adjusted	50,764	527.4	2.42	12,509	122.3	1.14	6,665	70.7	0.89	4,250	44.9	0.71	4,421	47.7	0.74	2,621	28.0	0.57

Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Delay Adjusted Counts and Rates[1] for the Five Most Commonly Diagnosed Cancers
Selected Areas in the NAACCR United States Combined[2], 2010-2014, All Races, Males

	All Sites			Prostate			Lung & Bronchus			Colon & Rectum			Bladder			Melanoma of the Skin		
	Count	*Rate*	*S.E.[3]*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
Delaware																		
Observed	14,089	558.4	4.81	3,837	141.6	2.33	2,025	81.3	1.85	1,069	43.5	1.36	1,017	42.2	1.35	969	39.5	1.30
Delay Adjusted	14,517	575.4	4.96	4,013	148.1	2.44	2,070	83.2	1.89	1,090	44.4	1.39	1,051	43.6	1.40	998	40.7	1.33
District of Columbia																		
Observed	7,188	529.6	6.46	2,188	159.7	3.52	883	67.4	2.34	641	48.7	1.99	297	23.3	1.39	168	12.4	0.99
Delay Adjusted	7,418	546.5	6.67	2,238	163.4	3.61	900	68.7	2.39	659	50.1	2.05	301	23.7	1.41	174	12.8	1.03
Florida																		
Observed	280,506	467.8	0.90	65,177	103.6	0.41	43,753	71.4	0.34	25,244	42.5	0.27	20,090	33.2	0.24	17,089	29.2	0.23
Delay Adjusted	291,471	486.1	0.93	68,947	109.5	0.43	44,086	72.0	0.35	25,967	43.8	0.28	20,809	34.4	0.24	18,305	31.2	0.24
Georgia																		
Observed	116,120	522.7	1.61	30,481	129.3	0.77	17,951	84.6	0.66	10,783	48.9	0.49	6,580	33.3	0.43	7,314	33.9	0.42
Delay Adjusted	118,058	531.6	1.64	31,040	131.8	0.79	18,150	85.5	0.67	10,903	49.4	0.50	6,684	33.8	0.44	7,474	34.6	0.42
Hawaii																		
Observed	16,765	434.7	3.44	3,582	90.1	1.55	2,178	56.9	1.25	2,003	51.3	1.18	893	23.7	0.81	1,023	26.5	0.85
Delay Adjusted	16,928	438.9	3.48	3,627	91.3	1.56	2,194	57.3	1.26	2,013	51.6	1.18	901	23.9	0.82	1,037	26.9	0.86
Idaho																		
Observed	19,216	467.8	3.46	5,187	119.7	1.70	2,219	55.7	1.21	1,612	39.8	1.01	1,479	37.9	1.01	1,311	32.2	0.91
Delay Adjusted	19,719	480.1	3.55	5,427	125.3	1.78	2,265	56.8	1.24	1,633	40.3	1.03	1,560	40.0	1.06	1,343	33.0	0.93
Illinois																		
Observed	161,813	511.8	1.30	39,845	119.4	0.61	24,273	79.2	0.52	16,315	52.5	0.42	11,139	37.5	0.36	7,142	23.0	0.28
Delay Adjusted	164,162	519.3	1.32	40,779	122.3	0.63	24,508	80.0	0.53	16,430	52.8	0.42	11,288	38.0	0.37	7,302	23.5	0.28
Indiana																		
Observed	80,261	485.8	1.76	16,713	95.7	0.76	14,434	88.9	0.76	7,932	48.8	0.56	5,747	36.7	0.50	3,539	21.8	0.38
Delay Adjusted	82,439	499.0	1.81	17,279	99.0	0.79	14,614	90.0	0.77	8,088	49.7	0.57	5,922	37.8	0.51	3,720	22.9	0.40
Iowa																		
Observed	43,899	517.1	2.51	10,064	112.2	1.14	6,611	78.3	0.98	4,310	51.7	0.80	3,190	38.4	0.69	2,394	29.1	0.61
Delay Adjusted	44,548	524.8	2.55	10,223	114.0	1.16	6,694	79.3	0.99	4,344	52.1	0.81	3,230	38.8	0.70	2,431	29.5	0.62
Kansas																		
Observed	37,656	511.0	2.69	9,518	124.1	1.30	5,305	73.1	1.02	3,439	47.4	0.82	2,748	38.8	0.75	2,167	30.2	0.66
Delay Adjusted	^	^	^	^	^	^	^	^	^	^	^	^	^	^	^	^	^	^

1. Rates are per 100,000 population and are age-adjusted by five-year age groups to the 2000 U.S. standard population based on single years of age.
2. See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.
3. Abbreviations: S.E. - Standard Error

^ There are insufficient historical registry data to calculate delay factors necessary for delay adjusted rates.

\- Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for All Sites combined.

Note: Delay adjusted counts are rounded to the nearest whole number. A small registry and/or rare cancer site may have the same delay adjusted count and observed count, but this is just due to rounding to whole counts. For more information about cancer incidence rates adjusted for reporting delay see https://surveillance.cancer.gov/delay/.

Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Delay Adjusted Counts and Rates[1] for the Five Most Commonly Diagnosed Cancers
Selected Areas in the NAACCR United States Combined[2], 2010-2014, All Races, Males

	All Sites			Prostate			Lung & Bronchus			Colon & Rectum			Bladder			Melanoma of the Skin		
	Count	*Rate*	*S.E.*[3]	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
Kentucky																		
Observed	66,211	578.2	2.32	13,778	113.0	0.99	13,202	116.2	1.04	6,687	59.3	0.75	4,304	40.2	0.63	3,403	30.7	0.54
Delay Adjusted	67,361	588.3	2.36	14,148	116.0	1.02	13,350	117.6	1.06	6,802	60.3	0.76	4,382	41.0	0.65	3,459	31.2	0.55
Louisiana																		
Observed	64,191	565.4	2.31	17,311	144.4	1.13	10,065	90.8	0.93	6,303	56.0	0.73	3,530	33.5	0.58	2,482	23.0	0.48
Delay Adjusted	64,936	571.9	2.33	17,528	146.3	1.15	10,144	91.5	0.94	6,344	56.3	0.73	3,566	33.9	0.59	2,523	23.3	0.48
Maine																		
Observed	20,845	505.3	3.61	4,491	99.8	1.53	3,464	83.9	1.46	1,717	42.7	1.06	1,892	47.5	1.12	1,134	28.1	0.86
Delay Adjusted	21,135	512.4	3.66	4,560	101.3	1.55	3,503	84.9	1.48	1,728	43.0	1.07	1,922	48.3	1.13	1,153	28.6	0.88
Maryland																		
Observed	71,651	494.2	1.90	20,406	131.5	0.95	9,265	66.6	0.71	6,025	42.6	0.56	4,916	37.0	0.54	4,032	28.5	0.46
Delay Adjusted	73,497	506.9	1.95	21,106	136.0	0.98	9,475	68.2	0.73	6,144	43.4	0.58	5,123	38.5	0.57	4,051	28.7	0.47
Massachusetts																		
Observed	86,303	495.1	1.72	21,277	114.2	0.80	11,875	70.3	0.66	7,327	42.9	0.51	6,718	40.5	0.50	4,332	25.4	0.40
Delay Adjusted	87,627	502.7	1.75	21,745	116.7	0.82	12,079	71.5	0.67	7,416	43.4	0.52	6,819	41.1	0.51	4,453	26.1	0.41
Michigan																		
Observed	136,177	506.3	1.41	36,195	126.3	0.68	20,447	77.2	0.55	11,508	44.0	0.42	9,972	39.1	0.40	5,904	22.7	0.30
Delay Adjusted	138,313	514.3	1.43	36,663	128.0	0.69	20,804	78.6	0.56	11,680	44.6	0.43	10,102	39.6	0.41	6,005	23.1	0.31
Detroit																		
Observed	56,711	571.2	2.47	16,298	155.2	1.25	8,151	85.0	0.97	4,722	48.5	0.73	3,965	42.4	0.69	2,041	21.2	0.48
Delay Adjusted	57,232	576.5	2.49	16,424	156.4	1.26	8,198	85.5	0.98	4,752	48.8	0.73	4,015	42.9	0.70	2,054	21.3	0.48
Mississippi																		
Observed	41,223	550.5	2.80	10,748	135.5	1.34	7,466	101.4	1.21	4,300	58.2	0.92	2,145	30.9	0.69	1,662	23.1	0.59
Delay Adjusted	42,074	561.8	2.85	11,018	139.0	1.38	7,568	102.8	1.23	4,347	58.8	0.93	2,193	31.6	0.70	1,691	23.5	0.60
Missouri																		
Observed	79,601	492.7	1.78	17,428	101.0	0.78	14,217	89.0	0.76	7,820	49.7	0.57	5,198	33.8	0.48	3,831	24.3	0.40
Delay Adjusted	81,678	505.6	1.83	18,224	105.7	0.82	14,465	90.5	0.77	7,955	50.5	0.58	5,342	34.7	0.49	3,940	25.0	0.41
Montana																		
Observed	14,426	473.8	4.07	3,856	116.6	1.93	1,810	60.9	1.47	1,334	44.2	1.25	1,056	36.0	1.13	897	30.3	1.05
Delay Adjusted	14,988	492.2	4.23	4,036	122.1	2.02	1,849	62.2	1.50	1,379	45.7	1.29	1,105	37.6	1.19	941	31.8	1.10
Nebraska																		
Observed	23,934	499.1	3.29	5,977	119.6	1.58	3,374	71.8	1.26	2,375	49.9	1.04	1,704	37.1	0.91	1,122	23.5	0.71
Delay Adjusted	24,713	515.4	3.39	6,288	125.8	1.66	3,446	73.4	1.28	2,429	51.1	1.07	1,791	39.0	0.96	1,165	24.4	0.74

Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Delay Adjusted Counts and Rates[1] for the Five Most Commonly Diagnosed Cancers
Selected Areas in the NAACCR United States Combined[2], 2010-2014, All Races, Males

	All Sites			Prostate			Lung & Bronchus			Colon & Rectum			Bladder			Melanoma of the Skin		
	Count	*Rate*	*S.E.*[3]	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
New Hampshire																		
Observed	19,457	517.0	3.84	5,094	123.5	1.79	2,580	71.1	1.45	1,504	40.9	1.09	1,714	48.1	1.20	1,218	33.3	0.99
Delay Adjusted	19,895	528.7	3.93	5,208	126.3	1.83	2,634	72.7	1.48	1,525	41.4	1.11	1,744	49.0	1.22	1,256	34.4	1.02
New Jersey																		
Observed	120,262	530.5	1.56	33,215	139.4	0.78	14,380	65.6	0.56	10,728	47.9	0.47	8,887	41.5	0.45	6,416	28.7	0.37
Delay Adjusted	122,331	539.6	1.59	33,820	142.0	0.80	14,554	66.4	0.57	10,846	48.4	0.48	9,022	42.2	0.46	6,498	29.1	0.37
New York																		
Observed	264,632	532.9	1.06	71,412	136.8	0.52	33,861	69.9	0.39	22,839	46.9	0.32	19,314	41.0	0.30	11,099	22.8	0.22
Delay Adjusted	267,652	539.0	1.07	72,670	139.3	0.53	34,024	70.2	0.39	22,983	47.2	0.32	19,561	41.5	0.30	11,343	23.3	0.23
North Carolina																		
Observed	127,959	517.9	1.49	32,878	125.0	0.71	21,414	88.5	0.62	10,565	43.7	0.44	8,158	35.5	0.40	7,449	31.4	0.37
Delay Adjusted	131,935	534.0	1.54	34,737	132.0	0.75	21,715	89.7	0.63	10,781	44.5	0.45	8,387	36.5	0.42	7,778	32.8	0.39
North Dakota																		
Observed	9,433	498.4	5.24	2,446	123.2	2.55	1,308	70.8	1.99	1,000	53.8	1.74	683	37.0	1.44	466	24.9	1.18
Delay Adjusted	9,702	512.6	5.39	2,527	127.3	2.63	1,340	72.6	2.04	1,030	55.4	1.79	710	38.5	1.50	479	25.6	1.21
Ohio																		
Observed	155,426	498.3	1.29	37,060	111.8	0.59	25,857	83.7	0.53	14,626	47.9	0.41	11,494	38.6	0.37	7,519	24.9	0.29
Delay Adjusted	160,029	513.1	1.33	38,302	115.5	0.61	26,465	85.7	0.54	14,943	48.9	0.41	11,802	39.7	0.38	7,783	25.8	0.30
Oklahoma																		
Observed	48,596	494.4	2.29	11,133	108.3	1.05	8,448	86.6	0.96	4,705	48.6	0.72	3,158	33.6	0.61	2,276	23.9	0.51
Delay Adjusted	49,322	501.8	2.33	11,308	110.0	1.06	8,549	87.6	0.97	4,754	49.1	0.73	3,196	34.0	0.62	2,284	24.0	0.51
Oregon																		
Observed	50,186	461.0	2.11	11,791	101.1	0.96	6,714	63.1	0.79	4,313	40.8	0.64	3,946	38.0	0.62	3,324	31.1	0.55
Delay Adjusted	50,986	468.4	2.15	12,003	103.0	0.98	6,819	64.1	0.80	4,360	41.2	0.64	3,987	38.4	0.63	3,369	31.6	0.56
Pennsylvania																		
Observed	192,296	532.1	1.24	45,104	117.5	0.56	28,145	78.3	0.47	17,716	49.9	0.38	15,434	43.9	0.36	10,098	28.7	0.29
Delay Adjusted	195,245	540.3	1.26	46,308	120.6	0.58	28,403	79.0	0.48	17,857	50.3	0.38	15,624	44.5	0.36	10,263	29.1	0.30
Rhode Island																		
Observed	14,377	511.4	4.37	3,240	108.5	1.95	2,139	77.7	1.72	1,135	41.2	1.25	1,234	45.1	1.31	781	28.5	1.04
Delay Adjusted	14,638	520.7	4.45	3,345	112.0	2.02	2,169	78.8	1.74	1,145	41.6	1.26	1,258	46.0	1.34	794	28.9	1.06

1. Rates are per 100,000 population and are age-adjusted by five-year age groups to the 2000 U.S. standard population based on single years of age.
2. See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.
3. Abbreviations: S.E. - Standard Error

^ There are insufficient historical registry data to calculate delay factors necessary for delay adjusted rates.

\- Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for All Sites combined.

Note: Delay adjusted counts are rounded to the nearest whole number. A small registry and/or rare cancer site may have the same delay adjusted count and observed count, but this is just due to rounding to whole counts. For more information about cancer incidence rates adjusted for reporting delay see https://surveillance.cancer.gov/delay/.

Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Delay Adjusted Counts and Rates[1] for the Five Most Commonly Diagnosed Cancers
Selected Areas in the NAACCR United States Combined[2], 2010-2014, All Races, Males

	All Sites			Prostate			Lung & Bronchus			Colon & Rectum			Bladder			Melanoma of the Skin		
	Count	*Rate*	*S.E.*[3]	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
South Carolina																		
Observed	64,615	512.6	2.09	16,502	121.4	0.98	10,737	85.7	0.86	5,531	44.7	0.62	4,026	34.7	0.57	3,466	28.4	0.50
Delay Adjusted	66,672	529.0	2.16	17,158	126.3	1.02	10,979	87.7	0.88	5,677	45.9	0.64	4,151	35.8	0.59	3,564	29.2	0.51
South Dakota																		
Observed	11,178	489.3	4.72	2,815	116.8	2.24	1,568	69.7	1.79	1,165	51.6	1.54	771	35.1	1.28	558	24.9	1.08
Delay Adjusted	11,694	511.9	4.94	3,013	125.0	2.40	1,623	72.1	1.85	1,205	53.3	1.59	830	37.8	1.38	599	26.7	1.16
Tennessee																		
Observed	88,122	520.4	1.81	21,512	119.1	0.84	16,027	95.7	0.78	7,724	46.8	0.55	5,466	34.5	0.48	4,232	26.0	0.41
Delay Adjusted	89,744	530.0	1.84	21,946	121.5	0.85	16,210	96.8	0.79	7,800	47.3	0.56	5,553	35.0	0.49	4,329	26.6	0.42
Texas																		
Observed	256,452	453.4	0.93	58,205	99.4	0.43	36,316	68.0	0.37	26,047	46.3	0.30	13,966	27.4	0.24	9,825	17.7	0.19
Delay Adjusted	263,690	466.2	0.95	60,569	103.5	0.44	36,909	69.1	0.38	26,411	46.9	0.30	14,309	28.1	0.25	10,190	18.4	0.19
Utah																		
Observed	25,089	449.1	2.91	7,404	130.6	1.56	1,717	32.7	0.81	1,942	35.5	0.82	1,515	29.8	0.78	2,643	47.7	0.95
Delay Adjusted	25,483	456.2	2.96	7,540	133.0	1.58	1,741	33.2	0.82	1,958	35.8	0.83	1,539	30.3	0.80	2,679	48.3	0.96
Vermont																		
Observed	8,998	486.2	5.31	1,989	98.4	2.28	1,325	71.8	2.04	721	39.3	1.52	691	38.9	1.52	715	39.5	1.54
Delay Adjusted	9,117	492.6	5.38	2,015	99.7	2.31	1,340	72.6	2.06	731	39.9	1.54	705	39.7	1.55	725	40.1	1.56
Virginia																		
Observed	92,237	454.4	1.55	23,473	107.6	0.72	14,290	73.1	0.63	8,205	41.1	0.47	5,791	31.1	0.42	4,861	24.6	0.36
Delay Adjusted	94,017	463.1	1.58	23,972	109.9	0.74	14,454	73.9	0.64	8,295	41.5	0.47	5,880	31.6	0.43	4,954	25.1	0.37
Washington																		
Observed	86,524	490.7	1.72	21,829	116.3	0.81	11,011	65.5	0.64	7,123	40.9	0.50	6,175	37.6	0.49	5,344	30.5	0.43
Delay Adjusted	87,636	497.0	1.74	22,113	117.9	0.82	11,163	66.4	0.65	7,195	41.3	0.50	6,273	38.2	0.50	5,436	31.0	0.44
Seattle																		
Observed	60,612	511.2	2.15	15,504	123.3	1.03	7,398	66.1	0.79	4,930	41.8	0.61	4,246	38.9	0.62	3,978	33.7	0.55
Delay Adjusted	60,968	514.2	2.16	15,596	124.1	1.03	7,417	66.3	0.80	4,942	41.9	0.62	4,271	39.1	0.62	4,030	34.1	0.56
West Virginia																		
Observed	29,078	514.8	3.11	6,057	99.6	1.31	5,639	98.7	1.35	2,939	53.1	1.01	2,175	39.9	0.88	1,350	25.3	0.71
Delay Adjusted	29,717	526.2	3.18	6,175	101.5	1.34	5,735	100.4	1.37	2,988	54.0	1.03	2,219	40.7	0.90	1,394	26.1	0.73
Wisconsin																		
Observed	78,197	502.4	1.84	19,236	116.0	0.86	10,632	69.5	0.69	6,648	43.6	0.55	5,912	39.9	0.53	4,173	27.3	0.43
Delay Adjusted	79,969	513.7	1.88	20,495	123.6	0.91	10,855	70.9	0.70	6,798	44.6	0.56	6,040	40.8	0.54	4,362	28.5	0.45

Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Delay Adjusted Counts and Rates[1] for the Five Most Commonly Diagnosed Cancers
Selected Areas in the NAACCR United States Combined[2], 2010-2014, All Races, Males

	All Sites			Prostate			Lung & Bronchus			Colon & Rectum			Bladder			Melanoma of the Skin		
	Count	*Rate*	*S.E.*[3]	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
Wyoming																		
Observed	6,874	441.1	5.53	1,823	108.5	2.64	738	49.4	1.89	634	41.7	1.72	524	36.0	1.63	425	28.5	1.43
Delay Adjusted	6,940	445.3	5.58	1,837	109.3	2.66	742	49.6	1.90	636	41.8	1.72	527	36.2	1.64	438	29.4	1.48

1. Rates are per 100,000 population and are age-adjusted by five-year age groups to the 2000 U.S. standard population based on single years of age.
2. See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.
3. Abbreviations: S.E. - Standard Error

^ There are insufficient historical registry data to calculate delay factors necessary for delay adjusted rates.

\- Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for All Sites combined.

Note: Delay adjusted counts are rounded to the nearest whole number. A small registry and/or rare cancer site may have the same delay adjusted count and observed count, but this is just due to rounding to whole counts. For more information about cancer incidence rates adjusted for reporting delay see https://surveillance.cancer.gov/delay/.

Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Delay Adjusted Counts and Rates[1] for the Five Most Commonly Diagnosed Cancers
Selected Areas in the NAACCR United States Combined[2], 2010-2014, All Races, Females

	All Sites			Breast			Lung & Bronchus			Colon & Rectum			Uterus Corpus & Uterus, NOS[3]			Thyroid		
	Count	*Rate*	*S.E.*[3]	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
NAACCR U.S. Combined																		
Observed	3,719,881	411.4	0.22	1,107,649	123.6	0.12	492,990	52.8	0.08	323,272	34.8	0.06	241,508	25.9	0.05	169,974	21.1	0.05
Delay Adjusted	3,800,087	420.3	0.22	1,125,695	125.6	0.12	499,516	53.5	0.08	327,845	35.3	0.06	244,146	26.1	0.05	172,649	21.5	0.05
Alabama																		
Observed	58,705	390.8	1.65	17,811	119.6	0.92	8,235	52.1	0.58	5,651	36.9	0.50	2,968	19.1	0.36	1,712	13.0	0.32
Delay Adjusted	59,362	395.3	1.67	18,058	121.2	0.93	8,279	52.4	0.58	5,693	37.2	0.50	2,982	19.2	0.36	1,725	13.1	0.32
Alaska																		
Observed	6,483	405.9	5.34	2,085	125.4	2.90	773	53.3	2.04	620	40.2	1.71	431	24.6	1.26	299	17.5	1.05
Delay Adjusted	6,715	420.4	5.53	2,126	127.9	2.96	790	54.5	2.09	635	41.2	1.76	445	25.4	1.29	307	17.9	1.07
Arizona																		
Observed	71,467	370.0	1.41	21,537	112.4	0.79	9,262	45.4	0.48	5,908	29.8	0.40	4,260	21.6	0.34	3,979	23.7	0.39
Delay Adjusted	74,887	387.7	1.48	22,597	118.0	0.82	9,475	46.4	0.49	6,194	31.3	0.41	4,460	22.6	0.35	4,186	24.9	0.41
Arkansas																		
Observed	35,609	392.7	2.13	10,122	112.7	1.15	5,818	60.7	0.81	3,456	37.1	0.64	1,997	21.6	0.49	1,180	15.1	0.45
Delay Adjusted	36,067	397.7	2.16	10,244	114.0	1.16	5,867	61.2	0.81	3,480	37.4	0.65	2,007	21.7	0.50	1,207	15.5	0.46
California																		
Observed	397,924	382.6	0.62	125,201	120.7	0.35	41,594	39.9	0.20	34,469	32.6	0.18	26,391	24.7	0.15	18,747	19.1	0.14
Delay Adjusted	404,285	388.7	0.63	126,328	121.8	0.35	42,089	40.4	0.20	34,787	32.9	0.18	26,532	24.8	0.15	18,920	19.2	0.14
Greater California																		
Observed	221,624	389.4	0.84	69,313	122.1	0.47	24,872	43.4	0.28	18,789	32.6	0.24	13,933	23.7	0.20	10,305	19.4	0.19
Delay Adjusted	225,414	396.0	0.85	70,018	123.3	0.48	25,176	44.0	0.28	18,981	32.9	0.24	14,015	23.9	0.21	10,409	19.5	0.20
Greater Bay																		
Observed	78,096	386.0	1.41	25,087	124.6	0.80	7,911	39.0	0.45	6,667	32.3	0.40	5,363	25.7	0.36	3,212	16.9	0.30
Delay Adjusted	78,978	390.3	1.42	25,227	125.3	0.80	7,977	39.3	0.45	6,696	32.4	0.40	5,379	25.7	0.36	3,231	17.0	0.30
Los Angeles																		
Observed	98,091	364.9	1.18	30,772	114.6	0.66	8,799	32.9	0.36	9,004	32.9	0.35	7,095	25.8	0.31	5,229	20.0	0.28
Delay Adjusted	99,775	371.2	1.20	31,052	115.7	0.67	8,924	33.4	0.36	9,101	33.3	0.36	7,137	26.0	0.31	5,279	20.2	0.28
Colorado																		
Observed	53,876	384.6	1.69	17,541	123.7	0.95	5,751	41.6	0.56	4,259	30.8	0.48	3,321	22.3	0.40	3,154	23.5	0.43
Delay Adjusted	54,994	392.6	1.73	17,615	124.2	0.96	5,795	41.9	0.57	4,289	31.0	0.48	3,339	22.4	0.40	3,183	23.8	0.43
Connecticut																		
Observed	51,826	451.0	2.04	15,767	139.2	1.14	6,766	56.8	0.71	4,159	34.3	0.55	3,483	29.2	0.50	2,827	29.1	0.56
Delay Adjusted	52,979	461.0	2.09	16,022	141.4	1.16	6,857	57.5	0.71	4,205	34.6	0.56	3,513	29.4	0.51	2,863	29.4	0.57

Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Delay Adjusted Counts and Rates[1] for the Five Most Commonly Diagnosed Cancers
Selected Areas in the NAACCR United States Combined[2], 2010-2014, All Races, Females

	All Sites			Breast			Lung & Bronchus			Colon & Rectum			Uterus Corpus & Uterus, NOS[3]			Thyroid		
	Count	*Rate*	*S.E.[3]*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
Delaware																		
Observed	13,172	449.8	4.02	3,886	133.1	2.20	1,930	62.9	1.45	986	32.9	1.07	890	29.0	1.00	578	23.6	1.01
Delay Adjusted	13,578	463.6	4.15	3,943	135.1	2.23	1,973	64.4	1.49	1,005	33.5	1.09	898	29.3	1.01	593	24.2	1.04
District of Columbia																		
Observed	7,556	443.0	5.23	2,389	143.5	3.00	845	49.5	1.74	690	40.3	1.57	549	31.2	1.36	396	22.5	1.18
Delay Adjusted	7,815	458.5	5.41	2,525	151.6	3.17	862	50.5	1.77	709	41.4	1.62	561	31.9	1.39	416	23.7	1.24
Florida																		
Observed	260,388	389.0	0.79	75,212	115.5	0.44	38,670	53.1	0.27	22,825	32.3	0.22	15,788	23.1	0.19	9,412	17.6	0.19
Delay Adjusted	270,808	404.7	0.82	78,166	120.0	0.45	38,964	53.5	0.28	23,458	33.3	0.23	16,292	23.9	0.20	9,473	17.7	0.19
Georgia																		
Observed	108,740	403.5	1.24	33,614	123.5	0.68	14,071	52.2	0.45	9,578	35.6	0.37	6,133	21.6	0.28	4,359	16.6	0.26
Delay Adjusted	110,546	410.3	1.26	33,933	124.7	0.69	14,226	52.7	0.45	9,684	36.0	0.37	6,173	21.8	0.28	4,400	16.8	0.26
Hawaii																		
Observed	16,988	401.4	3.19	5,670	136.0	1.87	1,717	38.1	0.94	1,599	36.5	0.94	1,237	30.0	0.88	831	23.2	0.83
Delay Adjusted	17,156	405.3	3.22	5,702	136.8	1.88	1,730	38.3	0.95	1,607	36.7	0.95	1,238	30.0	0.88	836	23.4	0.84
Idaho																		
Observed	17,729	403.5	3.10	5,295	120.5	1.69	2,117	46.8	1.03	1,437	32.5	0.88	1,111	24.2	0.74	980	24.9	0.81
Delay Adjusted	18,201	414.2	3.18	5,368	122.1	1.72	2,160	47.7	1.05	1,457	33.0	0.89	1,130	24.6	0.76	989	25.1	0.82
Illinois																		
Observed	163,740	432.4	1.09	48,665	130.0	0.60	22,371	57.8	0.39	14,948	38.1	0.32	11,336	29.0	0.28	7,370	21.7	0.26
Delay Adjusted	166,148	438.7	1.11	49,154	131.3	0.61	22,591	58.4	0.40	15,053	38.4	0.32	11,432	29.3	0.28	7,452	21.9	0.26
Indiana																		
Observed	81,298	419.5	1.50	23,084	120.1	0.81	12,158	60.8	0.56	7,666	38.5	0.45	5,390	27.0	0.38	3,015	17.8	0.33
Delay Adjusted	83,574	431.4	1.55	23,729	123.4	0.83	12,314	61.6	0.57	7,814	39.3	0.46	5,488	27.5	0.38	3,074	18.2	0.34
Iowa																		
Observed	41,560	432.1	2.20	11,520	122.8	1.19	5,376	53.0	0.74	4,114	39.8	0.64	3,042	31.2	0.58	1,575	20.1	0.52
Delay Adjusted	42,173	438.4	2.23	11,592	123.5	1.19	5,447	53.7	0.75	4,147	40.1	0.65	3,058	31.3	0.58	1,593	20.3	0.53
Kansas																		
Observed	35,187	421.9	2.31	10,182	123.5	1.26	4,596	52.8	0.79	3,129	36.1	0.66	2,104	24.7	0.55	1,833	25.3	0.61
Delay Adjusted	^	^	^	^	^	^	^	^	^	^	^	^	^	^	^	^	^	^

1. Rates are per 100,000 population and are age-adjusted by five-year age groups to the 2000 U.S. standard population based on single years of age.
2. See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.
3. Abbreviations: NOS - Not Otherwise Specified; S.E. - Standard Error

^ There are insufficient historical registry data to calculate delay factors necessary for delay adjusted rates.

\- Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for All Sites combined.

Note: Delay adjusted counts are rounded to the nearest whole number. A small registry and/or rare cancer site may have the same delay adjusted count and observed count, but this is just due to rounding to whole counts. For more information about cancer incidence rates adjusted for reporting delay see https://surveillance.cancer.gov/delay/.

Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Delay Adjusted Counts and Rates[1] for the Five Most Commonly Diagnosed Cancers
Selected Areas in the NAACCR United States Combined[2], 2010-2014, All Races, Females

	All Sites			Breast			Lung & Bronchus			Colon & Rectum			Uterus Corpus & Uterus, NOS[3]			Thyroid		
	Count	*Rate*	*S.E.*[3]	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
Kentucky																		
Observed	61,841	464.1	1.91	16,413	123.3	0.98	11,053	79.7	0.77	5,774	42.4	0.57	3,521	25.8	0.44	2,526	21.6	0.44
Delay Adjusted	62,924	472.2	1.94	16,544	124.3	0.99	11,177	80.6	0.78	5,872	43.1	0.58	3,547	25.9	0.45	2,557	21.9	0.45
Louisiana																		
Observed	55,010	414.1	1.80	16,330	123.2	0.99	7,512	54.9	0.64	5,551	41.4	0.57	2,610	19.1	0.38	2,371	19.4	0.41
Delay Adjusted	55,652	419.0	1.82	16,460	124.2	0.99	7,572	55.3	0.65	5,587	41.6	0.57	2,627	19.2	0.38	2,405	19.7	0.41
Maine																		
Observed	20,702	446.6	3.23	5,743	125.5	1.72	3,188	65.2	1.18	1,638	34.3	0.88	1,506	30.9	0.82	822	22.3	0.82
Delay Adjusted	20,996	452.9	3.28	5,842	127.8	1.75	3,223	66.0	1.19	1,649	34.5	0.88	1,516	31.1	0.83	827	22.4	0.82
Maryland																		
Observed	73,149	415.9	1.57	23,092	131.0	0.88	9,245	52.0	0.55	5,920	33.3	0.44	4,867	26.4	0.39	3,554	22.0	0.38
Delay Adjusted	75,055	426.8	1.61	23,569	133.7	0.90	9,456	53.1	0.56	6,038	33.9	0.45	4,918	26.7	0.39	3,632	22.5	0.39
Massachusetts																		
Observed	92,851	443.5	1.49	28,058	136.1	0.83	13,038	60.5	0.54	7,338	33.7	0.40	6,255	28.9	0.37	5,322	29.1	0.41
Delay Adjusted	94,274	450.1	1.52	28,219	136.9	0.84	13,263	61.5	0.55	7,423	34.0	0.41	6,280	29.1	0.38	5,348	29.3	0.41
Michigan																		
Observed	129,486	419.9	1.20	37,301	122.2	0.65	18,929	58.8	0.43	10,849	34.0	0.33	9,267	28.9	0.31	5,218	19.8	0.28
Delay Adjusted	131,547	426.6	1.22	37,732	123.6	0.66	19,249	59.8	0.44	11,008	34.5	0.34	9,352	29.2	0.31	5,284	20.1	0.29
Detroit																		
Observed	54,551	450.6	1.98	15,788	131.6	1.07	8,075	64.8	0.73	4,704	37.5	0.56	3,787	30.3	0.50	2,511	23.6	0.48
Delay Adjusted	55,063	454.9	2.00	15,897	132.5	1.08	8,122	65.2	0.74	4,733	37.7	0.56	3,805	30.5	0.51	2,530	23.8	0.49
Mississippi																		
Observed	35,533	402.4	2.18	10,104	115.5	1.17	5,172	56.3	0.79	3,786	41.8	0.69	1,778	19.6	0.47	1,261	15.9	0.46
Delay Adjusted	36,266	410.6	2.22	10,287	117.7	1.20	5,242	57.1	0.80	3,827	42.2	0.70	1,788	19.7	0.48	1,287	16.3	0.47
Missouri																		
Observed	78,784	421.7	1.54	23,218	125.9	0.85	12,540	64.3	0.58	7,077	36.6	0.45	4,997	26.1	0.38	2,866	17.8	0.34
Delay Adjusted	80,830	432.6	1.58	23,608	128.0	0.86	12,762	65.4	0.59	7,198	37.3	0.45	5,041	26.3	0.38	2,905	18.0	0.35
Montana																		
Observed	13,173	416.2	3.77	3,893	123.1	2.06	1,790	53.8	1.30	1,117	34.1	1.05	860	25.9	0.92	561	21.6	0.95
Delay Adjusted	13,693	432.6	3.92	4,029	127.4	2.13	1,829	55.0	1.32	1,154	35.3	1.09	881	26.5	0.94	571	22.0	0.97
Nebraska																		
Observed	22,306	410.3	2.83	6,538	121.8	1.55	2,834	50.2	0.96	2,170	38.2	0.84	1,470	26.6	0.71	975	21.0	0.69
Delay Adjusted	23,032	423.6	2.92	6,637	123.7	1.57	2,894	51.3	0.98	2,217	39.0	0.86	1,479	26.8	0.71	990	21.3	0.70

Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Delay Adjusted Counts and Rates[1] for the Five Most Commonly Diagnosed Cancers
Selected Areas in the NAACCR United States Combined[2], 2010-2014, All Races, Females

	All Sites			Breast			Lung & Bronchus			Colon & Rectum			Uterus Corpus & Uterus, NOS[3]			Thyroid		
	Count	*Rate*	*S.E.[3]*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
New Hampshire																		
Observed	19,220	456.0	3.39	5,914	140.4	1.88	2,726	63.3	1.24	1,481	34.2	0.91	1,466	32.8	0.88	845	23.3	0.83
Delay Adjusted	19,662	466.5	3.47	5,978	141.9	1.90	2,783	64.6	1.26	1,502	34.6	0.93	1,483	33.2	0.89	858	23.6	0.84
New Jersey																		
Observed	123,204	445.1	1.30	36,075	132.0	0.71	15,018	52.5	0.44	10,947	37.8	0.37	8,874	31.1	0.34	6,800	27.9	0.35
Delay Adjusted	125,364	452.9	1.32	36,560	133.8	0.72	15,204	53.1	0.44	11,064	38.3	0.37	8,946	31.4	0.34	6,906	28.3	0.35
New York																		
Observed	267,814	441.9	0.87	76,791	129.0	0.48	34,081	54.2	0.30	22,514	35.7	0.24	19,341	31.0	0.23	15,178	28.3	0.23
Delay Adjusted	270,879	446.9	0.88	77,291	129.8	0.48	34,247	54.5	0.30	22,657	36.0	0.25	19,388	31.1	0.23	15,304	28.5	0.24
North Carolina																		
Observed	122,150	414.9	1.21	37,914	129.4	0.68	17,108	56.1	0.43	9,839	33.0	0.34	7,459	24.4	0.29	4,854	18.3	0.27
Delay Adjusted	125,974	427.9	1.25	38,861	132.6	0.69	17,354	56.9	0.44	10,036	33.6	0.34	7,553	24.7	0.29	4,972	18.7	0.27
North Dakota																		
Observed	8,348	407.2	4.63	2,397	121.4	2.58	1,048	48.9	1.54	872	40.0	1.40	527	25.4	1.14	426	24.4	1.23
Delay Adjusted	8,590	419.0	4.77	2,419	122.6	2.61	1,075	50.1	1.58	898	41.2	1.44	537	25.9	1.16	438	25.1	1.26
Ohio																		
Observed	153,232	420.1	1.10	44,328	122.9	0.60	22,436	59.0	0.40	13,624	35.8	0.31	10,697	28.4	0.28	6,693	21.7	0.27
Delay Adjusted	157,817	432.6	1.14	45,333	125.6	0.61	22,969	60.4	0.41	13,920	36.6	0.32	10,850	28.8	0.29	6,837	22.1	0.28
Oklahoma																		
Observed	45,615	406.7	1.95	13,103	117.8	1.05	6,872	58.7	0.72	4,236	36.9	0.58	2,496	21.7	0.45	1,731	17.5	0.43
Delay Adjusted	46,314	413.0	1.98	13,277	119.4	1.07	6,955	59.4	0.72	4,279	37.3	0.58	2,523	22.0	0.45	1,753	17.7	0.44
Oregon																		
Observed	50,044	414.7	1.91	15,188	126.0	1.06	6,696	53.4	0.67	3,881	31.2	0.52	3,342	26.5	0.47	2,010	19.6	0.45
Delay Adjusted	50,838	421.2	1.94	15,354	127.4	1.07	6,799	54.2	0.68	3,923	31.6	0.52	3,365	26.7	0.48	2,022	19.7	0.45
Pennsylvania																		
Observed	192,175	453.8	1.07	53,851	129.8	0.58	25,221	56.1	0.36	16,964	37.5	0.30	14,004	32.1	0.28	10,515	30.5	0.31
Delay Adjusted	195,168	460.9	1.09	54,366	131.0	0.58	25,456	56.7	0.36	17,097	37.8	0.30	14,050	32.3	0.28	10,684	31.0	0.31
Rhode Island																		
Observed	15,446	452.9	3.78	4,331	130.3	2.04	2,265	63.7	1.37	1,271	34.7	1.01	1,117	31.9	0.98	784	27.1	1.00
Delay Adjusted	15,730	461.2	3.85	4,380	131.7	2.07	2,297	64.5	1.39	1,282	35.0	1.02	1,130	32.2	0.99	789	27.3	1.01

1. Rates are per 100,000 population and are age-adjusted by five-year age groups to the 2000 U.S. standard population based on single years of age.
2. See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.
3. Abbreviations: NOS - Not Otherwise Specified; S.E. - Standard Error

^ There are insufficient historical registry data to calculate delay factors necessary for delay adjusted rates.

\- Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for All Sites combined.

Note: Delay adjusted counts are rounded to the nearest whole number. A small registry and/or rare cancer site may have the same delay adjusted count and observed count, but this is just due to rounding to whole counts. For more information about cancer incidence rates adjusted for reporting delay see https://surveillance.cancer.gov/delay/.

Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Delay Adjusted Counts and Rates[1] for the Five Most Commonly Diagnosed Cancers
Selected Areas in the NAACCR United States Combined[2], 2010-2014, All Races, Females

	All Sites			Prostate			Lung & Bronchus			Colon & Rectum			Uterus Corpus & Uterus, NOS[3]			Thyroid		
	Count	*Rate*	*S.E.*[3]	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
South Carolina																		
Observed	59,570	405.6	1.70	18,597	127.2	0.96	8,267	53.7	0.60	5,051	33.8	0.49	3,505	22.7	0.39	2,222	17.2	0.38
Delay Adjusted	61,483	418.6	1.76	19,093	130.5	0.98	8,453	54.9	0.61	5,180	34.6	0.50	3,572	23.2	0.40	2,285	17.7	0.39
South Dakota																		
Observed	10,474	421.1	4.27	3,167	130.7	2.41	1,324	50.8	1.42	1,041	38.8	1.25	708	28.2	1.09	412	19.8	1.01
Delay Adjusted	10,954	440.4	4.47	3,230	133.3	2.46	1,369	52.5	1.47	1,076	40.1	1.29	717	28.5	1.11	422	20.2	1.03
Tennessee																		
Observed	81,739	413.2	1.48	23,909	121.1	0.80	12,555	61.0	0.55	7,307	36.4	0.43	4,633	22.7	0.34	3,231	18.5	0.33
Delay Adjusted	83,258	420.9	1.50	24,240	122.8	0.81	12,700	61.7	0.56	7,378	36.7	0.44	4,674	22.9	0.34	3,280	18.8	0.34
Texas																		
Observed	246,902	372.5	0.76	74,269	111.5	0.41	29,299	44.7	0.26	21,373	32.2	0.22	14,447	21.1	0.18	11,944	18.3	0.17
Delay Adjusted	253,934	383.1	0.78	75,969	114.0	0.42	29,783	45.4	0.27	21,672	32.6	0.23	14,643	21.4	0.18	12,416	19.0	0.18
Utah																		
Observed	22,861	368.8	2.47	7,012	114.7	1.39	1,448	24.1	0.64	1,686	27.7	0.68	1,643	25.9	0.65	1,856	28.8	0.68
Delay Adjusted	23,215	374.4	2.51	7,038	115.2	1.39	1,468	24.4	0.65	1,701	28.0	0.69	1,650	26.0	0.65	1,862	28.9	0.68
Vermont																		
Observed	8,932	432.6	4.76	2,662	130.1	2.62	1,285	59.7	1.71	710	33.0	1.28	710	32.8	1.27	316	19.0	1.13
Delay Adjusted	9,051	438.3	4.82	2,675	130.7	2.63	1,301	60.5	1.73	720	33.4	1.29	713	32.9	1.28	318	19.2	1.13
Virginia																		
Observed	93,909	394.8	1.31	30,271	126.9	0.74	12,254	50.8	0.47	7,934	33.1	0.38	5,915	23.8	0.31	4,277	19.6	0.31
Delay Adjusted	95,775	402.6	1.34	30,740	128.9	0.76	12,395	51.4	0.47	8,018	33.5	0.38	5,972	24.0	0.32	4,332	19.8	0.31
Washington																		
Observed	84,933	430.0	1.51	26,686	134.9	0.85	10,712	53.7	0.53	6,534	32.9	0.42	5,342	25.6	0.36	3,679	20.4	0.34
Delay Adjusted	86,026	435.5	1.53	26,838	135.6	0.85	10,855	54.4	0.54	6,600	33.2	0.42	5,371	25.8	0.36	3,709	20.6	0.35
Seattle																		
Observed	59,547	441.3	1.85	18,947	139.9	1.04	7,270	54.0	0.65	4,519	33.4	0.51	3,756	26.3	0.44	2,586	20.6	0.41
Delay Adjusted	59,901	444.0	1.86	18,968	140.1	1.04	7,289	54.1	0.65	4,530	33.4	0.51	3,771	26.4	0.44	2,596	20.7	0.42
West Virginia																		
Observed	27,296	437.9	2.75	7,112	114.8	1.42	4,416	66.2	1.01	2,664	40.9	0.81	1,935	30.4	0.72	1,226	24.3	0.72
Delay Adjusted	27,929	448.5	2.82	7,310	118.0	1.45	4,493	67.4	1.03	2,710	41.6	0.83	1,953	30.7	0.73	1,253	24.8	0.74
Wisconsin																		
Observed	74,902	428.9	1.61	21,974	127.9	0.89	9,883	54.8	0.56	6,139	33.6	0.44	5,428	29.9	0.41	2,947	19.7	0.37
Delay Adjusted	76,538	437.9	1.64	22,528	131.2	0.91	10,091	55.9	0.57	6,275	34.3	0.45	5,596	30.8	0.43	3,019	20.2	0.38

Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Delay Adjusted Counts and Rates[1] for the Five Most Commonly Diagnosed Cancers
Selected Areas in the NAACCR United States Combined[2], 2010-2014, All Races, Females

	All Sites			Prostate			Lung & Bronchus			Colon & Rectum			Uterus Corpus & Uterus, NOS[3]			Thyroid		
	Count	*Rate*	*S.E.[3]*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
Wyoming																		
Observed	6,032	377.1	5.00	1,827	113.5	2.74	725	44.1	1.67	490	30.5	1.41	357	21.5	1.18	305	21.3	1.26
Delay Adjusted	6,090	380.7	5.05	1,841	114.3	2.76	729	44.3	1.68	492	30.6	1.42	360	21.7	1.19	308	21.5	1.28

1. Rates are per 100,000 population and are age-adjusted by five-year age groups to the 2000 U.S. standard population based on single years of age.
2. See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.
3. Abbreviations: NOS - Not Otherwise Specified; S.E. - Standard Error

^ There are insufficient historical registry data to calculate delay factors necessary for delay adjusted rates.

\- Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for All Sites combined.

Note: Delay adjusted counts are rounded to the nearest whole number. A small registry and/or rare cancer site may have the same delay adjusted count and observed count, but this is just due to rounding to whole counts. For more information about cancer incidence rates adjusted for reporting delay see https://surveillance.cancer.gov/delay/.

NAACCR U.S. Combined Delay Adjusted Incidence 2010-2014, White Combined

UNITED STATES

V-15 Combined Delay Adjusted Incidence – White

Average Annual Age-adjusted Cancer Incidence Delay Adjusted Counts and Rates*, Selected Sites
NAACCR United States Combined[1], 2010-2014, White

	MALES (ALL RACES)						FEMALES (ALL RACES)					
	Observed			Delay Adjusted			Observed			Delay Adjusted		
SITES	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
All Sites (invasive & in situ bladder)	3,202,538	487.1	0.28	3,285,488	499.8	0.28	3,116,727	417.2	0.24	3,197,799	427.9	0.25
Oral Cavity & Pharynx	122,398	17.9	0.05	125,579	18.3	0.05	49,388	6.5	0.03	50,631	6.7	0.03
Digestive												
Esophagus	55,310	8.3	0.04	56,148	8.4	0.04	13,892	1.7	0.02	14,100	1.8	0.02
Stomach	53,959	8.4	0.04	54,766	8.5	0.04	30,863	4.0	0.02	31,319	4.1	0.02
Colon & Rectum	289,746	44.8	0.08	294,604	45.5	0.09	264,042	34.0	0.07	268,387	34.5	0.07
Liver & IBD[2]	76,330	10.9	0.04	78,153	11.2	0.04	29,910	3.8	0.02	30,635	3.9	0.02
Pancreas	92,145	14.1	0.05	93,504	14.3	0.05	85,916	10.7	0.04	87,157	10.9	0.04
Respiratory												
Larynx	40,378	5.9	0.03	41,123	6.0	0.03	10,553	1.4	0.01	10,747	1.4	0.01
Lung & bronchus	472,731	72.7	0.11	479,536	73.8	0.11	426,824	54.4	0.08	432,990	55.2	0.09
Skin												
Melanomas of the skin	190,947	29.7	0.07	199,002	31.0	0.07	132,012	18.8	0.05	137,396	19.6	0.06
Breast (female cases only)	-	-	-	-	-	-	919,063	124.5	0.13	935,962	126.8	0.14
Female Genital												
Cervix uteri	-	-	-	-	-	-	46,054	7.3	0.04	46,862	7.5	0.04
Corpus & uterus, NOS[2]	-	-	-	-	-	-	201,669	26.2	0.06	204,493	26.6	0.06
Ovary	-	-	-	-	-	-	88,008	11.8	0.04	90,333	12.1	0.04
Male Genital												
Prostate	735,242	105.5	0.13	766,456	110.0	0.13	-	-	-	-	-	-
Testis	36,726	6.4	0.03	37,351	6.5	0.03	-	-	-	-	-	-
Urinary												
Bladder (incl. *in situ*)	240,261	38.0	0.08	246,759	39.1	0.08	74,103	9.3	0.03	76,092	9.5	0.04
Kidney & renal pelvis	145,125	21.9	0.06	148,838	22.5	0.06	85,287	11.4	0.04	87,452	11.7	0.04
Brain & Other Nervous System	52,277	8.3	0.04	53,600	8.5	0.04	41,399	6.0	0.03	42,427	6.1	0.03
Endocrine												
Thyroid gland	48,427	7.6	0.04	49,350	7.7	0.04	140,681	22.1	0.06	143,339	22.6	0.06
Lymphomas												
Hodgkin Lymphoma	18,811	3.1	0.02	19,189	3.2	0.02	15,072	2.5	0.02	15,369	2.5	0.02
Non-Hodgkin Lymphoma	150,400	23.5	0.06	155,491	24.3	0.06	124,006	16.2	0.05	128,170	16.8	0.05
Multiple Myeloma	47,663	7.4	0.03	51,425	7.9	0.04	35,869	4.6	0.02	38,701	4.9	0.03
Leukemias	112,076	17.9	0.05	123,554	19.7	0.06	80,246	10.9	0.04	88,363	12.0	0.04

* Rates are per 100,000 population and are age-adjusted by five-year age groups to the 2000 U.S. standard population based on single years of age and to standard.

\- Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for All Sites combined.

[1] NAACCR United States Combined includes registries for which a U.S. average delay adjustment factor was calculated. See technical notes.

[2] Abbreviations: IBD – Intrahepatic Bile Duct; NOS - Not Otherwise Specified

Note: Delay adjusted counts are rounded to the nearest whole number. A small registry and/or rare cancer site may have the same delay adjusted count and observed count, but this is just due to rounding to whole counts. For more information about cancer incidence rates adjusted for reporting delay see https://surveillance.cancer.gov/delay/.

Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Delay Adjusted Counts and Rates[1] for the Five Most Commonly Diagnosed Cancers
Selected Areas in the NAACCR United States Combined[2], 2010-2014, White, Males

	All Sites			Prostate			Lung & Bronchus			Colon & Rectum			Bladder			Melanoma of the Skin		
	Count	*Rate*	*S.E.*[3]	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
NAACCR U.S. Combined																		
Observed	3,202,538	487.1	0.28	735,242	105.5	0.13	472,731	72.7	0.11	289,746	44.8	0.08	240,261	38.0	0.08	190,947	29.7	0.07
Delay Adjusted	3,285,488	499.8	0.28	766,456	110.0	0.13	479,536	73.8	0.11	294,604	45.5	0.09	246,759	39.1	0.08	199,002	31.0	0.07
Alabama																		
Observed	51,066	511.6	2.32	11,431	107.3	1.03	9,278	92.1	0.98	4,866	49.6	0.73	3,577	37.2	0.64	3,275	34.1	0.61
Delay Adjusted	51,834	519.2	2.36	11,757	110.5	1.06	9,330	92.7	0.98	4,908	50.0	0.74	3,621	37.6	0.65	3,364	35.0	0.63
Alaska																		
Observed	5,350	433.8	6.72	1,288	94.3	2.97	692	61.1	2.62	461	39.0	2.07	394	36.8	2.07	239	18.8	1.34
Delay Adjusted	5,565	451.5	6.99	1,363	99.8	3.16	710	62.7	2.69	474	40.1	2.14	402	37.5	2.12	263	20.7	1.48
Arizona																		
Observed	65,306	405.1	1.61	13,279	77.8	0.68	9,272	56.9	0.60	6,171	38.9	0.50	5,345	33.4	0.46	4,112	26.1	0.41
Delay Adjusted	68,673	426.0	1.69	14,363	84.2	0.74	9,493	58.2	0.61	6,478	40.8	0.53	5,599	35.0	0.48	4,468	28.3	0.45
Arkansas																		
Observed	35,169	502.2	2.73	7,998	107.7	1.22	6,865	96.7	1.19	3,339	48.5	0.86	2,544	37.4	0.75	1,832	27.3	0.65
Delay Adjusted	35,568	507.9	2.76	8,155	109.8	1.25	6,903	97.2	1.19	3,359	48.8	0.86	2,575	37.8	0.76	1,881	28.0	0.67
California																		
Observed	319,275	460.5	0.83	76,018	105.4	0.39	33,459	50.8	0.28	28,912	42.0	0.25	22,430	34.5	0.23	23,188	33.8	0.23
Delay Adjusted	324,008	467.4	0.85	78,265	108.7	0.40	33,751	51.2	0.29	29,090	42.2	0.25	22,755	35.0	0.24	23,837	34.8	0.23
Greater California																		
Observed	193,765	465.1	1.08	45,347	104.3	0.50	21,843	54.6	0.38	17,503	42.3	0.33	13,706	34.9	0.30	14,427	35.1	0.30
Delay Adjusted	197,272	473.6	1.10	46,903	108.0	0.52	22,057	55.2	0.38	17,638	42.6	0.33	13,961	35.6	0.31	14,891	36.2	0.31
Greater Bay																		
Observed	57,303	485.2	2.08	14,812	119.4	1.01	5,124	46.2	0.66	4,657	39.7	0.60	3,988	35.9	0.58	5,104	43.8	0.63
Delay Adjusted	57,592	487.7	2.09	14,933	120.4	1.02	5,149	46.4	0.66	4,668	39.8	0.60	4,011	36.1	0.58	5,214	44.8	0.64
Los Angeles																		
Observed	68,063	429.1	1.69	15,809	97.6	0.80	6,475	43.7	0.56	6,743	42.7	0.53	4,731	32.3	0.48	3,653	23.2	0.39
Delay Adjusted	68,993	435.0	1.71	16,374	101.2	0.83	6,527	44.1	0.56	6,775	42.9	0.54	4,778	32.6	0.48	3,728	23.7	0.40
Colorado																		
Observed	48,982	425.3	2.00	12,129	98.6	0.93	5,183	47.9	0.69	4,217	37.6	0.60	3,447	32.6	0.58	3,391	29.6	0.53
Delay Adjusted	50,364	437.3	2.05	12,990	106.0	1.01	5,226	48.3	0.70	4,267	38.0	0.61	3,578	33.8	0.60	3,486	30.5	0.54
Connecticut																		
Observed	44,169	513.0	2.50	10,261	111.7	1.13	5,968	69.9	0.92	3,709	43.8	0.74	4,125	48.9	0.78	2,456	29.3	0.61
Delay Adjusted	45,244	525.5	2.56	10,578	115.3	1.17	6,050	70.8	0.94	3,757	44.3	0.74	4,207	49.9	0.79	2,522	30.1	0.62

Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Delay Adjusted Counts and Rates[1] for the Five Most Commonly Diagnosed Cancers
Selected Areas in the NAACCR United States Combined[2], 2010-2014, White, Males

	All Sites			Prostate			Lung & Bronchus			Colon & Rectum			Bladder			Melanoma of the Skin		
	Count	*Rate*	*S.E.*[3]	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
Delaware																		
Observed	11,491	559.4	5.34	2,837	128.1	2.45	1,699	82.2	2.03	863	43.0	1.49	924	45.8	1.53	959	47.9	1.59
Delay Adjusted	11,852	577.0	5.51	2,972	134.2	2.56	1,737	84.1	2.08	876	43.6	1.52	956	47.4	1.59	993	49.6	1.64
District of Columbia																		
Observed	2,069	375.6	8.69	539	97.6	4.35	173	35.6	2.81	148	26.6	2.30	124	24.4	2.28	122	22.2	2.12
Delay Adjusted	2,180	395.9	9.16	559	101.3	4.52	177	36.4	2.87	153	27.4	2.37	127	25.1	2.34	128	23.3	2.24
Florida																		
Observed	243,510	464.4	0.96	52,312	94.6	0.42	39,119	72.1	0.37	21,501	41.3	0.29	18,793	34.7	0.26	16,379	32.0	0.26
Delay Adjusted	253,149	482.9	1.00	55,406	100.1	0.44	39,415	72.7	0.37	22,115	42.5	0.30	19,508	36.1	0.27	17,580	34.3	0.27
Georgia																		
Observed	83,934	515.2	1.84	18,972	109.0	0.82	13,581	85.4	0.76	7,439	46.2	0.55	5,651	37.2	0.51	7,091	44.2	0.54
Delay Adjusted	85,594	525.4	1.88	19,423	111.7	0.84	13,736	86.4	0.77	7,527	46.7	0.56	5,757	37.9	0.52	7,335	45.8	0.56
Hawaii																		
Observed	6,037	511.5	7.01	1,145	88.4	2.78	635	54.2	2.28	502	42.1	2.00	422	38.3	1.96	875	77.1	2.76
Delay Adjusted	6,191	524.5	7.19	1,176	90.8	2.85	641	54.7	2.30	506	42.4	2.01	429	38.9	2.00	910	80.3	2.88
Idaho																		
Observed	18,409	460.7	3.48	4,820	114.2	1.68	2,159	55.4	1.22	1,547	39.2	1.02	1,437	37.7	1.02	1,303	33.0	0.94
Delay Adjusted	18,915	473.3	3.57	5,068	120.1	1.77	2,204	56.5	1.24	1,570	39.7	1.04	1,518	39.8	1.07	1,354	34.2	0.97
Illinois																		
Observed	133,928	506.2	1.41	30,909	110.5	0.64	20,199	78.2	0.56	13,241	50.7	0.45	10,174	40.3	0.41	6,768	26.0	0.32
Delay Adjusted	136,495	515.9	1.44	32,041	114.7	0.67	20,424	79.1	0.57	13,364	51.2	0.45	10,395	41.2	0.42	7,012	26.9	0.33
Indiana																		
Observed	73,068	480.9	1.82	14,560	90.3	0.77	13,369	88.9	0.79	7,241	48.4	0.58	5,443	37.4	0.52	3,328	22.3	0.40
Delay Adjusted	75,803	498.9	1.89	15,410	95.8	0.82	13,668	90.9	0.81	7,446	49.8	0.60	5,709	39.2	0.54	3,579	24.0	0.43
Iowa																		
Observed	42,372	513.9	2.54	9,613	109.9	1.14	6,437	78.1	0.99	4,177	51.6	0.81	3,114	38.3	0.70	2,325	29.2	0.62
Delay Adjusted	43,185	523.8	2.59	9,904	113.3	1.17	6,520	79.1	1.00	4,213	52.0	0.82	3,182	39.2	0.71	2,392	30.0	0.64
Kansas																		
Observed	34,074	497.5	2.75	8,261	115.3	1.29	4,864	71.7	1.05	3,125	46.3	0.84	2,608	39.1	0.78	1,979	29.8	0.68
Delay Adjusted	^	^	^	^	^	^	^	^	^	^	^	^	^	^	^	^	^	^

1. Rates are per 100,000 population and are age-adjusted by five-year age groups to the 2000 U.S. standard population based on single years of age.
2. See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.
3. Abbreviations: S.E. - Standard Error

^ There are insufficient historical registry data to calculate delay factors necessary for delay adjusted rates.

\- Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for All Sites combined.

Note: Delay adjusted counts are rounded to the nearest whole number. A small registry and/or rare cancer site may have the same delay adjusted count and observed count, but this is just due to rounding to whole counts. For more information about cancer incidence rates adjusted for reporting delay see https://surveillance.cancer.gov/delay/.

Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Delay Adjusted Counts and Rates[1] for the Five Most Commonly Diagnosed Cancers
Selected Areas in the NAACCR United States Combined[2], 2010-2014, White, Males

	All Sites			Prostate			Lung & Bronchus			Colon & Rectum			Bladder			Melanoma of the Skin		
	Count	*Rate*	*S.E.*[3]	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
Kentucky																		
Observed	61,114	573.7	2.39	11,986	105.6	0.99	12,431	116.9	1.08	6,202	59.1	0.77	4,125	41.1	0.66	3,339	32.4	0.58
Delay Adjusted	62,498	586.7	2.45	12,566	110.5	1.04	12,576	118.2	1.09	6,317	60.2	0.79	4,234	42.1	0.68	3,434	33.2	0.59
Louisiana																		
Observed	45,570	551.4	2.65	11,177	127.1	1.23	6,992	85.5	1.05	4,295	52.5	0.82	3,017	38.1	0.71	2,436	30.6	0.64
Delay Adjusted	46,023	556.9	2.67	11,294	128.5	1.24	7,034	86.0	1.05	4,311	52.7	0.82	3,046	38.5	0.72	2,480	31.2	0.65
Maine																		
Observed	20,441	504.5	3.64	4,345	98.1	1.53	3,441	84.7	1.48	1,694	42.8	1.07	1,869	47.6	1.12	1,106	28.0	0.87
Delay Adjusted	20,790	513.1	3.70	4,461	100.8	1.57	3,480	85.6	1.50	1,707	43.1	1.08	1,903	48.5	1.15	1,139	28.8	0.89
Maryland																		
Observed	49,947	490.1	2.25	12,310	112.6	1.04	6,628	66.6	0.84	4,077	40.8	0.65	4,070	41.8	0.67	3,911	39.0	0.64
Delay Adjusted	51,219	502.7	2.31	12,830	117.6	1.09	6,785	68.1	0.86	4,137	41.4	0.66	4,252	43.7	0.70	3,945	39.3	0.64
Massachusetts																		
Observed	77,040	490.8	1.81	17,890	106.0	0.81	10,934	71.1	0.69	6,549	42.6	0.54	6,383	42.1	0.54	4,072	26.6	0.43
Delay Adjusted	78,477	499.9	1.84	18,459	109.4	0.84	11,138	72.4	0.71	6,638	43.1	0.54	6,498	42.9	0.55	4,253	27.8	0.45
Michigan																		
Observed	113,657	486.8	1.48	27,917	111.6	0.68	17,452	75.2	0.58	9,560	42.0	0.44	9,118	40.6	0.43	5,473	24.3	0.34
Delay Adjusted	115,929	496.6	1.51	28,655	114.6	0.70	17,765	76.5	0.59	9,720	42.7	0.45	9,251	41.2	0.44	5,632	24.9	0.35
Detroit																		
Observed	42,250	559.4	2.80	11,099	138.6	1.35	6,095	82.5	1.08	3,393	45.6	0.80	3,468	47.5	0.83	1,978	26.8	0.62
Delay Adjusted	42,806	566.8	2.83	11,366	141.9	1.39	6,133	83.0	1.09	3,419	46.0	0.81	3,530	48.4	0.84	1,996	27.1	0.62
Mississippi																		
Observed	28,196	530.2	3.24	6,278	110.4	1.42	5,169	96.9	1.38	2,787	53.2	1.03	1,826	35.6	0.85	1,633	31.8	0.81
Delay Adjusted	28,867	542.8	3.31	6,494	114.2	1.47	5,244	98.3	1.40	2,823	53.8	1.05	1,892	36.9	0.89	1,663	32.3	0.82
Missouri																		
Observed	70,749	486.2	1.86	14,865	95.4	0.80	12,713	87.7	0.79	6,934	48.8	0.60	4,929	35.1	0.51	3,766	26.6	0.44
Delay Adjusted	72,582	498.8	1.91	15,517	99.7	0.83	12,936	89.3	0.81	7,053	49.6	0.61	5,065	36.1	0.52	3,881	27.4	0.46
Montana																		
Observed	13,548	465.3	4.13	3,598	113.5	1.94	1,679	58.8	1.47	1,248	43.1	1.26	1,017	35.9	1.15	853	30.2	1.08
Delay Adjusted	14,129	485.1	4.31	3,812	120.3	2.06	1,715	60.0	1.50	1,291	44.6	1.30	1,071	37.8	1.21	904	32.0	1.14
Nebraska																		
Observed	22,443	491.8	3.34	5,514	115.5	1.58	3,199	71.1	1.28	2,223	49.0	1.06	1,661	37.7	0.94	1,055	23.3	0.73
Delay Adjusted	23,297	510.5	3.47	5,898	123.5	1.69	3,267	72.6	1.30	2,277	50.2	1.08	1,749	39.7	0.99	1,135	25.1	0.79

Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Delay Adjusted Counts and Rates[1] for the Five Most Commonly Diagnosed Cancers
Selected Areas in the NAACCR United States Combined[2], 2010-2014, White, Males

	All Sites			Prostate			Lung & Bronchus			Colon & Rectum			Bladder			Melanoma of the Skin		
	Count	*Rate*	*S.E.*[3]	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
New Hampshire																		
Observed	18,895	514.7	3.88	4,909	121.6	1.79	2,538	71.5	1.46	1,461	40.7	1.10	1,681	48.2	1.21	1,149	32.3	0.99
Delay Adjusted	19,334	526.7	3.97	5,041	124.9	1.84	2,589	73.0	1.49	1,482	41.3	1.12	1,711	49.1	1.23	1,188	33.4	1.02
New Jersey																		
Observed	99,033	536.5	1.74	25,269	130.0	0.83	12,200	67.2	0.62	8,838	48.4	0.52	8,005	44.8	0.51	6,053	33.2	0.44
Delay Adjusted	101,860	551.8	1.79	26,581	136.9	0.88	12,358	68.1	0.63	8,983	49.1	0.53	8,295	46.5	0.53	6,315	34.7	0.45
New York																		
Observed	210,728	539.6	1.20	51,944	126.4	0.56	27,822	71.9	0.44	17,872	46.5	0.35	17,475	46.0	0.35	10,686	27.9	0.27
Delay Adjusted	215,058	550.7	1.22	53,564	130.4	0.58	28,008	72.4	0.44	18,094	47.1	0.36	17,848	47.0	0.36	11,317	29.5	0.29
North Carolina																		
Observed	99,314	503.3	1.64	22,783	107.9	0.73	16,910	86.3	0.68	7,997	41.4	0.47	7,216	38.2	0.46	7,191	37.6	0.45
Delay Adjusted	102,237	518.1	1.68	23,872	113.1	0.77	17,184	87.7	0.69	8,145	42.2	0.48	7,417	39.2	0.47	7,464	39.0	0.47
North Dakota																		
Observed	9,018	496.0	5.34	2,357	123.3	2.59	1,238	69.3	2.00	949	53.5	1.78	667	37.3	1.47	461	25.9	1.24
Delay Adjusted	9,275	510.1	5.49	2,437	127.4	2.68	1,268	70.9	2.05	975	54.9	1.82	694	38.8	1.53	473	26.6	1.27
Ohio																		
Observed	134,500	484.6	1.35	29,179	98.3	0.59	22,969	83.0	0.56	12,822	47.1	0.43	10,594	39.5	0.39	7,012	26.1	0.32
Delay Adjusted	139,734	503.7	1.41	30,992	104.4	0.62	23,549	85.0	0.57	13,189	48.5	0.44	11,063	41.3	0.41	7,429	27.7	0.34
Oklahoma																		
Observed	40,116	472.8	2.41	8,690	97.1	1.06	7,147	83.8	1.01	3,827	45.9	0.76	2,839	34.4	0.66	2,071	25.1	0.56
Delay Adjusted	40,612	478.6	2.44	8,846	98.8	1.08	7,220	84.7	1.02	3,848	46.1	0.76	2,865	34.7	0.66	2,080	25.2	0.57
Oregon																		
Observed	47,047	458.7	2.17	10,983	99.5	0.97	6,278	62.3	0.80	4,003	40.2	0.65	3,765	38.1	0.64	3,267	32.6	0.59
Delay Adjusted	48,106	469.1	2.22	11,302	102.4	1.00	6,396	63.4	0.82	4,083	41.0	0.66	3,860	39.1	0.65	3,317	33.1	0.60
Pennsylvania																		
Observed	168,600	518.5	1.29	36,818	106.0	0.56	25,242	77.2	0.49	15,804	49.3	0.40	14,561	45.4	0.38	9,156	28.9	0.31
Delay Adjusted	171,754	528.2	1.31	38,094	109.7	0.58	25,521	78.1	0.50	15,964	49.8	0.40	14,802	46.2	0.39	9,453	29.9	0.32
Rhode Island																		
Observed	13,282	510.0	4.53	2,837	102.3	1.97	2,029	78.8	1.79	1,052	41.1	1.29	1,199	46.7	1.38	753	29.7	1.11
Delay Adjusted	13,563	520.8	4.63	2,954	106.5	2.05	2,058	79.9	1.81	1,062	41.5	1.31	1,224	47.7	1.41	770	30.4	1.14

1. Rates are per 100,000 population and are age-adjusted by five-year age groups to the 2000 U.S. standard population based on single years of age.
2. See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.
3. Abbreviations: S.E. - Standard Error

^ There are insufficient historical registry data to calculate delay factors necessary for delay adjusted rates.

\- Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for All Sites combined.

Note: Delay adjusted counts are rounded to the nearest whole number. A small registry and/or rare cancer site may have the same delay adjusted count and observed count, but this is just due to rounding to whole counts. For more information about cancer incidence rates adjusted for reporting delay see https://surveillance.cancer.gov/delay/.

Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Delay Adjusted Counts and Rates[1] for the Five Most Commonly Diagnosed Cancers
Selected Areas in the NAACCR United States Combined[2], 2010-2014, White, Males

	All Sites			Prostate			Lung & Bronchus			Colon & Rectum			Bladder			Melanoma of the Skin		
	Count	*Rate*	*S.E.*[3]	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
South Carolina																		
Observed	48,278	498.3	2.34	10,631	101.1	1.01	8,285	84.8	0.96	3,996	42.2	0.69	3,626	39.4	0.67	3,393	35.9	0.64
Delay Adjusted	50,013	516.2	2.43	11,138	105.9	1.06	8,480	86.8	0.98	4,104	43.3	0.71	3,756	40.8	0.70	3,522	37.2	0.66
South Dakota																		
Observed	10,532	489.1	4.87	2,680	117.4	2.31	1,459	67.9	1.80	1,079	50.7	1.58	756	36.0	1.33	553	26.4	1.15
Delay Adjusted	11,023	511.7	5.09	2,867	125.6	2.47	1,509	70.2	1.87	1,117	52.5	1.63	815	38.8	1.43	592	28.3	1.24
Tennessee																		
Observed	75,841	515.5	1.93	17,485	110.8	0.86	14,007	95.2	0.83	6,513	45.5	0.58	5,152	36.6	0.53	4,166	29.4	0.47
Delay Adjusted	77,335	525.6	1.96	17,903	113.4	0.88	14,165	96.3	0.84	6,574	45.9	0.59	5,248	37.3	0.54	4,308	30.4	0.49
Texas																		
Observed	218,983	451.5	0.99	47,193	93.8	0.44	30,881	66.8	0.39	21,976	45.6	0.32	12,834	28.8	0.26	9,431	19.7	0.21
Delay Adjusted	227,929	469.9	1.03	50,914	101.5	0.48	31,484	68.1	0.40	22,435	46.5	0.32	13,368	30.0	0.27	10,281	21.5	0.23
Utah																		
Observed	24,050	450.0	2.97	7,076	130.0	1.58	1,631	32.3	0.82	1,840	35.2	0.84	1,492	30.5	0.80	2,585	48.8	0.98
Delay Adjusted	24,428	457.1	3.02	7,253	133.2	1.62	1,652	32.7	0.83	1,855	35.5	0.85	1,516	30.9	0.82	2,622	49.6	1.00
Vermont																		
Observed	8,799	482.8	5.34	1,940	97.2	2.28	1,291	71.0	2.04	706	39.1	1.53	685	39.0	1.53	702	39.4	1.55
Delay Adjusted	8,910	488.8	5.40	1,966	98.5	2.31	1,304	71.7	2.06	715	39.6	1.55	698	39.8	1.56	713	40.0	1.57
Virginia																		
Observed	70,009	438.6	1.70	15,834	92.1	0.75	11,221	71.8	0.70	6,215	39.4	0.51	4,984	33.3	0.48	4,263	27.3	0.43
Delay Adjusted	71,305	446.7	1.74	16,174	94.1	0.77	11,346	72.6	0.70	6,269	39.8	0.52	5,064	33.8	0.49	4,355	27.9	0.44
Washington																		
Observed	77,589	492.3	1.82	19,294	114.4	0.85	10,010	65.8	0.68	6,233	40.1	0.52	5,771	38.8	0.52	5,115	32.8	0.47
Delay Adjusted	79,577	505.0	1.87	20,013	118.7	0.88	10,180	66.9	0.69	6,374	41.0	0.53	5,975	40.2	0.54	5,233	33.6	0.48
Seattle																		
Observed	53,513	521.3	2.32	13,649	124.6	1.10	6,542	66.6	0.85	4,175	41.0	0.65	3,947	41.1	0.67	3,854	37.9	0.63
Delay Adjusted	53,974	525.8	2.34	13,845	126.4	1.12	6,553	66.7	0.85	4,199	41.3	0.66	3,989	41.5	0.68	3,912	38.5	0.64
West Virginia																		
Observed	28,003	514.3	3.17	5,725	97.7	1.32	5,471	99.0	1.38	2,820	52.8	1.02	2,134	40.5	0.90	1,339	26.0	0.73
Delay Adjusted	28,641	526.0	3.24	5,852	99.8	1.35	5,564	100.7	1.40	2,867	53.7	1.04	2,177	41.3	0.92	1,379	26.8	0.76
Wisconsin																		
Observed	72,289	491.6	1.87	17,573	111.8	0.86	9,790	67.2	0.69	6,101	42.4	0.55	5,738	40.6	0.55	3,912	27.1	0.44
Delay Adjusted	74,618	507.2	1.93	18,863	120.0	0.93	10,119	69.6	0.72	6,299	43.7	0.57	5,881	41.6	0.56	4,123	28.6	0.47

Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Delay Adjusted Counts and Rates[1] for the Five Most Commonly Diagnosed Cancers
Selected Areas in the NAACCR United States Combined[2], 2010-2014, White, Males

	All Sites			Prostate			Lung & Bronchus			Colon & Rectum			Bladder			Melanoma of the Skin		
	Count	*Rate*	*S.E.*[3]	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
Wyoming																		
Observed	6,718	446.2	5.65	1,790	110.1	2.70	722	49.8	1.92	614	41.8	1.75	520	36.8	1.67	423	29.4	1.48
Delay Adjusted	6,794	451.2	5.72	1,808	111.3	2.73	726	50.1	1.93	618	42.2	1.76	524	37.1	1.68	437	30.3	1.53

1. Rates are per 100,000 population and are age-adjusted by five-year age groups to the 2000 U.S. standard population based on single years of age.
2. See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.
3. Abbreviations: S.E. - Standard Error

^ There are insufficient historical registry data to calculate delay factors necessary for delay adjusted rates.

\- Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for All Sites combined.

Note: Delay adjusted counts are rounded to the nearest whole number. A small registry and/or rare cancer site may have the same delay adjusted count and observed count, but this is just due to rounding to whole counts. For more information about cancer incidence rates adjusted for reporting delay see https://surveillance.cancer.gov/delay/.

Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Delay Adjusted Counts and Rates[1] for the Five Most Commonly Diagnosed Cancers
Selected Areas in the NAACCR United States Combined[2], 2010-2014, White, Females

	All Sites			Breast			Lung & Bronchus			Colon & Rectum			Uterus Corpus & Uterus, NOS[3]			Thyroid		
	Count	*Rate*	*S.E.*[3]	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
NAACCR U.S. Combined																		
Observed	3,116,727	417.2	0.24	919,063	124.5	0.13	426,824	54.4	0.08	264,042	34.0	0.07	201,669	26.2	0.06	140,681	22.1	0.06
Delay Adjusted	3,197,799	427.9	0.25	935,962	126.8	0.14	432,990	55.2	0.09	268,387	34.5	0.07	204,493	26.6	0.06	143,339	22.6	0.06
Alabama																		
Observed	44,900	395.8	1.93	13,184	117.1	1.05	6,850	56.3	0.69	4,052	34.7	0.56	2,187	18.8	0.41	1,331	14.2	0.40
Delay Adjusted	45,586	401.8	1.96	13,395	119.0	1.07	6,888	56.6	0.69	4,087	35.1	0.57	2,202	18.9	0.42	1,341	14.3	0.41
Alaska																		
Observed	4,770	398.0	6.13	1,558	124.9	3.37	573	52.6	2.35	392	33.3	1.79	348	26.4	1.51	200	15.6	1.14
Delay Adjusted	4,964	414.2	6.38	1,594	127.7	3.45	587	53.9	2.41	403	34.2	1.84	360	27.3	1.56	205	16.0	1.16
Arizona																		
Observed	65,048	372.9	1.50	19,729	114.4	0.84	8,714	46.4	0.50	5,372	29.7	0.42	3,778	21.1	0.35	3,520	24.2	0.42
Delay Adjusted	68,411	392.2	1.58	20,761	120.4	0.88	8,923	47.5	0.52	5,637	31.2	0.44	3,967	22.2	0.37	3,703	25.5	0.44
Arkansas																		
Observed	30,297	389.3	2.31	8,495	110.1	1.23	5,180	62.0	0.88	2,858	35.4	0.68	1,687	21.5	0.54	1,014	15.9	0.52
Delay Adjusted	30,646	393.8	2.33	8,571	111.1	1.25	5,211	62.4	0.88	2,875	35.6	0.68	1,695	21.6	0.54	1,032	16.2	0.53
California																		
Observed	316,198	398.5	0.72	98,648	125.3	0.41	33,783	42.0	0.23	26,344	32.4	0.20	20,725	25.5	0.18	14,394	19.7	0.17
Delay Adjusted	320,840	404.3	0.73	99,089	125.8	0.41	34,084	42.3	0.23	26,502	32.6	0.20	20,772	25.6	0.18	14,464	19.8	0.17
Greater California																		
Observed	189,554	401.5	0.94	59,165	126.2	0.53	21,944	45.5	0.31	15,712	32.5	0.26	11,829	24.4	0.23	8,449	19.8	0.22
Delay Adjusted	192,938	408.6	0.96	59,527	127.0	0.53	22,166	46.0	0.31	15,831	32.8	0.27	11,859	24.5	0.23	8,489	19.9	0.22
Greater Bay																		
Observed	55,458	416.8	1.81	17,509	133.3	1.03	5,664	41.6	0.57	4,489	32.7	0.50	3,748	27.2	0.45	2,112	17.6	0.39
Delay Adjusted	55,749	419.0	1.82	17,529	133.5	1.03	5,693	41.8	0.57	4,499	32.7	0.50	3,750	27.2	0.45	2,117	17.7	0.39
Los Angeles																		
Observed	71,087	377.3	1.43	21,949	117.1	0.80	6,165	33.0	0.43	6,134	31.9	0.41	5,148	26.9	0.38	3,832	20.8	0.34
Delay Adjusted	72,049	382.3	1.45	22,008	117.4	0.80	6,215	33.3	0.43	6,163	32.0	0.42	5,162	27.0	0.38	3,857	21.0	0.34
Colorado																		
Observed	50,288	388.7	1.77	16,336	124.9	1.00	5,397	41.9	0.58	3,947	30.7	0.50	3,100	22.5	0.41	2,884	23.8	0.45
Delay Adjusted	51,675	399.3	1.82	16,525	126.3	1.01	5,442	42.2	0.59	3,994	31.0	0.50	3,130	22.8	0.42	2,927	24.1	0.46
Connecticut																		
Observed	46,293	457.8	2.22	14,030	141.8	1.24	6,254	58.6	0.76	3,662	33.8	0.58	3,108	29.5	0.54	2,427	29.8	0.63
Delay Adjusted	47,427	469.0	2.27	14,279	144.3	1.26	6,341	59.4	0.77	3,711	34.2	0.59	3,139	29.8	0.55	2,462	30.2	0.64

Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Delay Adjusted Counts and Rates[1] for the Five Most Commonly Diagnosed Cancers
Selected Areas in the NAACCR United States Combined[2], 2010-2014, White, Females

	All Sites			Breast			Lung & Bronchus			Colon & Rectum			Uterus Corpus & Uterus, NOS[3]			Thyroid		
	Count	*Rate*	*S.E.[3]*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
Delaware																		
Observed	10,670	462.1	4.67	3,091	134.4	2.52	1,626	65.3	1.65	788	32.6	1.20	714	29.4	1.14	444	24.9	1.24
Delay Adjusted	11,008	476.7	4.81	3,135	136.3	2.56	1,663	66.8	1.69	800	33.1	1.22	722	29.7	1.15	456	25.6	1.27
District of Columbia																		
Observed	2,193	374.6	8.42	816	143.7	5.22	191	34.9	2.63	145	25.7	2.24	130	22.4	2.03	165	23.1	2.00
Delay Adjusted	2,312	395.0	8.88	861	151.5	5.51	195	35.7	2.69	149	26.5	2.31	135	23.2	2.10	173	24.2	2.10
Florida																		
Observed	225,114	393.4	0.87	64,149	115.6	0.48	35,342	55.8	0.30	19,377	31.5	0.24	13,217	22.9	0.21	7,948	18.6	0.22
Delay Adjusted	234,214	409.6	0.91	66,711	120.2	0.50	35,609	56.2	0.31	19,910	32.4	0.24	13,640	23.6	0.21	8,001	18.8	0.22
Georgia																		
Observed	77,186	415.0	1.52	23,143	123.9	0.83	10,946	56.7	0.55	6,273	33.3	0.43	4,112	21.2	0.34	3,212	19.2	0.35
Delay Adjusted	78,692	423.0	1.55	23,405	125.3	0.84	11,070	57.4	0.56	6,345	33.7	0.43	4,149	21.3	0.34	3,244	19.4	0.35
Hawaii																		
Observed	4,776	454.9	7.03	1,523	145.2	3.98	488	44.8	2.13	380	36.0	1.96	275	24.0	1.54	184	20.7	1.62
Delay Adjusted	4,901	466.7	7.21	1,533	146.1	4.01	493	45.2	2.15	383	36.3	1.98	279	24.3	1.56	188	21.2	1.65
Idaho																		
Observed	17,145	402.5	3.15	5,131	120.5	1.72	2,074	47.0	1.05	1,377	32.1	0.88	1,068	24.1	0.75	934	24.8	0.83
Delay Adjusted	17,626	413.8	3.23	5,205	122.2	1.75	2,116	47.9	1.07	1,397	32.5	0.90	1,091	24.6	0.77	940	24.9	0.84
Illinois																		
Observed	134,923	436.7	1.22	39,801	130.9	0.67	18,722	58.6	0.44	11,910	36.7	0.34	9,343	29.5	0.31	6,171	23.1	0.30
Delay Adjusted	137,525	445.1	1.24	40,251	132.4	0.68	18,934	59.3	0.44	12,020	37.0	0.35	9,446	29.8	0.31	6,243	23.4	0.31
Indiana																		
Observed	74,010	420.7	1.59	20,890	119.8	0.85	11,212	61.3	0.59	6,930	38.1	0.47	4,983	27.5	0.40	2,778	18.6	0.36
Delay Adjusted	76,814	436.7	1.65	21,528	123.5	0.88	11,467	62.6	0.60	7,123	39.1	0.48	5,094	28.1	0.41	2,846	19.1	0.37
Iowa																		
Observed	40,337	433.1	2.25	11,233	123.8	1.22	5,235	53.0	0.75	4,009	39.8	0.65	2,954	31.2	0.59	1,513	20.3	0.54
Delay Adjusted	41,110	441.3	2.29	11,324	124.8	1.23	5,306	53.7	0.76	4,044	40.1	0.66	2,974	31.4	0.60	1,535	20.6	0.55
Kansas																		
Observed	32,071	415.7	2.40	9,341	122.5	1.31	4,233	52.1	0.82	2,863	35.3	0.68	1,928	24.5	0.57	1,630	25.0	0.64
Delay Adjusted	^	^	^	^	^	^	^	^	^	^	^	^	^	^	^	^	^	^

1. Rates are per 100,000 population and are age-adjusted by five-year age groups to the 2000 U.S. standard population based on single years of age.
2. See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.
3. Abbreviations: NOS - Not Otherwise Specified; S.E. - Standard Error

^ There are insufficient historical registry data to calculate delay factors necessary for delay adjusted rates.

\- Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for All Sites combined.

Note: Delay adjusted counts are rounded to the nearest whole number. A small registry and/or rare cancer site may have the same delay adjusted count and observed count, but this is just due to rounding to whole counts. For more information about cancer incidence rates adjusted for reporting delay see https://surveillance.cancer.gov/delay/.

Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Delay Adjusted Counts and Rates[1] for the Five Most Commonly Diagnosed Cancers
Selected Areas in the NAACCR United States Combined[2], 2010-2014, White, Females

	All Sites			Breast			Lung & Bronchus			Colon & Rectum			Uterus Corpus & Uterus, NOS[3]			Thyroid		
	Count	*Rate*	*S.E.*[3]	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
Kentucky																		
Observed	57,454	466.9	1.99	15,170	123.4	1.03	10,362	80.4	0.80	5,293	42.0	0.59	3,280	26.1	0.47	2,349	22.2	0.47
Delay Adjusted	58,781	477.7	2.04	15,339	124.8	1.04	10,483	81.3	0.81	5,391	42.7	0.60	3,319	26.4	0.47	2,384	22.5	0.48
Louisiana																		
Observed	38,753	417.9	2.18	11,185	121.0	1.18	5,591	57.2	0.78	3,625	38.2	0.65	1,724	18.3	0.45	1,786	22.2	0.54
Delay Adjusted	39,138	422.0	2.20	11,269	121.9	1.18	5,625	57.5	0.78	3,639	38.3	0.65	1,736	18.4	0.46	1,810	22.5	0.55
Maine																		
Observed	20,337	446.7	3.27	5,655	126.0	1.75	3,145	65.3	1.19	1,612	34.2	0.88	1,488	31.0	0.83	787	22.0	0.83
Delay Adjusted	20,691	454.4	3.32	5,758	128.4	1.78	3,181	66.0	1.20	1,625	34.5	0.89	1,500	31.3	0.84	792	22.1	0.83
Maryland																		
Observed	50,054	428.8	1.98	15,248	131.4	1.10	6,781	55.6	0.69	3,831	31.6	0.53	3,299	27.1	0.48	2,419	24.7	0.52
Delay Adjusted	51,328	439.7	2.03	15,551	134.0	1.12	6,943	56.9	0.71	3,887	32.1	0.53	3,344	27.5	0.49	2,477	25.3	0.53
Massachusetts																		
Observed	84,165	449.6	1.60	25,550	139.2	0.90	12,209	62.5	0.58	6,590	33.4	0.43	5,736	29.7	0.40	4,530	29.1	0.45
Delay Adjusted	85,719	457.7	1.63	25,728	140.2	0.91	12,436	63.6	0.59	6,675	33.8	0.43	5,775	29.9	0.40	4,565	29.3	0.45
Michigan																		
Observed	109,106	416.4	1.31	31,272	120.8	0.71	16,129	58.3	0.47	8,910	32.5	0.35	7,957	29.3	0.34	4,451	20.8	0.33
Delay Adjusted	111,322	424.8	1.33	31,701	122.4	0.72	16,409	59.3	0.48	9,057	33.0	0.36	8,051	29.7	0.34	4,513	21.1	0.33
Detroit																		
Observed	40,761	460.2	2.36	11,601	132.5	1.27	6,127	66.0	0.86	3,322	35.3	0.63	2,897	31.9	0.61	1,965	26.9	0.63
Delay Adjusted	41,306	466.3	2.39	11,707	133.7	1.28	6,166	66.4	0.87	3,348	35.6	0.64	2,916	32.1	0.61	1,989	27.2	0.64
Mississippi																		
Observed	24,139	406.5	2.71	6,624	112.5	1.43	3,868	60.6	0.99	2,318	37.3	0.80	1,117	18.7	0.58	884	18.3	0.64
Delay Adjusted	24,719	416.2	2.77	6,743	114.6	1.46	3,923	61.5	1.00	2,348	37.8	0.81	1,128	18.9	0.58	905	18.8	0.66
Missouri																		
Observed	69,483	420.9	1.65	20,383	125.1	0.90	11,178	64.2	0.62	6,183	35.9	0.47	4,417	26.2	0.40	2,539	18.5	0.38
Delay Adjusted	71,273	431.6	1.69	20,717	127.2	0.92	11,376	65.3	0.63	6,288	36.5	0.48	4,452	26.4	0.41	2,564	18.6	0.38
Montana																		
Observed	12,333	410.9	3.86	3,674	122.9	2.12	1,659	52.0	1.30	1,026	33.0	1.07	805	25.4	0.93	523	21.8	1.00
Delay Adjusted	12,868	428.6	4.03	3,804	127.2	2.20	1,695	53.1	1.33	1,061	34.1	1.10	829	26.1	0.96	534	22.3	1.03
Nebraska																		
Observed	21,013	408.6	2.91	6,197	122.3	1.60	2,683	49.8	0.98	2,051	37.8	0.86	1,406	27.0	0.74	897	21.0	0.72
Delay Adjusted	21,816	424.2	3.02	6,296	124.2	1.63	2,739	50.8	1.00	2,099	38.7	0.88	1,415	27.1	0.74	911	21.3	0.73

Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Delay Adjusted Counts and Rates[1] for the Five Most Commonly Diagnosed Cancers
Selected Areas in the NAACCR United States Combined[2], 2010-2014, White, Females

	All Sites			Breast			Lung & Bronchus			Colon & Rectum			Uterus Corpus & Uterus, NOS[3]			Thyroid		
	Count	*Rate*	*S.E.*[3]	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
New Hampshire																		
Observed	18,755	456.6	3.45	5,809	141.7	1.92	2,689	63.8	1.26	1,447	34.2	0.93	1,432	32.9	0.90	804	23.0	0.85
Delay Adjusted	19,205	467.5	3.53	5,873	143.2	1.94	2,744	65.1	1.28	1,467	34.7	0.94	1,448	33.3	0.91	817	23.3	0.86
New Jersey																		
Observed	101,550	460.8	1.50	29,252	135.9	0.82	12,937	55.5	0.50	8,916	37.6	0.41	7,387	32.6	0.39	5,534	30.4	0.42
Delay Adjusted	104,491	474.2	1.54	29,728	138.1	0.83	13,108	56.3	0.50	9,054	38.2	0.42	7,492	33.1	0.39	5,688	31.2	0.43
New York																		
Observed	213,175	462.6	1.04	60,373	134.4	0.56	28,724	58.9	0.35	17,454	35.6	0.28	15,073	32.0	0.27	11,540	30.2	0.29
Delay Adjusted	217,603	472.2	1.06	61,248	136.3	0.57	28,919	59.3	0.36	17,668	36.1	0.28	15,228	32.3	0.27	11,757	30.7	0.30
North Carolina																		
Observed	94,838	417.4	1.39	28,898	128.0	0.77	14,045	58.5	0.50	7,335	31.6	0.38	5,696	24.2	0.33	3,771	19.4	0.33
Delay Adjusted	97,644	429.7	1.43	29,662	131.4	0.79	14,279	59.5	0.51	7,470	32.1	0.38	5,765	24.5	0.33	3,869	19.9	0.33
North Dakota																		
Observed	7,895	403.3	4.75	2,281	121.5	2.67	969	46.8	1.54	823	39.2	1.42	502	25.2	1.16	398	24.6	1.28
Delay Adjusted	8,120	414.5	4.87	2,302	122.6	2.69	993	48.0	1.58	845	40.2	1.46	513	25.8	1.19	409	25.3	1.32
Ohio																		
Observed	134,587	420.5	1.18	38,759	122.6	0.64	19,818	58.9	0.43	11,855	35.1	0.33	9,611	29.2	0.31	5,832	22.2	0.30
Delay Adjusted	139,829	436.7	1.23	39,791	125.8	0.66	20,323	60.4	0.44	12,195	36.1	0.34	9,796	29.7	0.31	6,001	22.9	0.31
Oklahoma																		
Observed	37,575	393.5	2.09	10,765	113.6	1.13	5,754	56.9	0.76	3,438	34.7	0.61	2,021	20.8	0.48	1,414	17.7	0.49
Delay Adjusted	38,042	398.4	2.11	10,851	114.5	1.14	5,813	57.5	0.77	3,457	34.9	0.61	2,039	21.0	0.48	1,425	17.8	0.49
Oregon																		
Observed	47,057	416.6	1.99	14,307	126.9	1.10	6,366	53.7	0.69	3,615	30.8	0.53	3,141	26.6	0.49	1,818	19.5	0.47
Delay Adjusted	48,105	425.8	2.03	14,550	129.0	1.12	6,484	54.7	0.70	3,687	31.5	0.54	3,177	26.9	0.50	1,849	19.8	0.48
Pennsylvania																		
Observed	169,599	452.3	1.15	47,785	130.3	0.62	22,296	55.3	0.38	15,001	36.9	0.31	12,590	32.8	0.30	9,230	31.5	0.34
Delay Adjusted	172,809	460.8	1.17	48,306	131.7	0.63	22,546	55.9	0.38	15,150	37.2	0.32	12,665	33.0	0.30	9,376	32.0	0.35
Rhode Island																		
Observed	14,486	461.7	4.01	4,066	133.5	2.18	2,155	64.9	1.44	1,185	34.6	1.06	1,054	32.6	1.04	723	28.4	1.10
Delay Adjusted	14,798	471.7	4.10	4,128	135.6	2.21	2,186	65.9	1.46	1,196	35.0	1.07	1,079	33.4	1.06	728	28.6	1.11

1. Rates are per 100,000 population and are age-adjusted by five-year age groups to the 2000 U.S. standard population based on single years of age.
2. See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.
3. Abbreviations: NOS - Not Otherwise Specified; S.E. - Standard Error

^ There are insufficient historical registry data to calculate delay factors necessary for delay adjusted rates.

\- Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for All Sites combined.

Note: Delay adjusted counts are rounded to the nearest whole number. A small registry and/or rare cancer site may have the same delay adjusted count and observed count, but this is just due to rounding to whole counts. For more information about cancer incidence rates adjusted for reporting delay see https://surveillance.cancer.gov/delay/.

Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Delay Adjusted Counts and Rates[1] for the Five Most Commonly Diagnosed Cancers
Selected Areas in the NAACCR United States Combined[2], 2010-2014, White, Females

	All Sites			Prostate			Lung & Bronchus			Colon & Rectum			Uterus Corpus & Uterus, NOS[3]			Thyroid		
	Count	*Rate*	*S.E.[3]*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
South Carolina																		
Observed	44,759	414.6	2.03	13,779	127.8	1.13	6,663	57.7	0.72	3,606	32.6	0.56	2,449	21.9	0.46	1,709	19.1	0.48
Delay Adjusted	46,365	429.3	2.10	14,158	131.3	1.16	6,820	59.0	0.74	3,701	33.4	0.57	2,502	22.4	0.47	1,758	19.7	0.50
South Dakota																		
Observed	9,772	421.3	4.46	2,951	131.0	2.52	1,222	49.7	1.46	970	38.1	1.28	676	28.8	1.15	386	20.7	1.10
Delay Adjusted	10,213	439.7	4.65	3,011	133.7	2.58	1,263	51.4	1.51	1,003	39.4	1.33	684	29.1	1.16	394	21.1	1.12
Tennessee																		
Observed	69,789	417.3	1.62	20,059	120.3	0.87	11,096	62.8	0.60	6,085	35.4	0.46	3,959	23.2	0.38	2,852	20.1	0.39
Delay Adjusted	71,176	425.6	1.65	20,370	122.2	0.89	11,221	63.5	0.61	6,142	35.7	0.47	3,996	23.4	0.38	2,898	20.5	0.40
Texas																		
Observed	209,645	376.0	0.83	62,605	112.2	0.45	25,239	45.0	0.29	17,579	31.3	0.24	12,211	21.3	0.20	10,162	19.0	0.19
Delay Adjusted	218,287	391.4	0.87	64,408	115.4	0.47	25,738	45.9	0.29	17,947	31.9	0.24	12,454	21.8	0.20	10,651	20.0	0.20
Utah																		
Observed	21,835	370.5	2.54	6,730	115.9	1.43	1,374	23.8	0.65	1,583	27.3	0.69	1,545	25.6	0.66	1,781	29.5	0.71
Delay Adjusted	22,189	376.6	2.58	6,754	116.3	1.44	1,391	24.1	0.66	1,596	27.5	0.70	1,550	25.7	0.66	1,788	29.7	0.72
Vermont																		
Observed	8,749	432.0	4.81	2,614	130.2	2.65	1,250	59.0	1.71	701	33.1	1.29	708	33.3	1.30	308	19.2	1.16
Delay Adjusted	8,860	437.4	4.87	2,628	130.9	2.67	1,263	59.6	1.73	711	33.5	1.31	711	33.4	1.30	310	19.3	1.16
Virginia																		
Observed	72,211	397.9	1.51	23,025	127.3	0.86	9,910	52.9	0.54	5,839	31.5	0.42	4,663	24.5	0.37	3,238	20.5	0.37
Delay Adjusted	73,584	405.5	1.54	23,364	129.1	0.87	10,022	53.5	0.55	5,889	31.8	0.42	4,708	24.8	0.37	3,276	20.7	0.37
Washington																		
Observed	75,846	437.1	1.63	23,797	137.1	0.92	9,766	54.8	0.57	5,767	32.8	0.44	4,782	26.0	0.39	3,089	20.4	0.38
Delay Adjusted	77,808	448.4	1.68	24,240	139.6	0.93	9,927	55.7	0.58	5,895	33.5	0.45	4,886	26.6	0.40	3,183	21.0	0.39
Seattle																		
Observed	51,932	454.7	2.06	16,502	144.2	1.16	6,475	55.7	0.71	3,853	33.3	0.55	3,287	27.1	0.49	2,080	20.6	0.46
Delay Adjusted	52,385	458.7	2.07	16,528	144.4	1.16	6,486	55.8	0.71	3,872	33.5	0.56	3,313	27.3	0.49	2,102	20.8	0.47
West Virginia																		
Observed	26,414	439.6	2.81	6,848	114.5	1.44	4,305	66.8	1.04	2,578	41.0	0.83	1,870	30.6	0.74	1,184	24.5	0.74
Delay Adjusted	27,027	449.7	2.88	7,043	117.8	1.48	4,380	68.0	1.06	2,622	41.7	0.85	1,889	30.9	0.75	1,211	25.1	0.76
Wisconsin																		
Observed	69,271	423.8	1.66	20,523	127.9	0.92	9,115	53.4	0.57	5,673	32.8	0.45	5,071	29.7	0.43	2,695	19.9	0.40
Delay Adjusted	71,402	436.1	1.71	21,160	131.8	0.95	9,425	55.2	0.59	5,854	33.8	0.46	5,268	30.9	0.45	2,773	20.5	0.41

Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Delay Adjusted Counts and Rates[1] for the Five Most Commonly Diagnosed Cancers
Selected Areas in the NAACCR United States Combined[2], 2010-2014, White, Females

	All Sites			Prostate			Lung & Bronchus			Colon & Rectum			Uterus Corpus & Uterus, NOS[3]			Thyroid		
	Count	*Rate*	*S.E.[3]*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
Wyoming																		
Observed	5,863	381.3	5.14	1,781	115.0	2.82	706	44.6	1.71	474	30.6	1.44	342	21.5	1.20	299	21.9	1.32
Delay Adjusted	5,930	385.7	5.20	1,799	116.2	2.85	710	44.8	1.72	478	30.8	1.45	345	21.7	1.21	302	22.1	1.33

1. Rates are per 100,000 population and are age-adjusted by five-year age groups to the 2000 U.S. standard population based on single years of age.
2. See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.
3. Abbreviations: NOS - Not Otherwise Specified; S.E. - Standard Error

^ There are insufficient historical registry data to calculate delay factors necessary for delay adjusted rates.

\- Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for All Sites combined.

Note: Delay adjusted counts are rounded to the nearest whole number. A small registry and/or rare cancer site may have the same delay adjusted count and observed count, but this is just due to rounding to whole counts. For more information about cancer incidence rates adjusted for reporting delay see https://surveillance.cancer.gov/delay/.

NAACCR U.S. Combined Delay Adjusted Incidence 2010-2014, Black Combined

Average Annual Age-adjusted Cancer Incidence Delay Adjusted Counts and Rates*, Selected Sites
NAACCR United States Combined[1], 2010-2014, Black

	MALES (ALL RACES)						FEMALES (ALL RACES)					
	Observed			Delay Adjusted			Observed			Delay Adjusted		
SITES	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
All Sites (invasive & in situ bladder)	427,618	542.4	0.89	441,499	560.1	0.92	412,162	394.3	0.63	425,740	407.3	0.65
Oral Cavity & Pharynx	12,501	14.5	0.14	12,823	14.9	0.14	5,489	5.1	0.07	5,631	5.2	0.07
Digestive												
Esophagus	5,558	7.1	0.10	5,633	7.2	0.10	2,380	2.3	0.05	2,414	2.3	0.05
Stomach	10,118	14.1	0.15	10,291	14.3	0.15	7,724	7.7	0.09	7,853	7.9	0.09
Colon & Rectum	42,108	55.2	0.29	43,081	56.5	0.30	42,301	41.0	0.20	43,274	41.9	0.21
Liver & IBD[2]	15,443	16.9	0.14	15,845	17.4	0.15	5,478	5.0	0.07	5,619	5.2	0.07
Pancreas	12,593	16.9	0.16	12,805	17.2	0.16	14,447	14.4	0.12	14,684	14.6	0.12
Respiratory												
Larynx	6,871	8.5	0.11	6,987	8.7	0.11	1,828	1.7	0.04	1,859	1.7	0.04
Lung & bronchus	62,843	85.9	0.37	63,827	87.3	0.37	50,093	49.1	0.22	50,895	49.9	0.23
Skin												
Melanomas of the skin	828	1.1	0.04	869	1.2	0.04	981	1.0	0.03	1,028	1.0	0.03
Breast (female cases only)	-	-	-	-	-	-	130,554	123.0	0.35	133,946	126.2	0.36
Female Genital												
Cervix uteri	-	-	-	-	-	-	9,792	9.3	0.10	9,972	9.5	0.10
Corpus & uterus, NOS[2]	-	-	-	-	-	-	27,125	25.2	0.16	27,597	25.6	0.16
Ovary	-	-	-	-	-	-	9,641	9.2	0.10	9,953	9.5	0.10
Male Genital												
Prostate	150,353	183.6	0.50	158,159	193.4	0.53	-	-	-	-	-	-
Testis	1,464	1.5	0.04	1,494	1.5	0.04	-	-	-	-	-	-
Urinary												
Bladder (incl. *in situ*)	13,090	19.5	0.18	13,501	20.1	0.19	6,440	6.6	0.08	6,636	6.8	0.09
Kidney & renal pelvis	19,982	24.2	0.18	20,582	24.9	0.19	13,230	12.6	0.11	13,620	13.0	0.12
Brain & Other Nervous System	4,255	4.8	0.08	4,394	4.9	0.08	3,756	3.5	0.06	3,876	3.6	0.06
Endocrine												
Thyroid gland	3,272	3.8	0.07	3,362	3.9	0.07	14,716	13.6	0.11	15,110	14.0	0.12
Lymphomas												
Hodgkin Lymphoma	3,042	3.1	0.06	3,114	3.2	0.06	2,542	2.3	0.05	2,601	2.3	0.05
Non-Hodgkin Lymphoma	13,793	16.9	0.15	14,329	17.5	0.16	12,476	12.0	0.11	12,962	12.4	0.11
Multiple Myeloma	11,309	15.4	0.16	12,333	16.8	0.17	11,556	11.4	0.11	12,580	12.4	0.12
Leukemias	10,295	13.3	0.14	11,290	14.6	0.16	8,829	8.5	0.09	9,677	9.4	0.10

* Rates are per 100,000 population and are age-adjusted by five-year age groups to the 2000 U.S. standard population based on single years of age and to standard.

\- Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for All Sites combined.

[1] NAACCR United States Combined includes registries for which a U.S. average delay adjustment factor was calculated. See technical notes.

[2] Abbreviations: IBD – Intrahepatic Bile Duct; NOS - Not Otherwise Specified

Note: Delay adjusted counts are rounded to the nearest whole number. A small registry and/or rare cancer site may have the same delay adjusted count and observed count, but this is just due to rounding to whole counts. For more information about cancer incidence rates adjusted for reporting delay see https://surveillance.cancer.gov/delay/.

Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Delay Adjusted Counts and Rates[1] for the Five Most Commonly Diagnosed Cancers
Selected Areas in the NAACCR United States Combined[2], 2010-2014, Black, Males

	All Sites			Prostate			Lung & Bronchus			Colon & Rectum			Kidney & Renal Pelvis			Liver & IBD[3]		
	Count	*Rate*	*S.E.[3]*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
NAACCR U.S. Combined																		
Observed	427,618	542.4	0.89	150,353	183.6	0.50	62,843	85.9	0.37	42,108	55.2	0.29	19,982	24.2	0.18	15,443	16.9	0.14
Delay Adjusted	441,499	560.1	0.92	158,159	193.4	0.53	63,827	87.3	0.37	43,081	56.5	0.30	20,582	24.9	0.19	15,845	17.4	0.15
Alabama																		
Observed	14,334	563.7	5.06	5,176	196.7	2.93	2,327	95.4	2.13	1,544	63.6	1.74	620	23.8	1.02	365	12.4	0.69
Delay Adjusted	14,617	574.8	5.16	5,401	205.8	3.08	2,342	96.0	2.14	1,558	64.1	1.75	628	24.1	1.03	370	12.6	0.70
Alaska																		
Observed	217	458.0	38.66	88	164.7	21.37	27	78.5	17.89	21	43.3	11.58	8	12.4	5.02	9	16.0	6.24
Delay Adjusted	225	474.7	40.06	93	174.2	22.67	28	80.3	18.27	21	44.3	11.85	8	12.8	5.13	9	16.4	6.37
Arizona																		
Observed	2,103	386.2	9.42	663	115.1	4.91	297	63.0	4.05	197	38.2	3.10	128	22.2	2.20	84	14.0	1.66
Delay Adjusted	2,270	417.1	10.18	747	129.8	5.56	305	64.6	4.16	207	40.0	3.25	134	23.2	2.30	87	14.5	1.71
Arkansas																		
Observed	5,048	573.3	8.71	1,672	178.9	4.70	969	115.7	4.02	510	59.8	2.85	208	23.6	1.75	138	13.4	1.22
Delay Adjusted	5,206	591.3	8.99	1,746	186.8	4.90	984	117.4	4.08	519	60.9	2.91	217	24.6	1.83	139	13.5	1.23
California																		
Observed	27,387	497.9	3.19	9,685	166.5	1.78	3,411	67.9	1.23	2,764	52.4	1.06	1,339	23.3	0.67	1,090	17.7	0.56
Delay Adjusted	28,922	526.1	3.37	10,440	179.8	1.93	3,541	70.5	1.28	2,879	54.5	1.10	1,409	24.5	0.70	1,139	18.5	0.59
Greater California																		
Observed	10,882	480.8	5.03	3,891	163.6	2.83	1,355	67.6	2.00	1,116	51.9	1.70	553	22.7	1.04	442	17.4	0.89
Delay Adjusted	11,601	513.2	5.38	4,222	178.1	3.09	1,418	70.7	2.09	1,165	54.2	1.78	591	24.3	1.12	464	18.2	0.94
Greater Bay																		
Observed	5,963	532.1	7.29	2,071	173.0	4.00	773	74.3	2.83	531	49.6	2.28	293	26.2	1.61	291	22.4	1.38
Delay Adjusted	6,259	558.7	7.65	2,219	185.6	4.30	800	77.0	2.94	556	52.0	2.39	304	27.2	1.68	302	23.3	1.43
Los Angeles																		
Observed	10,531	498.7	5.04	3,717	166.3	2.82	1,280	65.0	1.89	1,116	54.6	1.70	492	22.4	1.04	357	15.5	0.85
Delay Adjusted	11,051	523.4	5.29	3,992	178.7	3.03	1,320	67.1	1.95	1,158	56.7	1.76	513	23.3	1.09	373	16.2	0.89
Colorado																		
Observed	1,993	446.7	11.18	696	151.8	6.44	223	58.6	4.32	201	47.4	3.75	105	22.2	2.33	100	18.4	2.00
Delay Adjusted	2,057	461.3	11.55	740	162.4	6.93	225	59.0	4.35	204	48.0	3.79	107	22.8	2.39	101	18.5	2.02
Connecticut																		
Observed	3,724	515.5	9.07	1,351	179.0	5.18	465	71.7	3.57	347	50.3	2.92	172	20.3	1.63	111	13.9	1.40
Delay Adjusted	3,820	529.0	9.32	1,411	187.4	5.43	471	72.7	3.62	351	51.0	2.96	176	20.8	1.66	113	14.2	1.42

Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Delay Adjusted Counts and Rates[1] for the Five Most Commonly Diagnosed Cancers
Selected Areas in the NAACCR United States Combined[2], 2010-2014, Black, Males

	All Sites			Prostate			Lung & Bronchus			Colon & Rectum			Kidney & Renal Pelvis			Liver & IBD[3]		
	Count	*Rate*	*S.E.*[3]	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
Delaware																		
Observed	2,330	581.8	12.91	926	214.9	7.45	294	79.4	4.97	184	49.9	3.98	124	30.8	2.96	84	18.5	2.16
Delay Adjusted	2,404	600.2	13.33	971	225.5	7.82	301	81.3	5.09	192	52.1	4.15	129	32.0	3.08	88	19.3	2.24
District of Columbia																		
Observed	4,593	597.0	9.08	1,461	186.0	5.02	685	89.4	3.52	454	61.4	2.97	198	25.5	1.87	255	29.6	1.91
Delay Adjusted	4,732	615.1	9.36	1,513	192.7	5.20	699	91.3	3.60	467	63.2	3.06	205	26.4	1.93	259	30.1	1.94
Florida																		
Observed	28,929	470.9	2.93	10,680	169.7	1.73	3,787	65.5	1.13	2,940	49.5	0.97	1,098	17.0	0.54	838	12.0	0.44
Delay Adjusted	30,165	490.5	3.05	11,327	180.2	1.83	3,815	66.0	1.14	3,034	50.9	1.00	1,138	17.7	0.56	857	12.3	0.45
Georgia																		
Observed	29,468	558.6	3.66	10,901	201.0	2.14	4,073	84.8	1.50	3,016	58.0	1.19	1,302	23.0	0.71	813	12.9	0.50
Delay Adjusted	30,069	570.3	3.74	11,263	207.9	2.22	4,117	85.7	1.51	3,059	58.8	1.21	1,330	23.5	0.72	824	13.1	0.50
Hawaii																		
Observed	257	459.9	33.85	104	162.8	17.98	21	48.6	12.61	26	47.6	11.43	10	15.3	5.46	12	17.0	5.22
Delay Adjusted	264	472.7	34.85	107	167.7	18.54	21	49.1	12.72	26	48.3	11.62	10	15.7	5.56	12	17.7	5.45
Idaho																		
Observed	68	351.3	52.05	26	118.8	26.76	6	44.1	23.49	-	-	-	-	-	-	8	50.6	24.40
Delay Adjusted	75	390.6	58.25	27	124.9	28.21	6	45.0	23.96	-	-	-	-	-	-	8	53.5	25.93
Illinois																		
Observed	21,456	574.2	4.12	6,996	179.3	2.24	3,484	98.5	1.76	2,406	66.8	1.43	1,043	26.8	0.87	661	15.9	0.65
Delay Adjusted	21,907	586.5	4.21	7,260	186.5	2.34	3,526	99.7	1.78	2,437	67.6	1.45	1,063	27.3	0.89	675	16.3	0.66
Indiana																		
Observed	5,635	511.5	7.29	1,616	140.2	3.71	982	96.8	3.30	577	53.6	2.39	322	26.6	1.57	223	16.9	1.18
Delay Adjusted	5,905	536.0	7.64	1,711	149.1	3.97	1,004	99.0	3.37	593	55.1	2.45	334	27.5	1.63	232	17.6	1.23
Iowa																		
Observed	745	531.4	22.44	220	159.3	12.06	124	94.6	9.66	75	49.8	6.80	45	31.6	5.47	35	20.7	3.96
Delay Adjusted	770	549.7	23.23	237	171.3	12.95	126	96.1	9.82	75	50.1	6.84	46	32.2	5.60	36	21.1	4.07
Kansas																		
Observed	1,869	562.7	14.13	630	186.4	8.01	316	99.8	6.08	190	61.6	4.86	84	23.6	2.75	65	16.5	2.17
Delay Adjusted	^	^	^	^	^	^	^	^	^	^	^	^	^	^	^	^	^	^

1. Rates are per 100,000 population and are age-adjusted by five-year age groups to the 2000 U.S. standard population based on single years of age.
2. See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.
3. Abbreviations: IBD - Intrahepatic Bile Duct; S.E. - Standard Error

^ There are insufficient historical registry data to calculate delay factors necessary for delay adjusted rates.

\- Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for All Sites combined.

Note: Delay adjusted counts are rounded to the nearest whole number. A small registry and/or rare cancer site may have the same delay adjusted count and observed count, but this is just due to rounding to whole counts. For more information about cancer incidence rates adjusted for reporting delay see https://surveillance.cancer.gov/delay/.

Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Delay Adjusted Counts and Rates[1] for the Five Most Commonly Diagnosed Cancers
Selected Areas in the NAACCR United States Combined[2], 2010-2014, Black, Males

	All Sites			Prostate			Lung & Bronchus			Colon & Rectum			Kidney & Renal Pelvis			Liver & IBD[3]		
	Count	*Rate*	*S.E.[3]*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
Kentucky																		
Observed	3,997	593.0	10.19	1,158	165.2	5.24	733	118.4	4.75	415	62.8	3.36	236	31.6	2.19	142	17.8	1.61
Delay Adjusted	4,088	606.3	10.41	1,220	173.6	5.49	741	119.8	4.80	423	63.9	3.42	241	32.3	2.24	145	18.2	1.64
Louisiana																		
Observed	17,798	617.5	4.97	5,907	199.7	2.78	2,977	109.4	2.16	1,914	66.7	1.64	798	27.5	1.03	676	19.7	0.80
Delay Adjusted	18,105	628.3	5.06	6,059	204.9	2.85	3,012	110.8	2.19	1,937	67.5	1.66	806	27.8	1.04	688	20.1	0.82
Maine																		
Observed	111	359.6	37.96	36	121.0	22.09	8	31.9	12.31	7	17.8	6.96	-	-	-	7	25.3	11.19
Delay Adjusted	113	367.4	38.79	37	124.4	22.79	8	32.4	12.49	7	18.0	7.02	-	-	-	7	26.0	11.51
Maryland																		
Observed	18,583	529.6	4.19	7,160	192.7	2.43	2,365	74.2	1.65	1,660	50.1	1.33	805	22.2	0.84	759	18.6	0.72
Delay Adjusted	19,355	551.9	4.37	7,449	201.1	2.55	2,422	76.1	1.70	1,735	52.4	1.39	837	23.0	0.87	784	19.2	0.75
Massachusetts																		
Observed	4,753	488.9	7.65	1,853	183.3	4.54	526	62.0	2.91	408	43.2	2.32	192	19.0	1.47	204	18.8	1.42
Delay Adjusted	4,994	513.7	8.05	1,946	192.5	4.77	535	63.1	2.97	429	45.4	2.44	202	20.0	1.55	223	20.6	1.55
Michigan																		
Observed	16,993	584.3	4.73	5,903	194.9	2.67	2,615	96.5	1.99	1,551	55.1	1.48	927	31.2	1.08	657	19.6	0.80
Delay Adjusted	17,228	592.3	4.79	6,046	199.8	2.74	2,644	97.5	2.01	1,568	55.7	1.49	950	32.0	1.10	668	19.9	0.81
Detroit																		
Observed	12,917	630.5	5.82	4,594	214.8	3.32	1,934	101.4	2.42	1,191	60.2	1.83	705	33.8	1.33	510	21.5	1.00
Delay Adjusted	13,093	639.2	5.90	4,703	219.9	3.40	1,946	102.0	2.44	1,200	60.7	1.84	722	34.7	1.36	513	21.6	1.00
Mississippi																		
Observed	12,681	606.1	5.82	4,356	202.1	3.30	2,264	113.8	2.59	1,479	72.6	2.05	601	28.5	1.25	229	9.4	0.66
Delay Adjusted	12,984	620.4	5.95	4,504	209.2	3.42	2,296	115.4	2.62	1,498	73.5	2.07	616	29.3	1.28	237	9.7	0.69
Missouri																		
Observed	7,525	561.3	6.90	2,237	158.5	3.55	1,342	105.4	3.06	748	60.0	2.34	362	25.8	1.44	369	23.6	1.29
Delay Adjusted	7,719	576.0	7.09	2,392	170.0	3.83	1,362	107.0	3.11	759	60.9	2.38	373	26.6	1.48	377	24.1	1.32
Montana																		
Observed	50	554.1	90.53	17	182.5	47.17	9	119.2	43.10	-	-	-	-	-	-	-	-	-
Delay Adjusted	52	580.1	95.21	18	195.4	50.67	9	121.7	44.13	-	-	-	-	-	-	-	-	-
Nebraska																		
Observed	895	606.1	22.32	310	204.1	12.62	125	96.3	9.36	107	78.6	8.45	54	35.5	5.34	36	21.6	3.87
Delay Adjusted	929	629.5	23.18	333	219.3	13.57	127	98.1	9.54	110	80.4	8.65	57	37.3	5.61	38	22.9	4.07

Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Delay Adjusted Counts and Rates[1] for the Five Most Commonly Diagnosed Cancers
Selected Areas in the NAACCR United States Combined[2], 2010-2014, Black, Males

	All Sites			Prostate			Lung & Bronchus			Colon & Rectum			Kidney & Renal Pelvis			Liver & IBD[3]		
	Count	*Rate*	*S.E.*[3]	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
New Hampshire																		
Observed	135	399.6	38.46	52	151.0	23.13	12	35.8	11.09	15	43.4	12.22	6	17.1	7.78	7	23.2	9.74
Delay Adjusted	138	408.9	39.37	53	155.2	23.80	12	36.6	11.39	15	44.0	12.40	6	17.5	8.01	7	24.0	10.06
New Jersey																		
Observed	13,273	525.0	4.83	5,172	199.5	2.92	1,622	70.5	1.86	1,253	50.7	1.52	610	22.7	0.97	409	14.1	0.74
Delay Adjusted	13,926	550.7	5.07	5,634	217.6	3.19	1,651	71.8	1.89	1,315	53.1	1.59	629	23.4	0.99	418	14.5	0.75
New York																		
Observed	38,150	551.4	2.97	15,490	216.8	1.82	4,217	65.2	1.06	3,371	51.3	0.93	1,574	21.8	0.57	1,419	18.6	0.52
Delay Adjusted	39,024	564.0	3.04	16,214	227.2	1.91	4,245	65.6	1.07	3,425	52.1	0.95	1,611	22.3	0.59	1,451	19.0	0.53
North Carolina																		
Observed	24,403	562.6	3.88	8,576	189.5	2.19	4,030	99.4	1.69	2,232	52.8	1.21	1,120	25.1	0.80	737	14.5	0.57
Delay Adjusted	25,330	583.9	4.03	9,093	200.8	2.33	4,096	101.0	1.72	2,296	54.3	1.24	1,149	25.8	0.82	751	14.7	0.58
North Dakota																		
Observed	46	422.3	87.24	14	116.8	41.09	7	98.0	45.95	6	58.4	32.24	-	-	-	-	-	-
Delay Adjusted	47	435.3	89.90	15	121.9	43.09	7	100.1	46.80	6	59.9	32.96	-	-	-	-	-	-
Ohio																		
Observed	15,668	528.1	4.42	5,272	170.5	2.46	2,660	95.3	1.93	1,459	50.6	1.39	770	24.9	0.93	607	17.6	0.75
Delay Adjusted	16,257	548.5	4.60	5,601	181.1	2.61	2,727	97.7	1.98	1,501	52.1	1.43	796	25.7	0.96	631	18.3	0.78
Oklahoma																		
Observed	3,140	582.7	11.37	1,082	199.1	6.59	527	102.5	4.85	298	54.1	3.43	144	26.7	2.41	99	15.5	1.70
Delay Adjusted	3,248	602.7	11.76	1,128	207.4	6.86	532	103.5	4.89	303	55.0	3.49	146	27.0	2.44	103	16.1	1.76
Oregon																		
Observed	813	498.2	19.82	263	156.4	10.87	119	81.9	8.55	75	50.4	6.51	44	23.2	3.93	50	29.2	4.64
Delay Adjusted	854	524.2	20.89	284	169.3	11.79	121	83.2	8.68	79	52.8	6.82	45	23.7	4.02	51	29.9	4.75
Pennsylvania																		
Observed	16,392	579.2	4.78	5,209	176.7	2.58	2,506	97.0	2.04	1,475	53.9	1.48	905	30.0	1.04	915	28.2	0.98
Delay Adjusted	16,966	599.5	4.95	5,645	191.5	2.79	2,533	98.0	2.06	1,494	54.6	1.50	936	31.0	1.08	938	28.9	1.00
Rhode Island																		
Observed	572	432.1	19.81	181	124.5	10.02	75	65.2	8.22	43	37.0	6.12	24	17.8	4.01	22	11.9	2.68
Delay Adjusted	585	441.5	20.24	197	135.6	10.91	76	66.3	8.34	43	37.4	6.19	24	18.1	4.09	22	12.2	2.72

1. Rates are per 100,000 population and are age-adjusted by five-year age groups to the 2000 U.S. standard population based on single years of age.
2. See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.
3. Abbreviations: IBD - Intrahepatic Bile Duct; S.E. - Standard Error

^ There are insufficient historical registry data to calculate delay factors necessary for delay adjusted rates.

\- Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for All Sites combined.

Note: Delay adjusted counts are rounded to the nearest whole number. A small registry and/or rare cancer site may have the same delay adjusted count and observed count, but this is just due to rounding to whole counts. For more information about cancer incidence rates adjusted for reporting delay see https://surveillance.cancer.gov/delay/.

Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Delay Adjusted Counts and Rates[1] for the Five Most Commonly Diagnosed Cancers
Selected Areas in the NAACCR United States Combined[2], 2010-2014, Black, Males

	All Sites			Prostate			Lung & Bronchus			Colon & Rectum			Kidney & Renal Pelvis			Liver & IBD[3]		
	Count	*Rate*	*S.E.[3]*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
South Carolina																		
Observed	15,310	556.5	4.84	5,431	188.1	2.73	2,373	90.4	2.00	1,464	54.8	1.54	641	22.7	0.95	388	12.3	0.67
Delay Adjusted	15,874	577.3	5.02	5,726	198.5	2.89	2,434	92.8	2.06	1,511	56.4	1.59	664	23.5	0.99	404	12.8	0.70
South Dakota																		
Observed	61	239.8	35.14	10	32.6	10.55	8	39.6	17.91	7	28.5	11.20	6	21.3	8.93	-	-	-
Delay Adjusted	64	252.1	36.93	11	35.1	11.42	8	40.6	18.30	7	29.6	11.65	6	22.5	9.41	-	-	-
Tennessee																		
Observed	11,111	558.1	5.83	3,743	182.1	3.26	1,900	103.3	2.61	1,094	57.8	1.92	504	24.5	1.19	384	16.0	0.88
Delay Adjusted	11,413	573.2	5.99	3,849	187.3	3.35	1,930	105.0	2.65	1,114	58.9	1.96	521	25.3	1.24	395	16.4	0.91
Texas																		
Observed	29,472	528.4	3.41	9,029	157.5	1.83	4,572	91.3	1.49	3,236	58.4	1.14	1,514	25.5	0.72	1,516	22.4	0.62
Delay Adjusted	30,689	550.2	3.55	9,716	170.3	1.99	4,663	93.2	1.52	3,304	59.7	1.17	1,564	26.4	0.74	1,546	22.9	0.63
Utah																		
Observed	239	492.0	39.28	89	194.2	24.50	24	55.6	13.51	29	57.5	13.48	14	28.4	9.25	16	30.1	8.68
Delay Adjusted	251	517.1	41.28	91	199.2	25.14	24	56.1	13.61	29	58.0	13.59	14	29.1	9.45	17	31.2	8.97
Vermont																		
Observed	35	270.5	52.15	14	108.5	33.13	-	-	-	-	-	-	-	-	-	-	-	-
Delay Adjusted	35	274.7	53.14	14	110.0	33.62	-	-	-	-	-	-	-	-	-	-	-	-
Virginia																		
Observed	17,774	532.8	4.27	6,428	180.7	2.39	2,676	88.0	1.82	1,601	49.6	1.33	902	25.7	0.90	545	13.8	0.62
Delay Adjusted	18,217	545.7	4.37	6,611	185.8	2.46	2,707	88.9	1.84	1,628	50.5	1.35	926	26.4	0.93	561	14.2	0.64
Washington																		
Observed	2,729	511.7	11.24	936	165.7	6.11	315	72.7	4.62	225	43.0	3.27	165	27.2	2.37	140	23.7	2.26
Delay Adjusted	2,799	525.1	11.54	980	173.5	6.41	319	73.7	4.68	229	43.8	3.33	169	27.8	2.43	142	24.0	2.28
Seattle																		
Observed	2,481	528.8	12.17	865	171.9	6.57	286	76.6	5.08	199	43.9	3.56	145	27.2	2.53	134	26.3	2.57
Delay Adjusted	2,535	540.3	12.44	902	179.1	6.85	288	77.1	5.12	202	44.5	3.60	147	27.6	2.58	135	26.4	2.58
West Virginia																		
Observed	851	517.7	19.37	282	159.1	10.23	134	92.4	8.71	92	55.8	6.37	46	25.3	3.96	38	18.7	3.23
Delay Adjusted	871	529.9	19.83	289	163.0	10.48	136	93.9	8.86	94	56.8	6.48	48	26.2	4.10	39	19.2	3.30
Wisconsin																		
Observed	3,870	662.4	11.94	1,243	207.0	6.51	606	116.7	5.28	375	66.2	3.85	193	29.4	2.35	172	24.5	1.98
Delay Adjusted	3,971	680.5	12.27	1,337	222.8	7.01	610	117.5	5.32	387	68.4	3.98	206	31.4	2.51	173	24.7	2.00

Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Delay Adjusted Counts and Rates[1] for the Five Most Commonly Diagnosed Cancers
Selected Areas in the NAACCR United States Combined[2], 2010-2014, Black, Males

	All Sites			Prostate			Lung & Bronchus			Colon & Rectum			Kidney & Renal Pelvis			Liver & IBD[3]		
	Count	*Rate*	*S.E.[3]*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
Wyoming																		
Observed	32	179.2	32.84	9	47.9	16.53	-	-	-	6	40.8	16.67	-	-	-	-	-	-
Delay Adjusted	32	181.8	33.29	9	48.6	16.73	-	-	-	6	41.1	16.79	-	-	-	-	-	-

1. Rates are per 100,000 population and are age-adjusted by five-year age groups to the 2000 U.S. standard population based on single years of age.
2. See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.
3. Abbreviations: IBD - Intrahepatic Bile Duct; S.E. - Standard Error

^ There are insufficient historical registry data to calculate delay factors necessary for delay adjusted rates.

\- Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for All Sites combined.

Note: Delay adjusted counts are rounded to the nearest whole number. A small registry and/or rare cancer site may have the same delay adjusted count and observed count, but this is just due to rounding to whole counts. For more information about cancer incidence rates adjusted for reporting delay see https://surveillance.cancer.gov/delay/.

Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Delay Adjusted Counts and Rates[1] for the Five Most Commonly Diagnosed Cancers
Selected Areas in the NAACCR United States Combined[2], 2010-2014, Black, Females

	All Sites			Breast			Lung & Bronchus			Colon & Rectum			Uterus Corpus & Uterus, NOS[3]			Thyroid		
	Count	*Rate*	*S.E.*[3]	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
NAACCR U.S. Combined																		
Observed	412,162	394.3	0.63	130,554	123.0	0.35	50,093	49.1	0.22	42,301	41.0	0.20	27,125	25.2	0.16	14,716	13.6	0.11
Delay Adjusted	425,740	407.3	0.65	133,946	126.2	0.36	50,895	49.9	0.23	43,274	41.9	0.21	27,597	25.6	0.16	15,110	14.0	0.12
Alabama																		
Observed	12,813	373.8	3.37	4,314	124.4	1.93	1,313	38.5	1.09	1,508	44.4	1.17	750	21.6	0.81	339	9.9	0.55
Delay Adjusted	13,069	381.3	3.44	4,383	126.4	1.97	1,322	38.7	1.09	1,521	44.7	1.18	757	21.8	0.82	342	9.9	0.55
Alaska																		
Observed	157	324.8	27.47	61	126.1	16.94	11	27.1	8.36	12	22.2	6.77	-	-	-	9	15.5	5.53
Delay Adjusted	163	338.0	28.59	61	127.0	17.06	11	27.8	8.58	12	22.6	6.89	-	-	-	9	15.9	5.68
Arizona																		
Observed	1,971	332.2	7.80	625	101.3	4.19	235	42.5	2.88	193	33.9	2.55	134	21.9	1.96	116	17.1	1.63
Delay Adjusted	2,130	358.9	8.43	658	106.6	4.42	241	43.6	2.95	203	35.6	2.67	141	23.0	2.06	122	18.0	1.71
Arkansas																		
Observed	4,312	381.2	5.96	1,332	116.3	3.27	543	48.9	2.16	511	45.6	2.07	252	21.7	1.41	114	9.7	0.92
Delay Adjusted	4,447	393.1	6.14	1,392	121.5	3.41	551	49.6	2.19	520	46.4	2.11	258	22.2	1.44	116	9.8	0.94
California																		
Observed	25,510	382.3	2.44	8,312	123.0	1.37	3,110	47.7	0.87	2,680	40.9	0.80	1,715	24.7	0.61	797	11.8	0.43
Delay Adjusted	26,943	403.8	2.58	8,740	129.3	1.44	3,228	49.5	0.91	2,792	42.6	0.84	1,756	25.2	0.62	823	12.2	0.44
Greater California																		
Observed	9,346	365.7	3.91	3,092	117.0	2.17	1,118	46.8	1.45	959	39.5	1.32	589	22.0	0.93	337	11.8	0.66
Delay Adjusted	9,969	390.3	4.18	3,305	125.1	2.32	1,170	49.0	1.52	1,001	41.3	1.38	606	22.6	0.96	353	12.4	0.69
Greater Bay																		
Observed	5,319	399.8	5.60	1,691	126.2	3.13	701	53.3	2.06	563	42.3	1.83	362	26.3	1.41	107	8.2	0.81
Delay Adjusted	5,587	419.9	5.89	1,775	132.5	3.29	726	55.1	2.13	589	44.3	1.91	370	26.9	1.44	110	8.4	0.83
Los Angeles																		
Observed	10,841	390.1	3.80	3,529	126.8	2.17	1,290	46.1	1.30	1,158	41.5	1.24	764	26.4	0.97	353	13.3	0.72
Delay Adjusted	11,383	409.5	3.99	3,660	131.5	2.25	1,331	47.5	1.34	1,201	43.1	1.28	781	27.0	0.99	360	13.6	0.73
Colorado																		
Observed	1,633	343.0	8.75	555	113.5	4.95	185	42.5	3.22	149	33.3	2.81	88	18.5	2.03	81	14.8	1.68
Delay Adjusted	1,684	353.8	9.03	561	114.8	5.00	186	42.9	3.24	151	33.7	2.84	89	18.6	2.04	82	15.0	1.70
Connecticut																		
Observed	3,790	381.0	6.32	1,191	116.0	3.43	408	42.9	2.18	360	37.1	2.00	274	26.5	1.63	182	17.2	1.29
Delay Adjusted	3,888	390.9	6.49	1,216	118.4	3.50	413	43.5	2.21	364	37.6	2.03	277	26.8	1.65	186	17.5	1.32

Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Delay Adjusted Counts and Rates[1] for the Five Most Commonly Diagnosed Cancers
Selected Areas in the NAACCR United States Combined[2], 2010-2014, Black, Females

	All Sites			Breast			Lung & Bronchus			Colon & Rectum			Uterus Corpus & Uterus, NOS[3]			Thyroid		
	Count	*Rate*	*S.E.[3]*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
Delaware																		
Observed	2,193	418.6	9.19	696	129.4	5.03	278	56.7	3.50	184	35.7	2.71	158	29.5	2.41	95	17.2	1.79
Delay Adjusted	2,264	432.1	9.48	708	131.7	5.12	284	58.0	3.59	192	37.3	2.83	160	29.8	2.44	98	17.6	1.84
District of Columbia																		
Observed	4,895	460.8	6.78	1,451	140.0	3.79	626	56.9	2.32	508	47.0	2.14	387	34.7	1.81	174	17.9	1.42
Delay Adjusted	5,047	475.1	6.99	1,530	147.6	3.99	639	58.1	2.37	522	48.3	2.20	402	36.1	1.88	182	18.8	1.48
Florida																		
Observed	27,263	346.0	2.13	8,474	105.8	1.17	2,677	35.0	0.69	2,776	36.0	0.70	2,052	25.3	0.57	973	11.9	0.39
Delay Adjusted	28,466	361.0	2.23	8,821	110.1	1.22	2,698	35.2	0.69	2,864	37.1	0.72	2,123	26.1	0.59	980	11.9	0.39
Georgia																		
Observed	28,618	385.6	2.37	9,650	125.4	1.32	2,876	41.0	0.80	3,061	42.4	0.80	1,863	24.5	0.59	932	11.4	0.38
Delay Adjusted	29,200	393.6	2.42	9,759	126.9	1.34	2,907	41.4	0.81	3,105	43.0	0.81	1,882	24.8	0.60	948	11.6	0.39
Hawaii																		
Observed	129	315.2	30.79	54	118.4	17.38	11	41.9	13.13	6	19.1	8.88	6	15.4	6.85	-	-	-
Delay Adjusted	132	323.0	31.58	55	120.2	17.65	11	42.4	13.28	6	19.2	8.95	6	15.6	6.93	-	-	-
Idaho																		
Observed	41	319.0	55.40	12	96.0	31.41	-	-	-	-	-	-	-	-	-	-	-	-
Delay Adjusted	45	353.5	61.38	12	97.3	31.83	-	-	-	-	-	-	-	-	-	-	-	-
Illinois																		
Observed	22,300	431.1	2.93	6,781	130.4	1.61	3,228	62.8	1.12	2,489	48.5	0.99	1,469	27.3	0.72	655	12.6	0.50
Delay Adjusted	22,781	440.5	3.00	6,884	132.3	1.63	3,268	63.6	1.14	2,521	49.1	1.00	1,486	27.6	0.73	669	12.9	0.51
Indiana																		
Observed	6,090	416.6	5.46	1,928	129.5	3.01	875	61.6	2.13	625	43.7	1.79	342	22.9	1.27	154	9.9	0.81
Delay Adjusted	6,385	436.8	5.73	1,997	134.1	3.12	895	63.0	2.18	642	44.9	1.84	348	23.3	1.29	158	10.2	0.83
Iowa																		
Observed	659	434.2	17.78	163	103.1	8.43	102	73.0	7.54	65	46.4	5.99	45	29.5	4.57	17	9.2	2.31
Delay Adjusted	681	449.0	18.40	170	107.8	8.81	103	74.0	7.64	65	46.7	6.03	45	29.7	4.61	17	9.3	2.34
Kansas																		
Observed	1,633	419.5	10.58	488	124.2	5.72	238	63.4	4.18	153	39.8	3.27	96	24.8	2.59	77	18.8	2.19
Delay Adjusted	^	^	^	^	^	^	^	^	^	^	^	^	^	^	^	^	^	^

1. Rates are per 100,000 population and are age-adjusted by five-year age groups to the 2000 U.S. standard population based on single years of age.
2. See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.
3. Abbreviations: NOS - Not Otherwise Specified; S.E. - Standard Error

^ There are insufficient historical registry data to calculate delay factors necessary for delay adjusted rates.

\- Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for All Sites combined.

Note: Delay adjusted counts are rounded to the nearest whole number. A small registry and/or rare cancer site may have the same delay adjusted count and observed count, but this is just due to rounding to whole counts. For more information about cancer incidence rates adjusted for reporting delay see https://surveillance.cancer.gov/delay/.

Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Delay Adjusted Counts and Rates[1] for the Five Most Commonly Diagnosed Cancers
Selected Areas in the NAACCR United States Combined[2], 2010-2014, Black, Females

	All Sites			Breast			Lung & Bronchus			Colon & Rectum			Uterus Corpus & Uterus, NOS[3]			Thyroid		
	Count	*Rate*	*S.E.*[3]	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
Kentucky																		
Observed	3,754	438.9	7.34	1,096	126.2	3.90	643	77.9	3.16	423	49.5	2.46	200	23.2	1.68	133	15.1	1.34
Delay Adjusted	3,841	448.9	7.51	1,109	127.7	3.95	650	78.8	3.19	431	50.4	2.51	203	23.6	1.71	135	15.4	1.36
Louisiana																		
Observed	15,563	416.7	3.42	4,975	131.6	1.91	1,858	50.3	1.20	1,863	50.4	1.20	841	21.8	0.77	531	14.0	0.62
Delay Adjusted	15,835	423.9	3.48	5,017	132.7	1.93	1,880	50.9	1.21	1,886	51.0	1.21	847	21.9	0.78	538	14.2	0.63
Maine																		
Observed	76	305.2	37.43	18	69.0	16.81	10	46.3	15.72	-	-	-	7	29.6	11.85	-	-	-
Delay Adjusted	78	312.3	38.31	18	70.6	17.20	10	46.8	15.91	-	-	-	7	29.8	11.93	-	-	-
Maryland																		
Observed	19,357	397.3	2.94	6,575	131.3	1.66	2,235	47.3	1.03	1,779	37.2	0.91	1,321	26.2	0.74	775	15.2	0.55
Delay Adjusted	20,167	414.1	3.06	6,781	135.4	1.71	2,291	48.4	1.06	1,860	38.9	0.95	1,351	26.8	0.76	793	15.5	0.57
Massachusetts																		
Observed	4,569	358.4	5.42	1,387	106.7	2.92	504	42.2	1.92	425	33.6	1.67	296	22.7	1.35	302	21.8	1.27
Delay Adjusted	4,807	377.1	5.70	1,433	110.2	3.02	514	43.0	1.96	446	35.3	1.75	298	22.9	1.36	314	22.7	1.33
Michigan																		
Observed	16,166	422.8	3.40	4,855	126.4	1.85	2,403	63.1	1.32	1,617	42.3	1.07	1,059	26.6	0.84	530	14.0	0.62
Delay Adjusted	16,399	428.8	3.45	4,908	127.8	1.87	2,429	63.8	1.33	1,634	42.8	1.09	1,068	26.8	0.84	538	14.2	0.63
Detroit																		
Observed	12,423	440.3	4.04	3,739	132.1	2.21	1,831	64.6	1.55	1,281	45.5	1.30	798	27.1	0.98	433	15.6	0.76
Delay Adjusted	12,598	446.5	4.09	3,773	133.4	2.23	1,842	65.0	1.56	1,291	45.9	1.31	803	27.2	0.99	440	15.8	0.77
Mississippi																		
Observed	11,046	399.4	3.89	3,399	120.5	2.11	1,271	47.5	1.37	1,436	52.4	1.42	640	22.7	0.92	350	12.1	0.66
Delay Adjusted	11,310	408.9	3.98	3,477	123.3	2.16	1,289	48.2	1.39	1,454	53.1	1.43	645	22.9	0.93	357	12.4	0.67
Missouri																		
Observed	7,961	438.1	5.00	2,439	132.7	2.74	1,215	67.8	1.99	788	43.8	1.59	499	26.8	1.23	228	12.2	0.82
Delay Adjusted	8,165	449.3	5.13	2,482	135.0	2.78	1,234	68.9	2.02	800	44.5	1.61	503	27.0	1.23	235	12.6	0.84
Montana																		
Observed	19	286.3	69.79	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Delay Adjusted	20	296.7	72.12	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Nebraska																		
Observed	734	424.3	16.18	216	125.7	8.79	108	64.3	6.36	71	42.7	5.22	35	19.0	3.32	31	15.3	2.82
Delay Adjusted	762	440.3	16.79	219	127.4	8.91	110	65.7	6.50	73	43.7	5.34	35	19.2	3.35	32	15.6	2.87

Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Delay Adjusted Counts and Rates[1] for the Five Most Commonly Diagnosed Cancers
Selected Areas in the NAACCR United States Combined[2], 2010-2014, Black, Females

	All Sites			Breast			Lung & Bronchus			Colon & Rectum			Uterus Corpus & Uterus, NOS[3]			Thyroid		
	Count	*Rate*	*S.E.[3]*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
New Hampshire																		
Observed	67	207.8	26.51	24	75.9	16.05	8	26.1	9.34	6	19.7	8.52	6	18.6	8.05	-	-	-
Delay Adjusted	69	212.8	27.16	24	76.7	16.23	8	26.5	9.51	6	20.0	8.66	6	18.7	8.10	-	-	-
New Jersey																		
Observed	13,961	393.8	3.38	4,351	120.3	1.85	1,642	47.4	1.19	1,451	41.8	1.11	987	27.2	0.88	490	13.5	0.62
Delay Adjusted	14,670	413.8	3.55	4,533	125.3	1.93	1,673	48.3	1.21	1,522	43.8	1.17	1,009	27.8	0.90	504	13.9	0.63
New York																		
Observed	37,976	382.5	1.99	11,487	115.3	1.09	4,031	40.6	0.65	3,680	37.3	0.62	3,148	30.7	0.55	1,602	16.0	0.40
Delay Adjusted	38,864	391.4	2.04	11,670	117.1	1.11	4,058	40.9	0.65	3,739	37.9	0.63	3,181	31.0	0.56	1,663	16.6	0.42
North Carolina																		
Observed	23,505	401.1	2.67	7,874	132.8	1.53	2,732	47.2	0.92	2,224	38.2	0.83	1,525	25.3	0.66	826	14.0	0.49
Delay Adjusted	24,403	416.4	2.77	8,098	136.6	1.57	2,777	48.0	0.94	2,288	39.3	0.85	1,546	25.7	0.67	848	14.3	0.51
North Dakota																		
Observed	26	295.8	74.27	6	53.3	21.94	-	-	-	-	-	-	-	-	-	-	-	-
Delay Adjusted	27	305.7	76.93	6	53.8	22.16	-	-	-	-	-	-	-	-	-	-	-	-
Ohio																		
Observed	15,372	398.2	3.26	4,751	122.5	1.81	2,429	63.3	1.30	1,454	37.8	1.00	920	23.1	0.77	603	16.1	0.66
Delay Adjusted	15,964	413.7	3.39	4,878	125.8	1.86	2,491	64.9	1.34	1,496	38.9	1.03	948	23.8	0.80	620	16.5	0.68
Oklahoma																		
Observed	2,704	396.7	7.85	845	121.3	4.28	373	57.2	3.05	272	39.8	2.48	165	23.4	1.87	85	11.6	1.29
Delay Adjusted	2,795	410.2	8.12	870	124.9	4.41	377	57.7	3.08	276	40.4	2.52	166	23.6	1.89	86	11.7	1.30
Oregon																		
Observed	614	386.6	16.16	196	120.2	8.84	78	53.5	6.27	54	36.7	5.18	38	22.9	3.82	30	16.6	3.10
Delay Adjusted	645	406.6	17.00	206	126.5	9.30	79	54.4	6.38	57	38.4	5.42	38	23.2	3.88	31	16.9	3.16
Pennsylvania																		
Observed	17,345	452.3	3.49	4,875	127.2	1.85	2,603	68.7	1.37	1,600	41.6	1.05	1,141	28.9	0.87	841	22.0	0.77
Delay Adjusted	17,955	468.2	3.61	4,975	129.8	1.89	2,632	69.5	1.38	1,622	42.1	1.07	1,153	29.2	0.88	874	22.9	0.80
Rhode Island																		
Observed	586	344.4	14.64	179	102.4	7.86	77	47.7	5.58	51	30.9	4.46	33	20.1	3.60	39	20.7	3.38
Delay Adjusted	598	351.6	14.95	182	103.9	7.98	78	48.3	5.65	51	31.3	4.50	34	20.5	3.68	39	20.8	3.40

1. Rates are per 100,000 population and are age-adjusted by five-year age groups to the 2000 U.S. standard population based on single years of age.
2. See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.
3. Abbreviations: NOS - Not Otherwise Specified; S.E. - Standard Error

^ There are insufficient historical registry data to calculate delay factors necessary for delay adjusted rates.

\- Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for All Sites combined.

Note: Delay adjusted counts are rounded to the nearest whole number. A small registry and/or rare cancer site may have the same delay adjusted count and observed count, but this is just due to rounding to whole counts. For more information about cancer incidence rates adjusted for reporting delay see https://surveillance.cancer.gov/delay/.

Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Delay Adjusted Counts and Rates[1] for the Five Most Commonly Diagnosed Cancers
Selected Areas in the NAACCR United States Combined[2], 2010-2014, Black, Females

	All Sites			Prostate			Lung & Bronchus			Colon & Rectum			Uterus Corpus & Uterus, NOS[3]			Thyroid		
	Count	*Rate*	*S.E.*[3]	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
South Carolina																		
Observed	13,949	383.0	3.32	4,583	124.9	1.89	1,540	42.1	1.10	1,370	37.1	1.03	993	26.4	0.86	461	12.8	0.61
Delay Adjusted	14,474	397.5	3.45	4,724	128.8	1.95	1,578	43.2	1.13	1,413	38.2	1.06	1,017	27.0	0.88	474	13.2	0.63
South Dakota																		
Observed	38	200.6	33.45	16	88.0	22.25	-	-	-	-	-	-	-	-	-	-	-	-
Delay Adjusted	40	208.7	34.80	16	89.7	22.66	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee																		
Observed	10,842	398.1	3.93	3,518	125.7	2.18	1,359	52.1	1.46	1,127	42.6	1.31	620	22.0	0.91	297	10.5	0.62
Delay Adjusted	11,142	409.2	4.04	3,580	127.9	2.22	1,381	52.9	1.48	1,148	43.4	1.33	628	22.3	0.92	303	10.7	0.63
Texas																		
Observed	28,768	389.6	2.37	9,030	118.1	1.28	3,377	48.9	0.87	3,062	41.8	0.78	1,696	22.4	0.56	1,019	12.6	0.40
Delay Adjusted	29,958	405.7	2.47	9,360	122.4	1.33	3,443	49.9	0.89	3,126	42.7	0.80	1,740	23.0	0.58	1,069	13.2	0.42
Utah																		
Observed	134	304.2	28.82	35	76.5	13.77	13	34.6	10.26	16	43.2	11.46	6	18.8	8.15	8	13.2	5.18
Delay Adjusted	141	319.9	30.31	36	79.2	14.26	13	35.3	10.48	16	43.4	11.53	6	18.8	8.19	8	13.2	5.19
Vermont																		
Observed	23	237.9	55.07	8	70.6	26.78	-	-	-	-	-	-	-	-	-	-	-	-
Delay Adjusted	23	241.1	55.89	8	70.9	26.90	-	-	-	-	-	-	-	-	-	-	-	-
Virginia																		
Observed	16,864	389.6	3.06	5,844	132.7	1.77	2,015	47.6	1.08	1,710	39.9	0.98	953	21.6	0.71	541	12.3	0.54
Delay Adjusted	17,293	399.4	3.14	5,957	135.3	1.80	2,038	48.1	1.09	1,738	40.6	1.00	966	21.9	0.72	553	12.6	0.55
Washington																		
Observed	1,991	374.9	8.92	670	122.0	4.97	222	47.7	3.37	156	29.3	2.49	103	19.5	2.03	100	15.5	1.61
Delay Adjusted	2,040	384.1	9.14	676	123.2	5.02	224	48.2	3.40	159	29.8	2.52	104	19.7	2.06	102	15.8	1.64
Seattle																		
Observed	1,859	395.1	9.74	632	129.1	5.42	209	51.5	3.75	145	31.1	2.73	96	20.7	2.24	96	16.8	1.79
Delay Adjusted	1,900	403.7	9.95	636	129.9	5.45	210	51.8	3.77	147	31.5	2.77	97	20.8	2.26	97	17.0	1.81
West Virginia																		
Observed	645	375.1	15.14	201	119.3	8.65	91	51.4	5.49	64	37.1	4.76	36	19.1	3.29	25	15.4	3.16
Delay Adjusted	660	383.8	15.49	206	122.5	8.88	93	52.4	5.60	65	37.7	4.83	36	19.4	3.32	26	15.7	3.24
Wisconsin																		
Observed	3,471	469.5	8.34	1,003	130.7	4.28	522	76.4	3.49	308	42.5	2.53	208	27.8	2.01	130	15.3	1.40
Delay Adjusted	3,555	481.3	8.55	1,035	134.8	4.42	526	76.9	3.51	318	43.9	2.62	216	29.0	2.09	134	15.8	1.44

Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Delay Adjusted Counts and Rates[1] for the Five Most Commonly Diagnosed Cancers
Selected Areas in the NAACCR United States Combined[2], 2010-2014, Black, Females

	All Sites			Prostate			Lung & Bronchus			Colon & Rectum			Uterus Corpus & Uterus, NOS[3]			Thyroid		
	Count	*Rate*	*S.E.[3]*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
Wyoming																		
Observed	29	165.1	31.52	7	43.8	16.99	-	-	-	-	-	-	-	-	-	-	-	-
Delay Adjusted	29	167.4	31.99	7	44.2	17.14	-	-	-	-	-	-	-	-	-	-	-	-

1. Rates are per 100,000 population and are age-adjusted by five-year age groups to the 2000 U.S. standard population based on single years of age.
2. See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.
3. Abbreviations: NOS - Not Otherwise Specified; S.E. - Standard Error

^ There are insufficient historical registry data to calculate delay factors necessary for delay adjusted rates.

\- Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for All Sites combined.

Note: Delay adjusted counts are rounded to the nearest whole number. A small registry and/or rare cancer site may have the same delay adjusted count and observed count, but this is just due to rounding to whole counts. For more information about cancer incidence rates adjusted for reporting delay see https://surveillance.cancer.gov/delay/.

NAACCR U.S. Combined Delay Adjusted Incidence 2010-2014, Asian/Pacific Islander (API) Combined

UNITED STATES

V-43 Combined Delay Adjusted Incidence – Asian/Pacific Islander (API)

Average Annual Age-adjusted Cancer Incidence Delay Adjusted Counts and Rates*, Selected Sites
NAACCR United States Combined[1], 2010-2014, Asian/Pacific Islander (API)

	MALES (ALL RACES)						FEMALES (ALL RACES)					
	Observed			Delay Adjusted			Observed			Delay Adjusted		
SITES	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
All Sites (invasive & in situ bladder)	100,751	295.9	0.98	104,765	307.8	1.01	126,204	283.0	0.81	131,302	294.5	0.85
Oral Cavity & Pharynx	4,207	11.2	0.18	4,356	11.6	0.19	2,232	5.0	0.11	2,310	5.2	0.11
Digestive												
Esophagus	1,248	3.7	0.11	1,277	3.8	0.11	398	1.0	0.05	407	1.0	0.05
Stomach	4,553	14.1	0.22	4,641	14.3	0.22	3,376	8.1	0.14	3,443	8.2	0.14
Colon & Rectum	12,803	37.0	0.34	13,154	38.0	0.35	11,595	27.0	0.26	11,916	27.7	0.26
Liver & IBD[2]	7,146	20.0	0.25	7,359	20.6	0.25	3,145	7.6	0.14	3,241	7.8	0.14
Pancreas	3,190	10.0	0.18	3,266	10.2	0.19	3,522	8.7	0.15	3,605	8.9	0.15
Respiratory												
Larynx	740	2.2	0.08	752	2.2	0.09	123	0.3	0.03	125	0.3	0.03
Lung & bronchus	14,155	45.2	0.40	14,501	46.3	0.41	11,508	27.9	0.27	11,789	28.6	0.27
Skin												
Melanomas of the skin	534	1.5	0.07	565	1.6	0.07	530	1.2	0.05	562	1.3	0.06
Breast (female cases only)	-	-	-	-	-	-	42,016	90.7	0.45	43,660	94.2	0.47
Female Genital												
Cervix uteri	-	-	-	-	-	-	2,839	6.0	0.11	2,907	6.2	0.12
Corpus & uterus, NOS[2]	-	-	-	-	-	-	8,626	18.2	0.20	8,868	18.7	0.20
Ovary	-	-	-	-	-	-	4,212	9.2	0.14	4,385	9.5	0.15
Male Genital												
Prostate	19,972	58.4	0.43	21,223	62.1	0.46	-	-	-	-	-	-
Testis	893	1.9	0.06	908	1.9	0.07	-	-	-	-	-	-
Urinary												
Bladder (incl. *in situ*)	4,516	15.0	0.23	4,709	15.6	0.24	1,509	3.8	0.10	1,574	3.9	0.10
Kidney & renal pelvis	3,892	10.9	0.18	4,041	11.3	0.19	2,115	4.9	0.11	2,196	5.1	0.11
Brain & Other Nervous System	1,622	4.2	0.11	1,693	4.4	0.11	1,430	3.2	0.09	1,491	3.4	0.09
Endocrine												
Thyroid gland	2,793	6.9	0.13	2,879	7.1	0.14	10,254	21.3	0.21	10,568	21.9	0.22
Lymphomas												
Hodgkin Lymphoma	628	1.5	0.06	647	1.5	0.06	461	1.0	0.05	476	1.0	0.05
Non-Hodgkin Lymphoma	5,381	15.6	0.22	5,626	16.4	0.23	4,557	10.6	0.16	4,766	11.1	0.17
Multiple Myeloma	1,576	4.8	0.13	1,725	5.2	0.14	1,299	3.1	0.09	1,424	3.4	0.10
Leukemias	3,369	9.6	0.17	3,636	10.3	0.19	2,666	6.2	0.12	2,880	6.7	0.13

* Rates are per 100,000 population and are age-adjusted by five-year age groups to the 2000 U.S. standard population based on single years of age and to standard.

\- Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for All Sites combined.

[1] NAACCR United States Combined includes registries for which a U.S. average delay adjustment factor was calculated. See technical notes.

[2] Abbreviations: IBD – Intrahepatic Bile Duct; NOS - Not Otherwise Specified

Note: Delay adjusted counts are rounded to the nearest whole number. A small registry and/or rare cancer site may have the same delay adjusted count and observed count, but this is just due to rounding to whole counts. For more information about cancer incidence rates adjusted for reporting delay see https://surveillance.cancer.gov/delay/.

Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Delay Adjusted Counts and Rates[1] for the Five Most Commonly Diagnosed Cancers
Selected Areas in the NAACCR United States Combined[2], 2010-2014, Asian/Pacific Islander, Males

	All Sites			Prostate			Lung & Bronchus			Colon & Rectum			Liver & IBD[3]			Non-Hodgkin Lymphoma		
	Count	*Rate*	*S.E.*[3]	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
NAACCR U.S. Combined																		
Observed	100,751	295.9	0.98	19,972	58.4	0.43	14,155	45.2	0.40	12,803	37.0	0.34	7,146	20.0	0.25	5,381	15.6	0.22
Delay Adjusted	104,765	307.8	1.01	21,223	62.1	0.46	14,501	46.3	0.41	13,154	38.0	0.35	7,359	20.6	0.25	5,626	16.4	0.23
Alabama																		
Observed	269	292.2	21.29	67	72.1	9.96	29	34.2	7.61	21	17.8	4.41	23	27.1	6.30	13	10.5	3.04
Delay Adjusted	274	298.0	21.72	70	75.4	10.46	29	34.4	7.65	21	18.0	4.45	23	27.5	6.39	13	10.7	3.11
Alaska																		
Observed	291	351.0	24.12	57	73.6	10.94	38	55.8	10.38	45	51.7	8.96	21	23.7	6.16	19	27.2	7.55
Delay Adjusted	303	365.2	25.10	60	78.2	11.66	39	57.1	10.62	46	53.3	9.21	22	24.7	6.43	20	28.7	7.97
Arizona																		
Observed	842	225.1	8.44	150	40.4	3.54	122	36.5	3.58	105	28.4	3.03	59	14.4	1.99	41	10.8	1.82
Delay Adjusted	909	243.1	9.11	169	45.6	4.00	125	37.4	3.66	110	29.8	3.18	61	14.8	2.06	44	11.5	1.93
Arkansas																		
Observed	403	627.1	37.21	56	96.7	15.29	45	80.3	13.21	46	61.5	10.28	29	36.7	7.54	18	29.8	8.92
Delay Adjusted	423	657.0	38.95	59	101.2	15.94	46	81.9	13.46	47	62.9	10.58	29	37.1	7.63	19	31.1	9.36
California																		
Observed	38,384	299.8	1.57	7,730	59.0	0.69	5,608	46.2	0.63	5,132	39.5	0.57	2,804	21.0	0.41	2,121	16.6	0.37
Delay Adjusted	40,352	315.2	1.65	8,258	63.0	0.74	5,825	47.9	0.66	5,304	40.8	0.58	2,920	21.9	0.42	2,237	17.5	0.39
Greater California																		
Observed	13,395	280.2	2.51	2,664	54.8	1.10	1,977	44.3	1.03	1,743	35.9	0.89	1,004	19.8	0.65	706	14.8	0.57
Delay Adjusted	14,276	298.8	2.67	2,897	59.7	1.20	2,070	46.4	1.08	1,822	37.5	0.93	1,043	20.6	0.67	748	15.6	0.61
Greater Bay																		
Observed	13,547	321.5	2.82	2,911	68.4	1.30	2,011	50.3	1.15	1,708	39.8	0.98	1,003	22.7	0.74	786	18.5	0.67
Delay Adjusted	14,213	337.3	2.96	3,103	72.9	1.38	2,083	52.1	1.19	1,764	41.1	1.02	1,045	23.7	0.77	832	19.6	0.71
Los Angeles																		
Observed	11,431	299.9	2.86	2,151	53.8	1.18	1,620	43.8	1.11	1,681	43.7	1.08	795	20.5	0.74	628	16.6	0.68
Delay Adjusted	11,850	310.9	2.96	2,253	56.4	1.24	1,672	45.2	1.14	1,719	44.6	1.11	830	21.4	0.77	656	17.4	0.71
Colorado																		
Observed	719	259.9	10.43	124	47.4	4.50	103	43.3	4.54	80	29.2	3.50	51	16.6	2.49	48	17.4	2.73
Delay Adjusted	736	266.1	10.69	134	51.3	4.88	104	43.6	4.57	81	29.6	3.54	51	16.7	2.50	50	18.1	2.84
Connecticut																		
Observed	633	243.3	11.01	99	37.6	3.99	89	36.3	4.31	68	24.1	3.24	50	19.6	3.13	34	13.8	2.83
Delay Adjusted	651	250.4	11.35	103	39.4	4.19	90	36.8	4.36	69	24.4	3.28	51	20.0	3.19	36	14.4	2.95

Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Delay Adjusted Counts and Rates[1] for the Five Most Commonly Diagnosed Cancers
Selected Areas in the NAACCR United States Combined[2], 2010-2014, Asian/Pacific Islander, Males

	All Sites			Prostate			Lung & Bronchus			Colon & Rectum			Liver & IBD[3]			Non-Hodgkin Lymphoma		
	Count	*Rate*	*S.E.[3]*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
Delaware																		
Observed	152	254.8	22.46	35	55.5	9.66	16	33.9	9.44	13	20.1	5.87	10	16.8	5.57	10	18.6	6.61
Delay Adjusted	156	262.4	23.15	37	58.2	10.16	16	34.8	9.72	14	21.0	6.13	10	17.5	5.80	11	19.6	6.96
District of Columbia																		
Observed	142	453.3	41.02	21	72.3	16.47	17	62.6	15.77	19	64.0	15.80	10	30.6	10.28	6	18.5	8.17
Delay Adjusted	146	466.8	42.26	22	74.5	16.97	17	63.9	16.07	20	65.9	16.29	10	31.0	10.41	6	18.9	8.34
Florida																		
Observed	2,479	215.2	4.57	510	43.4	2.01	323	30.4	1.80	316	26.5	1.58	138	11.5	1.02	141	12.0	1.06
Delay Adjusted	2,598	225.2	4.78	549	46.8	2.17	328	30.9	1.83	327	27.4	1.62	141	11.7	1.04	150	12.7	1.13
Georgia																		
Observed	1,600	262.7	7.61	285	50.2	3.33	181	32.3	2.84	226	36.0	2.77	129	20.3	2.06	82	12.8	1.68
Delay Adjusted	1,681	276.3	8.02	303	53.3	3.55	184	32.9	2.90	230	36.6	2.82	130	20.5	2.08	85	13.2	1.74
Hawaii																		
Observed	10,135	390.8	3.94	2,251	85.8	1.84	1,511	58.0	1.52	1,452	55.3	1.48	487	18.5	0.85	519	20.4	0.91
Delay Adjusted	10,301	397.2	4.01	2,313	88.2	1.89	1,525	58.6	1.53	1,468	55.9	1.49	500	19.0	0.87	534	21.0	0.94
Idaho																		
Observed	107	289.2	30.67	19	46.2	11.25	14	52.1	14.53	13	33.5	10.27	7	13.2	5.29	6	15.9	6.99
Delay Adjusted	119	321.6	34.12	20	48.4	11.78	14	53.3	14.88	13	34.0	10.43	7	13.8	5.49	6	16.2	7.13
Illinois																		
Observed	3,586	279.1	5.06	741	56.5	2.24	486	41.2	2.02	466	36.2	1.82	198	15.2	1.17	182	14.7	1.18
Delay Adjusted	3,665	285.3	5.17	774	59.2	2.35	493	41.8	2.05	472	36.6	1.84	201	15.4	1.19	189	15.3	1.23
Indiana																		
Observed	436	277.5	15.91	65	43.7	6.14	52	41.9	6.77	48	25.2	4.07	28	18.6	4.18	24	10.7	2.38
Delay Adjusted	456	290.4	16.65	69	46.9	6.64	53	42.7	6.91	50	26.1	4.22	29	19.3	4.32	25	11.2	2.47
Iowa																		
Observed	256	343.3	26.07	35	51.4	10.06	29	51.6	11.08	32	45.0	9.46	37	52.2	10.42	15	19.5	5.83
Delay Adjusted	265	354.7	26.93	37	55.0	10.73	29	52.1	11.19	32	45.4	9.54	38	53.2	10.60	15	20.0	5.99
Kansas																		
Observed	359	301.7	18.38	61	59.1	8.37	42	38.7	6.84	37	27.2	4.88	32	23.3	4.38	25	17.3	4.09
Delay Adjusted	^	^	^	^	^	^	^	^	^	^	^	^	^	^	^	^	^	^

1. Rates are per 100,000 population and are age-adjusted by five-year age groups to the 2000 U.S. standard population based on single years of age.
2. See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.
3. Abbreviations: IBD - Intrahepatic Bile Duct; S.E. - Standard Error

^ There are insufficient historical registry data to calculate delay factors necessary for delay adjusted rates.

\- Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for All Sites combined.

Note: Delay adjusted counts are rounded to the nearest whole number. A small registry and/or rare cancer site may have the same delay adjusted count and observed count, but this is just due to rounding to whole counts. For more information about cancer incidence rates adjusted for reporting delay see https://surveillance.cancer.gov/delay/.

Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Delay Adjusted Counts and Rates[1] for the Five Most Commonly Diagnosed Cancers
Selected Areas in the NAACCR United States Combined[2], 2010-2014, Asian/Pacific Islander, Males

	All Sites			Prostate			Lung & Bronchus			Colon & Rectum			Liver & IBD[3]			Non-Hodgkin Lymphoma		
	Count	*Rate*	*S.E.*[3]	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
Kentucky																		
Observed	183	217.2	18.98	30	48.3	9.83	23	28.9	7.50	24	26.2	6.30	13	15.6	5.38	10	11.7	4.68
Delay Adjusted	187	222.2	19.46	31	50.5	10.30	23	29.2	7.54	24	26.7	6.46	13	16.1	5.64	10	12.1	4.83
Louisiana																		
Observed	461	310.2	16.21	80	60.4	7.38	75	52.6	6.88	60	41.8	6.07	50	28.3	4.35	15	7.6	2.02
Delay Adjusted	468	314.9	16.45	82	61.8	7.56	76	53.2	6.96	61	42.3	6.15	51	28.8	4.43	15	7.7	2.04
Maine																		
Observed	50	209.8	33.33	9	37.9	13.69	-	-	-	6	18.0	7.38	6	24.3	11.16	6	28.6	13.02
Delay Adjusted	51	214.8	34.16	9	38.8	14.00	-	-	-	6	18.1	7.43	6	25.4	11.72	6	29.0	13.18
Maryland																		
Observed	1,920	258.1	6.27	403	52.1	2.73	235	35.3	2.46	208	27.8	2.05	130	17.0	1.58	100	13.3	1.39
Delay Adjusted	2,002	269.2	6.55	438	56.8	2.98	241	36.2	2.52	216	28.9	2.12	135	17.6	1.64	106	14.0	1.47
Massachusetts																		
Observed	2,082	303.5	7.10	362	55.7	3.06	342	56.9	3.25	235	33.4	2.31	199	27.7	2.07	81	10.7	1.26
Delay Adjusted	2,131	310.8	7.27	380	58.5	3.21	348	57.9	3.31	244	34.6	2.39	202	28.1	2.11	82	10.8	1.28
Michigan																		
Observed	1,185	260.0	8.38	313	68.5	4.14	126	31.0	3.08	150	36.0	3.30	62	12.2	1.69	65	13.0	1.78
Delay Adjusted	1,234	270.8	8.73	321	70.3	4.26	127	31.3	3.11	153	36.6	3.36	63	12.4	1.72	67	13.3	1.83
Detroit																		
Observed	826	284.3	10.93	247	83.9	5.72	86	32.3	3.88	96	34.5	3.95	39	12.5	2.18	40	11.9	2.06
Delay Adjusted	872	300.1	11.54	253	86.1	5.87	87	32.5	3.90	97	34.9	4.01	39	12.6	2.18	40	12.1	2.09
Mississippi																		
Observed	151	311.1	28.73	27	62.5	13.79	27	54.9	11.35	18	37.3	9.66	15	30.6	9.06	6	14.0	6.58
Delay Adjusted	154	318.1	29.37	28	64.8	14.30	27	55.5	11.48	18	37.7	9.75	15	31.5	9.31	6	14.9	7.04
Missouri																		
Observed	487	268.1	13.47	75	42.4	5.27	72	45.6	5.86	61	35.6	5.11	45	22.4	3.57	27	14.9	3.34
Delay Adjusted	509	280.5	14.11	80	45.4	5.67	73	46.3	5.96	62	36.1	5.19	46	22.8	3.64	28	15.4	3.43
Montana																		
Observed	31	272.3	52.96	7	48.9	19.27	-	-	-	-	-	-	-	-	-	-	-	-
Delay Adjusted	32	284.7	55.38	7	51.8	20.36	-	-	-	-	-	-	-	-	-	-	-	-
Nebraska																		
Observed	168	323.4	28.33	21	50.9	11.98	23	46.8	10.76	22	38.8	9.26	20	36.7	9.08	9	19.1	7.37
Delay Adjusted	174	335.6	29.42	23	54.7	12.88	24	48.0	11.06	22	39.5	9.43	21	38.5	9.44	9	20.1	7.75

Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Delay Adjusted Counts and Rates[1] for the Five Most Commonly Diagnosed Cancers
Selected Areas in the NAACCR United States Combined[2], 2010-2014, Asian/Pacific Islander, Males

	All Sites			Prostate			Lung & Bronchus			Colon & Rectum			Liver & IBD[3]			Non-Hodgkin Lymphoma		
	Count	*Rate*	*S.E.[3]*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
New Hampshire																		
Observed	108	238.5	26.46	28	64.0	13.68	12	31.9	10.00	10	21.4	7.38	6	11.0	5.34	-	-	-
Delay Adjusted	112	247.1	27.46	29	66.0	14.12	12	32.4	10.13	10	21.7	7.46	6	11.5	5.56	-	-	-
New Jersey																		
Observed	4,234	266.7	4.52	1,004	61.8	2.11	468	33.4	1.69	469	28.3	1.44	223	14.1	1.05	219	13.2	0.98
Delay Adjusted	4,447	280.1	4.75	1,096	67.6	2.32	485	34.6	1.75	495	29.9	1.52	228	14.4	1.07	234	14.1	1.04
New York																		
Observed	11,558	335.4	3.30	2,160	61.2	1.39	1,651	52.8	1.38	1,325	38.4	1.12	831	22.7	0.83	550	15.8	0.71
Delay Adjusted	11,834	343.4	3.38	2,313	65.6	1.49	1,667	53.3	1.39	1,351	39.1	1.14	850	23.2	0.85	568	16.3	0.74
North Carolina																		
Observed	901	247.6	9.58	188	52.4	4.16	127	38.6	3.94	88	24.0	3.00	50	13.3	2.17	53	13.6	2.22
Delay Adjusted	939	258.2	10.00	200	55.6	4.42	128	39.0	3.99	91	24.8	3.10	51	13.6	2.22	55	14.1	2.30
North Dakota																		
Observed	28	337.5	78.47	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Delay Adjusted	29	347.5	80.78	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Ohio																		
Observed	874	239.3	9.00	202	54.5	4.11	119	38.5	3.92	94	26.9	3.13	58	14.5	2.10	39	10.4	1.87
Delay Adjusted	939	257.3	9.68	215	58.0	4.37	125	40.5	4.12	97	27.7	3.21	61	15.2	2.19	41	10.9	1.96
Oklahoma																		
Observed	426	359.5	19.64	72	61.1	7.90	62	56.3	7.85	68	56.7	8.00	31	26.7	5.31	22	16.8	4.06
Delay Adjusted	432	364.5	19.91	75	63.6	8.23	63	56.8	7.92	69	57.6	8.11	32	27.7	5.52	22	17.0	4.10
Oregon																		
Observed	959	307.2	10.56	157	50.8	4.26	142	51.4	4.54	112	38.8	3.89	106	32.5	3.34	50	16.0	2.43
Delay Adjusted	1,009	323.4	11.13	170	55.0	4.60	145	52.4	4.63	117	40.6	4.07	108	33.2	3.42	52	16.5	2.52
Pennsylvania																		
Observed	1,854	264.8	6.75	339	51.0	2.99	256	42.1	2.89	200	27.9	2.14	150	19.9	1.76	107	14.7	1.53
Delay Adjusted	1,921	274.5	7.00	368	55.3	3.24	259	42.5	2.92	204	28.5	2.19	154	20.5	1.81	111	15.2	1.59
Rhode Island																		
Observed	152	268.6	24.31	29	52.3	10.48	16	32.4	9.11	14	26.9	8.00	17	29.1	7.63	11	22.1	7.41
Delay Adjusted	156	276.0	24.96	32	56.9	11.38	16	32.7	9.19	14	27.2	8.09	17	29.6	7.74	11	22.9	7.69

1. Rates are per 100,000 population and are age-adjusted by five-year age groups to the 2000 U.S. standard population based on single years of age.
2. See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.
3. Abbreviations: IBD - Intrahepatic Bile Duct; S.E. - Standard Error

^ There are insufficient historical registry data to calculate delay factors necessary for delay adjusted rates.

\- Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for All Sites combined.

Note: Delay adjusted counts are rounded to the nearest whole number. A small registry and/or rare cancer site may have the same delay adjusted count and observed count, but this is just due to rounding to whole counts. For more information about cancer incidence rates adjusted for reporting delay see https://surveillance.cancer.gov/delay/.

Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Delay Adjusted Counts and Rates[1] for the Five Most Commonly Diagnosed Cancers
Selected Areas in the NAACCR United States Combined[2], 2010-2014, Asian/Pacific Islander, Males

	All Sites			Prostate			Lung & Bronchus			Colon & Rectum			Liver & IBD[3]			Non-Hodgkin Lymphoma		
	Count	*Rate*	*S.E.*[3]	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
South Carolina																		
Observed	299	238.6	16.08	53	47.3	7.31	52	42.1	6.79	36	28.9	5.53	11	8.7	2.98	25	21.7	5.58
Delay Adjusted	308	246.3	16.66	55	49.5	7.68	54	43.5	7.03	37	29.8	5.68	11	9.1	3.08	26	23.2	6.00
South Dakota																		
Observed	29	314.1	69.54	-	-	-	6	87.7	42.23	6	35.5	15.84	-	-	-	-	-	-
Delay Adjusted	30	329.7	72.88	-	-	-	6	90.3	43.35	6	37.1	16.68	-	-	-	-	-	-
Tennessee																		
Observed	484	290.0	15.48	97	64.0	7.22	54	36.7	5.91	62	37.9	5.61	37	19.6	3.53	33	20.1	4.28
Delay Adjusted	498	298.2	15.92	100	65.8	7.43	55	37.3	6.01	63	38.6	5.71	38	20.2	3.67	34	20.9	4.46
Texas																		
Observed	4,668	239.8	3.99	816	44.1	1.72	644	37.9	1.69	580	29.4	1.40	444	21.3	1.13	270	13.2	0.91
Delay Adjusted	4,925	253.0	4.21	887	48.1	1.88	657	38.6	1.72	600	30.4	1.44	452	21.7	1.15	290	14.2	0.99
Utah																		
Observed	463	337.9	17.40	118	94.2	9.28	38	34.6	6.04	42	30.9	5.22	32	22.5	4.41	23	15.7	3.74
Delay Adjusted	488	356.1	18.34	121	96.7	9.52	39	35.1	6.13	42	31.1	5.26	33	23.5	4.64	23	15.9	3.78
Vermont																		
Observed	35	299.5	56.93	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Delay Adjusted	35	303.1	57.63	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Virginia																		
Observed	2,182	251.5	6.02	391	43.4	2.38	308	39.5	2.50	263	30.2	2.12	154	16.9	1.50	110	12.3	1.31
Delay Adjusted	2,237	257.7	6.17	402	44.7	2.45	312	39.9	2.53	267	30.6	2.15	159	17.3	1.54	113	12.7	1.35
Washington																		
Observed	3,403	315.2	5.75	586	55.3	2.41	489	52.1	2.48	448	39.7	2.00	282	24.2	1.53	196	17.6	1.34
Delay Adjusted	3,487	323.0	5.89	613	57.9	2.52	495	52.7	2.51	455	40.3	2.02	286	24.6	1.55	200	18.0	1.37
Seattle																		
Observed	3,188	328.8	6.19	558	59.1	2.63	457	53.6	2.64	423	41.8	2.16	258	24.9	1.64	189	18.7	1.44
Delay Adjusted	3,258	336.0	6.32	582	61.6	2.74	461	54.1	2.67	429	42.3	2.19	259	25.1	1.65	192	19.0	1.47
West Virginia																		
Observed	52	189.7	30.60	10	30.9	10.06	13	43.2	13.36	11	44.3	15.30	-	-	-	-	-	-
Delay Adjusted	53	193.8	31.23	10	31.4	10.23	13	43.8	13.50	11	45.2	15.58	-	-	-	-	-	-
Wisconsin																		
Observed	514	305.8	15.68	77	58.9	7.33	54	42.2	6.36	62	32.8	4.80	50	28.8	4.66	36	18.8	3.82
Delay Adjusted	517	307.4	15.76	83	63.5	7.88	54	42.5	6.41	64	34.0	4.99	50	29.1	4.71	36	19.0	3.85

Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Delay Adjusted Counts and Rates[1] for the Five Most Commonly Diagnosed Cancers
Selected Areas in the NAACCR United States Combined[2], 2010-2014, Asian/Pacific Islander, Males

	All Sites			Prostate			Lung & Bronchus			Colon & Rectum			Liver & IBD[3]			Non-Hodgkin Lymphoma		
	Count	*Rate*	*S.E.*[3]	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
Wyoming																		
Observed	17	154.5	38.94	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Delay Adjusted	17	156.3	39.42	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-

1. Rates are per 100,000 population and are age-adjusted by five-year age groups to the 2000 U.S. standard population based on single years of age.
2. See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.
3. Abbreviations: IBD - Intrahepatic Bile Duct; S.E. - Standard Error

^ There are insufficient historical registry data to calculate delay factors necessary for delay adjusted rates.

\- Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for All Sites combined.

Note: Delay adjusted counts are rounded to the nearest whole number. A small registry and/or rare cancer site may have the same delay adjusted count and observed count, but this is just due to rounding to whole counts. For more information about cancer incidence rates adjusted for reporting delay see https://surveillance.cancer.gov/delay/.

Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Delay Adjusted Counts and Rates[1] for the Five Most Commonly Diagnosed Cancers
Selected Areas in the NAACCR United States Combined[2], 2010-2014, Asian/Pacific Islander, Females

	All Sites			Breast			Colon & Rectum			Lung & Bronchus			Thyroid			Uterus Corpus & Uterus, NOS[3]		
	Count	*Rate*	*S.E.*[3]	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
NAACCR U.S. Combined																		
Observed	126,204	283.0	0.81	42,016	90.7	0.45	11,595	27.0	0.26	11,508	27.9	0.27	10,254	21.3	0.21	8,626	18.2	0.20
Delay Adjusted	131,302	294.5	0.85	43,660	94.2	0.47	11,916	27.7	0.26	11,789	28.6	0.27	10,568	21.9	0.22	8,868	18.7	0.20
Alabama																		
Observed	399	280.2	15.32	136	87.4	8.03	33	24.6	4.70	42	35.5	5.88	24	12.5	2.57	15	9.3	2.55
Delay Adjusted	407	285.9	15.64	138	88.8	8.16	33	24.8	4.73	42	35.7	5.92	24	12.6	2.59	15	9.4	2.57
Alaska																		
Observed	369	295.5	16.15	120	92.3	8.77	28	22.6	4.48	43	38.4	6.19	33	25.0	4.46	34	23.3	4.14
Delay Adjusted	385	308.0	16.83	121	93.0	8.84	29	23.2	4.61	44	39.3	6.36	34	25.6	4.55	35	24.2	4.30
Arizona																		
Observed	1,395	252.3	7.09	450	75.6	3.68	99	19.9	2.10	150	30.8	2.65	134	21.0	1.86	85	13.5	1.52
Delay Adjusted	1,508	272.7	7.66	474	79.6	3.88	104	20.9	2.21	154	31.5	2.72	141	22.2	1.97	89	14.2	1.59
Arkansas																		
Observed	498	528.5	25.96	119	126.9	12.67	48	53.5	8.54	51	62.1	9.47	26	24.5	5.36	25	24.4	5.30
Delay Adjusted	522	553.7	27.19	125	133.1	13.29	49	54.5	8.73	52	63.4	9.67	27	25.1	5.51	26	25.0	5.44
California																		
Observed	48,091	289.1	1.33	16,207	95.7	0.76	4,796	29.0	0.42	4,325	26.6	0.41	3,169	19.1	0.34	3,467	19.9	0.34
Delay Adjusted	50,557	303.9	1.40	17,076	100.8	0.80	4,955	29.9	0.44	4,494	27.7	0.43	3,278	19.8	0.35	3,619	20.7	0.36
Greater California																		
Observed	17,420	277.5	2.13	5,743	88.5	1.18	1,709	27.9	0.68	1,552	26.0	0.67	1,271	19.7	0.56	1,221	18.3	0.53
Delay Adjusted	18,557	295.7	2.27	6,139	94.7	1.26	1,785	29.1	0.72	1,627	27.3	0.70	1,332	20.6	0.58	1,299	19.5	0.57
Greater Bay																		
Observed	16,185	298.7	2.37	5,585	101.4	1.37	1,516	28.0	0.73	1,503	28.5	0.74	936	17.2	0.57	1,190	21.0	0.62
Delay Adjusted	16,984	313.5	2.49	5,876	106.6	1.44	1,564	28.9	0.75	1,557	29.5	0.77	964	17.8	0.59	1,236	21.8	0.64
Los Angeles																		
Observed	14,479	293.9	2.48	4,876	98.8	1.44	1,571	31.2	0.80	1,269	25.5	0.72	962	20.6	0.68	1,057	20.8	0.65
Delay Adjusted	15,008	304.6	2.57	5,057	102.5	1.49	1,606	31.9	0.82	1,309	26.3	0.75	982	21.0	0.69	1,085	21.3	0.67
Colorado																		
Observed	1,035	248.6	8.01	339	76.1	4.25	99	27.5	2.85	106	28.5	2.86	112	22.6	2.18	66	14.1	1.77
Delay Adjusted	1,058	254.5	8.20	343	77.0	4.30	100	27.8	2.88	107	28.7	2.88	113	22.9	2.21	67	14.2	1.79
Connecticut																		
Observed	835	255.5	9.68	314	89.2	5.41	69	25.6	3.38	64	26.8	3.60	97	23.3	2.54	40	11.4	1.90
Delay Adjusted	860	263.2	9.98	338	96.1	5.82	70	26.0	3.43	65	27.2	3.66	99	23.8	2.59	40	11.5	1.92

Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Delay Adjusted Counts and Rates[1] for the Five Most Commonly Diagnosed Cancers
Selected Areas in the NAACCR United States Combined[2], 2010-2014, Asian/Pacific Islander, Females

	All Sites			Breast			Colon & Rectum			Lung & Bronchus			Thyroid			Uterus Corpus & Uterus, NOS[3]		
	Count	*Rate*	*S.E.[3]*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
Delaware																		
Observed	214	274.1	19.87	74	85.9	10.19	10	15.7	5.32	17	26.6	6.93	29	31.4	5.96	10	12.5	4.16
Delay Adjusted	220	282.1	20.46	75	87.4	10.37	10	16.4	5.56	17	27.1	7.09	30	32.3	6.13	10	12.6	4.20
District of Columbia																		
Observed	181	333.1	26.29	52	90.4	13.15	13	24.0	6.93	17	35.2	9.05	28	34.5	7.18	9	16.2	5.48
Delay Adjusted	187	343.6	27.12	55	95.7	13.92	13	24.6	7.07	17	35.9	9.24	29	36.4	7.56	9	16.8	5.69
Florida																		
Observed	3,440	219.7	3.88	1,165	70.1	2.10	284	18.6	1.15	323	22.3	1.29	260	15.5	0.98	237	14.0	0.94
Delay Adjusted	3,608	230.3	4.06	1,214	73.1	2.19	293	19.2	1.18	327	22.6	1.31	262	15.6	0.98	245	14.5	0.97
Georgia																		
Observed	1,960	244.2	6.08	662	74.4	3.09	171	23.9	2.04	137	20.3	1.89	165	16.4	1.32	107	12.2	1.25
Delay Adjusted	2,057	256.5	6.39	669	75.2	3.12	174	24.3	2.07	140	20.7	1.93	168	16.7	1.35	108	12.4	1.26
Hawaii																		
Observed	11,730	377.0	3.59	4,015	132.1	2.14	1,177	36.3	1.09	1,207	35.9	1.06	621	23.9	0.98	942	32.1	1.07
Delay Adjusted	11,926	383.3	3.65	4,083	134.4	2.18	1,191	36.7	1.11	1,218	36.2	1.07	631	24.3	1.00	954	32.6	1.09
Idaho																		
Observed	165	276.0	22.48	50	76.0	11.13	13	22.3	6.47	18	36.2	8.90	9	13.3	4.58	13	19.0	5.39
Delay Adjusted	184	307.3	25.04	51	77.1	11.29	13	22.6	6.54	18	36.9	9.07	9	13.4	4.61	13	19.4	5.51
Illinois																		
Observed	4,454	268.1	4.17	1,641	92.7	2.35	394	25.7	1.35	324	21.8	1.26	406	21.7	1.10	324	18.2	1.04
Delay Adjusted	4,551	274.0	4.26	1,705	96.4	2.44	399	26.0	1.37	329	22.1	1.28	415	22.2	1.13	328	18.4	1.05
Indiana																		
Observed	565	244.4	11.29	170	66.1	5.44	54	29.7	4.43	48	25.2	3.95	43	13.9	2.24	42	16.8	2.76
Delay Adjusted	593	256.3	11.85	176	68.6	5.66	56	30.7	4.56	49	25.7	4.04	44	14.3	2.29	43	17.1	2.80
Iowa																		
Observed	267	250.9	17.10	80	68.2	8.07	19	20.0	5.11	23	25.8	6.02	22	13.1	3.06	19	15.7	3.87
Delay Adjusted	276	259.9	17.74	84	71.4	8.44	19	20.1	5.15	23	26.3	6.14	22	13.2	3.10	19	15.8	3.89
Kansas																		
Observed	442	268.8	13.87	115	63.2	6.22	40	26.8	4.61	61	46.6	6.37	57	26.6	3.65	29	17.5	3.51
Delay Adjusted	^	^	^	^	^	^	^	^	^	^	^	^	^	^	^	^	^	^

1. Rates are per 100,000 population and are age-adjusted by five-year age groups to the 2000 U.S. standard population based on single years of age.
2. See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.
3. Abbreviations: NOS - Not Otherwise Specified; S.E. - Standard Error

^ There are insufficient historical registry data to calculate delay factors necessary for delay adjusted rates.

\- Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for All Sites combined.

Note: Delay adjusted counts are rounded to the nearest whole number. A small registry and/or rare cancer site may have the same delay adjusted count and observed count, but this is just due to rounding to whole counts. For more information about cancer incidence rates adjusted for reporting delay see https://surveillance.cancer.gov/delay/.

Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Delay Adjusted Counts and Rates[1] for the Five Most Commonly Diagnosed Cancers
Selected Areas in the NAACCR United States Combined[2], 2010-2014, Asian/Pacific Islander, Females

	All Sites			Breast			Colon & Rectum			Lung & Bronchus			Thyroid			Uterus Corpus & Uterus, NOS[3]		
	Count	*Rate*	*S.E.*[3]	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
Kentucky																		
Observed	303	254.4	16.00	89	64.4	7.21	34	35.1	6.56	36	39.8	7.16	27	17.8	3.52	22	16.1	3.56
Delay Adjusted	310	260.6	16.39	90	65.2	7.29	35	35.7	6.68	36	40.3	7.27	27	18.1	3.57	22	16.4	3.61
Louisiana																		
Observed	461	253.3	12.60	117	62.6	6.10	30	17.8	3.52	47	30.5	4.74	40	17.2	2.76	32	14.0	2.49
Delay Adjusted	468	257.3	12.81	118	63.2	6.14	30	18.1	3.56	48	30.8	4.80	41	17.5	2.80	32	14.1	2.51
Maine																		
Observed	102	280.8	30.59	28	76.3	15.97	8	24.5	9.06	13	50.0	14.94	19	38.8	8.96	6	16.6	7.19
Delay Adjusted	104	287.3	31.31	29	77.7	16.24	8	24.6	9.14	13	50.5	15.11	19	39.1	9.02	6	16.8	7.26
Maryland																		
Observed	2,485	261.1	5.43	861	84.3	2.95	212	24.1	1.73	185	22.1	1.69	264	25.4	1.59	160	15.6	1.27
Delay Adjusted	2,591	272.4	5.67	889	87.0	3.04	221	25.1	1.80	189	22.7	1.73	269	25.8	1.62	163	15.9	1.30
Massachusetts																		
Observed	2,521	289.6	6.02	816	87.7	3.17	221	27.7	1.95	244	32.6	2.17	280	27.0	1.67	118	12.9	1.22
Delay Adjusted	2,581	296.7	6.17	824	88.6	3.20	229	28.8	2.02	249	33.3	2.21	283	27.3	1.69	119	13.0	1.23
Michigan																		
Observed	1,497	260.9	7.18	534	85.3	3.86	104	19.6	2.06	152	31.5	2.71	110	15.9	1.58	106	17.0	1.72
Delay Adjusted	1,555	271.0	7.46	558	89.2	4.04	106	19.9	2.09	154	31.9	2.74	112	16.2	1.61	107	17.1	1.74
Detroit																		
Observed	954	274.3	9.42	358	94.8	5.24	68	21.4	2.78	80	27.9	3.30	78	19.6	2.31	67	17.9	2.30
Delay Adjusted	1,006	289.3	9.94	380	100.6	5.56	69	21.7	2.82	80	28.1	3.32	79	19.9	2.35	67	18.0	2.31
Mississippi																		
Observed	189	273.0	21.74	47	60.6	9.41	17	26.7	7.21	28	53.4	10.83	17	17.3	4.23	9	9.3	3.12
Delay Adjusted	193	279.5	22.27	48	62.1	9.65	17	27.1	7.30	28	54.2	10.99	17	17.7	4.31	9	9.3	3.13
Missouri																		
Observed	732	291.4	11.54	238	88.2	6.03	60	28.3	3.93	72	33.8	4.20	61	19.0	2.53	40	14.2	2.33
Delay Adjusted	767	305.5	12.11	243	89.8	6.15	61	28.8	4.00	73	34.3	4.26	63	19.6	2.61	40	14.3	2.35
Montana																		
Observed	77	341.3	40.70	27	112.5	22.63	6	29.0	12.30	9	41.4	14.42	-	-	-	9	35.5	12.20
Delay Adjusted	80	355.6	42.45	28	116.3	23.40	6	29.6	12.58	9	42.3	14.73	-	-	-	9	36.4	12.50
Nebraska																		
Observed	196	257.2	20.06	51	65.6	9.82	18	24.6	6.24	18	31.2	7.95	30	31.7	6.34	11	15.0	4.95
Delay Adjusted	203	267.0	20.82	52	66.4	9.95	18	25.3	6.42	18	31.8	8.07	31	32.3	6.46	11	15.2	5.00

Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Delay Adjusted Counts and Rates[1] for the Five Most Commonly Diagnosed Cancers
Selected Areas in the NAACCR United States Combined[2], 2010-2014, Asian/Pacific Islander, Females

	All Sites			Breast			Colon & Rectum			Lung & Bronchus			Thyroid			Uterus Corpus & Uterus, NOS[3]		
	Count	*Rate*	*S.E.[3]*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
New Hampshire																		
Observed	166	244.2	21.14	55	72.1	10.50	16	24.6	6.88	12	21.0	6.78	17	18.5	4.54	15	24.4	6.89
Delay Adjusted	172	253.4	21.96	56	73.0	10.64	16	24.9	6.96	12	21.7	7.04	17	18.8	4.62	15	24.8	6.99
New Jersey																		
Observed	5,190	264.0	3.86	1,938	92.2	2.17	401	22.3	1.18	377	22.5	1.22	509	22.3	1.01	352	16.4	0.91
Delay Adjusted	5,455	277.6	4.06	2,020	96.1	2.26	423	23.5	1.25	392	23.3	1.27	525	23.0	1.04	366	17.0	0.94
New York																		
Observed	13,166	310.2	2.78	4,076	92.2	1.47	1,106	27.5	0.85	1,173	30.3	0.91	1,567	33.5	0.86	858	19.2	0.67
Delay Adjusted	13,481	317.6	2.84	4,140	93.6	1.49	1,128	28.1	0.87	1,184	30.6	0.92	1,627	34.8	0.89	867	19.4	0.68
North Carolina																		
Observed	1,285	253.5	7.79	434	77.7	4.03	94	19.8	2.25	126	30.3	2.91	109	15.9	1.62	81	14.9	1.78
Delay Adjusted	1,338	264.0	8.12	488	87.4	4.52	97	20.4	2.32	127	30.7	2.95	112	16.3	1.66	82	15.1	1.80
North Dakota																		
Observed	39	291.9	55.10	-	-	-	-	-	-	-	-	-	9	51.7	21.75	-	-	-
Delay Adjusted	40	301.9	57.12	-	-	-	-	-	-	-	-	-	9	53.3	22.46	-	-	-
Ohio																		
Observed	1,161	236.0	7.34	419	79.8	4.07	85	19.0	2.20	109	26.5	2.68	119	19.6	1.85	58	11.0	1.50
Delay Adjusted	1,244	253.0	7.88	456	86.8	4.43	87	19.6	2.25	114	27.8	2.80	123	20.2	1.91	60	11.3	1.54
Oklahoma																		
Observed	559	322.2	14.67	159	80.5	6.63	62	41.9	5.74	65	44.9	5.95	29	16.0	3.30	44	22.8	3.61
Delay Adjusted	567	326.9	14.89	164	82.9	6.83	63	42.5	5.83	66	45.3	6.00	29	16.1	3.32	44	23.0	3.64
Oregon																		
Observed	1,268	277.4	8.08	380	76.9	4.04	123	28.9	2.71	120	31.5	2.97	109	21.2	2.08	75	14.2	1.67
Delay Adjusted	1,334	292.1	8.52	403	81.6	4.28	129	30.3	2.85	122	32.2	3.02	111	21.6	2.12	76	14.3	1.69
Pennsylvania																		
Observed	2,314	266.6	5.90	699	73.8	2.91	184	23.2	1.82	211	28.5	2.09	262	24.1	1.54	113	11.8	1.15
Delay Adjusted	2,398	276.3	6.11	714	75.4	2.97	188	23.7	1.86	214	28.8	2.11	283	26.1	1.66	114	11.9	1.16
Rhode Island																		
Observed	168	238.8	19.70	52	68.9	10.01	15	24.0	6.57	13	21.1	6.24	10	11.2	3.57	14	17.5	4.82
Delay Adjusted	174	247.1	20.39	53	69.9	10.16	15	24.2	6.65	13	21.4	6.33	10	11.3	3.61	14	18.0	4.96

1. Rates are per 100,000 population and are age-adjusted by five-year age groups to the 2000 U.S. standard population based on single years of age.
2. See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.
3. Abbreviations: NOS - Not Otherwise Specified; S.E. - Standard Error

^ There are insufficient historical registry data to calculate delay factors necessary for delay adjusted rates.

\- Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for All Sites combined.

Note: Delay adjusted counts are rounded to the nearest whole number. A small registry and/or rare cancer site may have the same delay adjusted count and observed count, but this is just due to rounding to whole counts. For more information about cancer incidence rates adjusted for reporting delay see https://surveillance.cancer.gov/delay/.

Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Delay Adjusted Counts and Rates[1] for the Five Most Commonly Diagnosed Cancers
Selected Areas in the NAACCR United States Combined[2], 2010-2014, Asian/Pacific Islander, Females

	All Sites			Prostate			Lung & Bronchus			Lung & Bronchus			Thyroid			Uterus Corpus & Uterus, NOS[3]		
	Count	*Rate*	*S.E.*[3]	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
South Carolina																		
Observed	459	263.3	13.24	145	76.3	6.66	39	24.8	4.30	45	30.9	4.95	32	15.3	2.72	25	12.7	2.61
Delay Adjusted	472	271.1	13.65	149	78.4	6.85	40	25.6	4.45	46	31.8	5.07	33	15.8	2.81	26	13.1	2.69
South Dakota																		
Observed	56	288.3	42.19	13	72.8	22.26	7	44.7	18.15	-	-	-	-	-	-	-	-	-
Delay Adjusted	58	300.6	43.97	13	74.2	22.66	7	46.4	18.76	-	-	-	-	-	-	-	-	-
Tennessee																		
Observed	592	248.1	11.26	182	70.2	5.62	59	26.4	3.79	64	33.4	4.57	43	15.2	2.50	29	11.2	2.20
Delay Adjusted	608	254.8	11.56	185	71.4	5.72	60	26.9	3.86	65	34.0	4.64	44	15.5	2.54	29	11.4	2.23
Texas																		
Observed	5,699	221.3	3.16	1,854	65.0	1.59	531	22.1	1.04	515	23.7	1.12	503	16.2	0.75	337	11.7	0.67
Delay Adjusted	6,024	233.9	3.34	1,923	67.4	1.65	550	22.9	1.07	526	24.2	1.15	528	16.9	0.79	346	12.0	0.69
Utah																		
Observed	590	321.1	14.06	175	91.2	7.28	59	36.0	4.93	45	31.0	4.83	45	19.6	3.03	72	34.3	4.19
Delay Adjusted	621	338.2	14.81	181	94.5	7.54	60	36.3	4.98	46	31.3	4.89	45	19.7	3.05	72	34.4	4.20
Vermont																		
Observed	42	246.9	42.73	14	89.6	26.76	-	-	-	8	59.4	22.56	-	-	-	-	-	-
Delay Adjusted	42	249.3	43.10	14	90.0	26.90	-	-	-	8	59.7	22.68	-	-	-	-	-	-
Virginia																		
Observed	3,118	250.6	4.75	1,057	79.0	2.53	260	22.5	1.48	272	25.9	1.67	368	25.2	1.35	170	12.3	0.98
Delay Adjusted	3,201	257.1	4.87	1,082	80.8	2.59	264	22.9	1.51	275	26.2	1.68	377	25.9	1.38	173	12.5	0.99
Washington																		
Observed	4,913	311.4	4.55	1,618	98.3	2.49	424	27.5	1.37	513	35.6	1.61	352	20.3	1.10	331	19.4	1.09
Delay Adjusted	5,035	319.1	4.66	1,658	100.7	2.55	431	27.9	1.39	519	36.0	1.63	357	20.7	1.12	335	19.7	1.10
Seattle																		
Observed	4,553	321.3	4.87	1,483	100.3	2.65	408	29.4	1.49	474	36.7	1.72	334	21.5	1.19	308	20.2	1.17
Delay Adjusted	4,654	328.4	4.97	1,515	102.4	2.71	413	29.8	1.51	478	37.0	1.74	337	21.7	1.20	310	20.3	1.18
West Virginia																		
Observed	87	221.9	25.03	33	77.8	14.01	-	-	-	6	18.6	8.13	9	21.8	7.29	7	17.6	7.12
Delay Adjusted	89	227.2	25.66	34	80.0	14.42	-	-	-	6	19.2	8.37	9	22.3	7.45	7	17.8	7.18
Wisconsin																		
Observed	703	299.6	12.14	184	74.2	5.74	62	31.5	4.23	62	32.8	4.36	66	20.7	2.73	60	22.4	3.04
Delay Adjusted	707	301.4	12.21	190	76.7	5.93	64	32.6	4.37	63	33.1	4.41	68	21.2	2.81	62	23.3	3.17

Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Delay Adjusted Counts and Rates[1] for the Five Most Commonly Diagnosed Cancers
Selected Areas in the NAACCR United States Combined[2], 2010-2014, Asian/Pacific Islander, Females

	All Sites			Prostate			Lung & Bronchus			Lung & Bronchus			Thyroid			Uterus Corpus & Uterus, NOS[3]		
	Count	*Rate*	*S.E.*[3]	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
Wyoming																		
Observed	26	196.2	39.91	8	58.4	21.46	-	-	-	-	-	-	-	-	-	-	-	-
Delay Adjusted	26	198.8	40.46	8	59.0	21.69	-	-	-	-	-	-	-	-	-	-	-	-

1. Rates are per 100,000 population and are age-adjusted by five-year age groups to the 2000 U.S. standard population based on single years of age.
2. See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.
3. Abbreviations: NOS - Not Otherwise Specified; S.E. - Standard Error

^ There are insufficient historical registry data to calculate delay factors necessary for delay adjusted rates.

\- Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for All Sites combined.

Note: Delay adjusted counts are rounded to the nearest whole number. A small registry and/or rare cancer site may have the same delay adjusted count and observed count, but this is just due to rounding to whole counts. For more information about cancer incidence rates adjusted for reporting delay see https://surveillance.cancer.gov/delay/.

NAACCR U.S. Combined Delay Adjusted Incidence 2010-2014, American Indian/Alaskan Native for CHSDA Combined

UNITED STATES

V-57 Combined Delay Adjusted Incidence – American Indian/Alaskan Native for CHSDA

Average Annual Age-adjusted Cancer Incidence Delay Adjusted Counts and Rates*, Selected Sites NAACCR United States Combined[1], 2010-2014, American Indian/Alaskan Native for CHSDA[2]

	MALES (ALL RACES)						FEMALES (ALL RACES)					
	Observed			Delay Adjusted			Observed			Delay Adjusted		
SITES	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
All Sites (invasive & in situ bladder)	12,594	417.8	4.13	13,562	450.0	4.45	14,068	382.5	3.38	15,149	411.9	3.64
Oral Cavity & Pharynx	554	16.3	0.76	591	17.3	0.81	228	6.1	0.43	244	6.6	0.46
Digestive												
Esophagus	236	7.8	0.56	245	8.1	0.58	80	2.3	0.27	83	2.4	0.28
Stomach	329	11.6	0.71	340	12.0	0.73	222	6.5	0.46	229	6.8	0.47
Colon & Rectum	1,513	50.1	1.43	1,622	53.7	1.53	1,463	41.3	1.13	1,570	44.4	1.22
Liver & IBD[3]	687	20.1	0.84	722	21.1	0.88	320	8.8	0.52	335	9.3	0.55
Pancreas	358	12.0	0.70	373	12.5	0.73	359	10.8	0.60	374	11.2	0.62
Respiratory												
Larynx	159	5.1	0.44	170	5.4	0.47	35	0.8	0.15	37	0.9	0.16
Lung & bronchus	1,911	71.9	1.80	2,006	75.5	1.89	1,875	55.9	1.35	1,967	58.6	1.42
Skin												
Melanomas of the skin	251	8.6	0.61	288	9.9	0.70	210	5.6	0.40	239	6.3	0.46
Breast (female cases only)	-	-	-	-	-	-	3,839	100.7	1.69	4,173	109.4	1.84
Female Genital												
Cervix uteri	-	-	-	-	-	-	374	9.1	0.49	404	9.9	0.52
Corpus & uterus, NOS[3]	-	-	-	-	-	-	896	22.4	0.77	941	23.5	0.81
Ovary	-	-	-	-	-	-	399	10.6	0.56	434	11.6	0.61
Male Genital												
Prostate	2,311	78.3	1.78	2,572	87.1	1.98	-	-	-	-	-	-
Testis	243	4.9	0.32	261	5.3	0.35	-	-	-	-	-	-
Urinary												
Bladder (incl. *in situ*)	489	19.4	0.98	537	21.4	1.07	194	5.9	0.45	212	6.5	0.49
Kidney & renal pelvis	994	30.0	1.03	1,072	32.3	1.12	670	17.4	0.70	722	18.7	0.76
Brain & Other Nervous System	210	5.4	0.42	228	5.9	0.46	154	3.6	0.31	168	3.9	0.33
Endocrine												
Thyroid gland	164	4.4	0.37	181	4.8	0.41	618	14.5	0.60	684	16.1	0.67
Lymphomas												
Hodgkin Lymphoma	63	1.5	0.20	66	1.5	0.21	43	1.0	0.17	45	1.1	0.18
Non-Hodgkin Lymphoma	500	16.7	0.84	544	18.2	0.91	472	13.5	0.65	515	14.7	0.71
Multiple Myeloma	214	7.8	0.59	244	8.8	0.67	173	5.0	0.40	195	5.7	0.45
Leukemias	408	12.4	0.70	456	13.9	0.78	345	8.8	0.50	382	9.7	0.56

* Rates are per 100,000 population and are age-adjusted by five-year age groups to the 2000 U.S. standard population based on single years of age and to standard.

\- Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for All Sites combined.

[1] NAACCR United States Combined includes registries for which a U.S. average delay adjustment factor was calculated. See technical notes.
[2] CHSDA - Indian Health Services Contract Health Service Delivery Areas (See Appendix G)
[3] Abbreviations: IBD – Intrahepatic Bile Duct; NOS - Not Otherwise Specified

Note: Delay adjusted counts are rounded to the nearest whole number. A small registry and/or rare cancer site may have the same delay adjusted count and observed count, but this is just due to rounding to whole counts. For more information about cancer incidence rates adjusted for reporting delay see https://surveillance.cancer.gov/delay/.

Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Delay Adjusted Counts and Rates[1] for the Five Most Commonly Diagnosed Cancers Selected Areas in the NAACCR United States Combined[2], 2010-2014, American Indian/Alaskan Native for CHSDA[3] by Region, Males

	All Sites			Prostate			Lung & Bronchus			Colon & Rectum		
	Count	*Rate*	*S.E.[4]*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
NAACCR U.S. Combined												
Observed	12,594	417.8	4.13	2,311	78.3	1.78	1,911	71.9	1.80	1,513	50.1	1.43
Delay Adjusted	13,562	450.0	4.45	2,572	87.1	1.98	2,006	75.5	1.89	1,622	53.7	1.53
Alaska												
Observed	967	499.9	17.97	99	54.1	6.03	178	108.8	8.95	178	91.7	7.60
Delay Adjusted	988	510.8	18.36	102	56.0	6.26	181	110.4	9.08	181	93.4	7.75
East												
Observed	816	279.0	10.86	183	62.2	5.04	123	45.0	4.49	96	35.4	4.06
Delay Adjusted	890	304.1	11.84	207	70.5	5.72	134	48.8	4.86	103	37.9	4.36
Northern Plains												
Observed	2,254	545.6	13.09	438	99.1	5.23	402	117.0	6.52	284	66.8	4.50
Delay Adjusted	2,363	572.1	13.74	478	108.2	5.70	418	121.7	6.80	294	69.3	4.67
Southern Plains												
Observed	3,931	584.3	10.17	737	110.7	4.43	692	110.6	4.54	459	68.7	3.50
Delay Adjusted	4,215	626.5	10.90	814	122.0	4.88	719	114.9	4.73	490	73.2	3.73
Pacific Coast												
Observed	2,828	325.6	6.88	518	60.1	2.90	400	53.5	2.99	292	34.7	2.28
Delay Adjusted	3,136	361.1	7.64	589	68.3	3.30	426	57.0	3.18	321	38.2	2.51
Southwest												
Observed	1,798	309.6	7.99	336	66.6	3.88	116	23.4	2.33	204	33.0	2.50
Delay Adjusted	1,970	339.2	8.76	381	75.4	4.40	129	26.0	2.59	233	37.6	2.86

1. Rates are per 100,000 population and are age-adjusted by five-year age groups to the 2000 U.S. standard population based on single years of age.
2. See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.
3. CHSDA – Indian Health Services Contract Service Delivery Areas (See Appendix G).
4. Abbreviations: S.E. - Standard Error

^ There are insufficient historical registry data to calculate delay factors necessary for delay adjusted rates.

\- Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for All Sites combined.

Note: Delay adjusted counts are rounded to the nearest whole number. A small registry and/or rare cancer site may have the same delay adjusted count and observed count, but this is just due to rounding to whole counts. For more information about cancer incidence rates adjusted for reporting delay see https://surveillance.cancer.gov/delay/.

Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Delay Adjusted Counts and Rates[1] for the Five Most Commonly Diagnosed Cancers Selected Areas in the NAACCR United States Combined[2], 2010-2014, American Indian/Alaskan Native for CHSDA[3] by Region, Females

	All Sites			Breast			Lung & Bronchus			Colon & Rectum		
	Count	*Rate*	*S.E.[4]*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
NAACCR U.S. Combined												
Observed	14,068	382.5	3.38	3,839	100.7	1.69	1,875	55.9	1.35	1,463	41.3	1.13
Delay Adjusted	15,149	411.9	3.64	4,173	109.4	1.84	1,967	58.6	1.42	1,570	44.4	1.22
Alaska												
Observed	1,160	509.8	15.70	343	144.8	8.14	144	68.8	5.99	184	86.4	6.66
Delay Adjusted	1,185	520.9	16.05	349	147.4	8.29	146	69.7	6.08	188	88.2	6.79
East												
Observed	873	261.2	9.23	244	70.7	4.71	123	38.6	3.63	83	25.0	2.86
Delay Adjusted	954	285.5	10.10	268	77.6	5.17	135	42.4	3.99	89	26.9	3.08
Northern Plains												
Observed	2,495	485.4	10.33	691	125.5	5.00	435	96.6	4.91	268	52.2	3.38
Delay Adjusted	2,615	508.9	10.84	729	132.5	5.28	452	100.4	5.11	278	54.1	3.51
Southern Plains												
Observed	4,258	519.8	8.23	1,194	143.1	4.26	661	84.5	3.40	437	54.9	2.72
Delay Adjusted	4,569	557.8	8.84	1,297	155.5	4.63	686	87.7	3.53	466	58.6	2.90
Pacific Coast												
Observed	3,176	310.3	5.87	908	85.8	3.01	388	42.9	2.32	307	32.1	1.95
Delay Adjusted	3,519	343.6	6.50	1,026	97.0	3.40	410	45.3	2.45	340	35.5	2.16
Southwest												
Observed	2,106	277.4	6.28	459	58.3	2.81	124	18.1	1.68	184	25.2	1.93
Delay Adjusted	2,307	304.0	6.89	503	64.0	3.08	138	20.1	1.87	209	28.7	2.21

1. Rates are per 100,000 population and are age-adjusted by five-year age groups to the 2000 U.S. standard population based on single years of age.
2. See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.
3. CHSDA – Indian Health Services Contract Service Delivery Areas (See Appendix G).
4. Abbreviations: S.E. - Standard Error

^ There are insufficient historical registry data to calculate delay factors necessary for delay adjusted rates.

\- Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for All Sites combined.

Note: Delay adjusted counts are rounded to the nearest whole number. A small registry and/or rare cancer site may have the same delay adjusted count and observed count, but this is just due to rounding to whole counts. For more information about cancer incidence rates adjusted for reporting delay see https://surveillance.cancer.gov/delay/.

NAACCR U.S. Combined Delay Adjusted Incidence 2010-2014, Hispanic/Latino, All Races Combined

UNITED STATES

V-61 Combined Delay Adjusted Incidence – Hispanic/Latino, All Races

Average Annual Age-adjusted Cancer Incidence Delay Adjusted Counts and Rates*, Selected Sites
NAACCR United States Combined[1], 2010-2014, Hispanic/Latino, All Races

	MALES (ALL RACES)						FEMALES (ALL RACES)					
	Observed			Delay Adjusted			Observed			Delay Adjusted		
SITES	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
All Sites (invasive & in situ bladder)	262,878	378.8	0.81	271,114	390.8	0.83	288,386	324.0	0.63	297,556	334.3	0.65
Oral Cavity & Pharynx	7,836	10.5	0.13	8,095	10.9	0.13	3,649	4.2	0.07	3,766	4.3	0.07
Digestive												
Esophagus	3,152	4.8	0.09	3,205	4.9	0.09	842	1.1	0.04	856	1.1	0.04
Stomach	8,479	12.9	0.15	8,618	13.1	0.15	6,462	7.7	0.10	6,568	7.9	0.10
Colon & Rectum	28,612	41.9	0.27	29,222	42.8	0.28	24,205	29.3	0.19	24,726	29.9	0.20
Liver & IBD[2]	14,558	19.7	0.18	14,885	20.2	0.18	6,069	7.6	0.10	6,202	7.8	0.10
Pancreas	7,607	12.0	0.15	7,743	12.2	0.15	7,924	10.3	0.12	8,064	10.5	0.12
Respiratory												
Larynx	3,211	4.8	0.09	3,272	4.9	0.09	543	0.6	0.03	552	0.7	0.03
Lung & bronchus	23,420	40.6	0.28	23,781	41.2	0.29	19,371	25.2	0.19	19,681	25.6	0.19
Skin												
Melanomas of the skin	3,394	4.9	0.09	3,544	5.1	0.10	3,985	4.2	0.07	4,141	4.4	0.07
Breast (female cases only)	-	-	-	-	-	-	83,873	91.9	0.33	86,639	94.9	0.34
Female Genital												
Cervix uteri	-	-	-	-	-	-	10,009	9.7	0.10	10,197	9.9	0.10
Corpus & uterus, NOS[2]	-	-	-	-	-	-	20,579	22.2	0.16	20,924	22.6	0.16
Ovary	-	-	-	-	-	-	9,452	10.3	0.11	9,779	10.6	0.11
Male Genital												
Prostate	63,511	97.1	0.41	66,185	101.2	0.43	-	-	-	-	-	-
Testis	6,559	4.5	0.06	6,667	4.6	0.06	-	-	-	-	-	-
Urinary												
Bladder (incl. *in situ*)	10,984	19.5	0.20	11,339	20.1	0.20	3,791	5.0	0.08	3,911	5.1	0.09
Kidney & renal pelvis	15,570	20.8	0.18	16,035	21.4	0.19	10,632	12.0	0.12	10,953	12.4	0.12
Brain & Other Nervous System	5,555	5.9	0.09	5,738	6.1	0.09	4,659	4.5	0.07	4,809	4.6	0.07
Endocrine												
Thyroid gland	4,987	5.4	0.08	5,086	5.5	0.08	21,544	20.2	0.14	21,956	20.5	0.14
Lymphomas												
Hodgkin Lymphoma	2,981	2.8	0.06	3,051	2.9	0.06	2,343	2.1	0.05	2,399	2.1	0.05
Non-Hodgkin Lymphoma	14,532	19.8	0.18	15,038	20.5	0.19	12,764	15.1	0.14	13,215	15.6	0.14
Multiple Myeloma	5,092	7.9	0.12	5,468	8.4	0.13	4,262	5.3	0.08	4,585	5.7	0.09
Leukemias	11,225	13.0	0.14	12,025	14.0	0.15	8,904	8.9	0.10	9,551	9.6	0.11

* Rates are per 100,000 population and are age-adjusted by five-year age groups to the 2000 U.S. standard population based on single years of age and to standard.

\- Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for All Sites combined.

[1] NAACCR United States Combined includes registries for which a U.S. average delay adjustment factor was calculated. See technical notes.

[2] Abbreviations: IBD – Intrahepatic Bile Duct; NOS - Not Otherwise Specified

Note: Delay adjusted counts are rounded to the nearest whole number. A small registry and/or rare cancer site may have the same delay adjusted count and observed count, but this is just due to rounding to whole counts. For more information about cancer incidence rates adjusted for reporting delay see https://surveillance.cancer.gov/delay/.

Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Delay Adjusted Counts and Rates[1] for the Five Most Commonly Diagnosed Cancers
Selected Areas in the NAACCR United States Combined[2], 2010-2014, Hispanic/Latino, All Races, Males

	All Sites			Prostate			Colon & Rectum			Lung & Bronchus			Kidney & Renal Pelvis			Non-Hodgkin Lymphoma		
	Count	*Rate*	*S.E.*[3]	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
NAACCR U.S. Combined																		
Observed	262,878	378.8	0.81	63,511	97.1	0.41	28,612	41.9	0.27	23,420	40.6	0.28	15,570	20.8	0.18	14,532	19.8	0.18
Delay Adjusted	271,114	390.8	0.83	66,185	101.2	0.43	29,222	42.8	0.28	23,781	41.2	0.29	16,035	21.4	0.19	15,038	20.5	0.19
Alabama																		
Observed	478	218.1	12.20	101	55.8	6.18	45	22.1	3.93	38	24.2	4.45	31	13.1	2.85	25	11.6	2.92
Delay Adjusted	484	220.5	12.33	103	56.8	6.28	45	22.3	3.97	38	24.3	4.47	31	13.3	2.87	25	11.7	2.95
Alaska																		
Observed	147	295.8	30.64	25	57.4	14.56	19	27.2	6.67	18	50.4	14.35	-	-	-	15	27.4	8.32
Delay Adjusted	159	319.1	33.06	26	60.1	15.26	19	27.7	6.80	18	51.2	14.54	-	-	-	16	28.8	8.71
Arizona																		
Observed	8,739	336.2	3.96	1,654	66.2	1.75	1,086	43.5	1.44	779	38.1	1.45	648	23.5	1.00	420	15.7	0.85
Delay Adjusted	9,357	360.0	4.25	1,801	72.1	1.90	1,143	45.8	1.52	797	39.0	1.49	706	25.6	1.09	447	16.6	0.90
Arkansas																		
Observed	1,218	722.4	25.27	192	122.6	10.11	133	78.1	8.31	171	115.9	10.41	77	45.3	6.16	58	35.0	5.93
Delay Adjusted	1,249	740.8	25.91	196	125.1	10.31	134	78.9	8.39	173	117.1	10.51	79	46.4	6.32	59	35.9	6.09
California																		
Observed	71,237	362.5	1.49	17,520	95.9	0.77	7,706	39.2	0.49	4,988	31.8	0.48	4,358	20.7	0.34	4,213	20.6	0.35
Delay Adjusted	72,535	369.2	1.52	17,978	98.4	0.79	7,795	39.7	0.49	5,063	32.3	0.48	4,439	21.1	0.35	4,294	21.0	0.36
Greater California																		
Observed	37,357	364.3	2.07	8,802	93.4	1.06	4,102	39.5	0.67	2,751	33.7	0.68	2,452	22.4	0.49	2,181	20.6	0.49
Delay Adjusted	38,020	370.9	2.11	9,053	96.1	1.09	4,148	39.9	0.68	2,788	34.2	0.69	2,497	22.8	0.50	2,227	21.0	0.50
Greater Bay																		
Observed	9,742	398.2	4.43	2,644	115.3	2.39	932	38.9	1.39	678	35.5	1.44	536	19.3	0.91	592	23.3	1.06
Delay Adjusted	9,925	405.7	4.51	2,668	116.4	2.42	943	39.4	1.41	691	36.2	1.47	546	19.7	0.93	602	23.7	1.08
Los Angeles																		
Observed	24,119	346.8	2.43	6,066	92.5	1.26	2,671	39.0	0.82	1,558	27.6	0.74	1,370	18.5	0.54	1,440	19.7	0.57
Delay Adjusted	24,571	353.4	2.48	6,248	95.3	1.30	2,703	39.4	0.83	1,583	28.0	0.75	1,397	18.9	0.56	1,465	20.0	0.58
Colorado																		
Observed	5,509	377.2	5.61	1,206	86.5	2.67	645	43.5	1.88	486	41.2	2.00	385	24.9	1.41	253	17.1	1.20
Delay Adjusted	5,623	385.1	5.73	1,272	91.2	2.82	649	43.8	1.90	489	41.5	2.02	392	25.3	1.43	258	17.4	1.22
Connecticut																		
Observed	3,172	507.5	10.33	722	120.5	4.95	317	51.6	3.35	329	62.7	3.82	135	19.5	1.94	180	26.1	2.32
Delay Adjusted	3,362	538.4	10.97	783	131.1	5.40	331	53.9	3.50	335	63.9	3.89	147	21.2	2.10	187	27.2	2.41

Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Delay Adjusted Counts and Rates[1] for the Five Most Commonly Diagnosed Cancers
Selected Areas in the NAACCR United States Combined[2], 2010-2014, Hispanic/Latino, All Races, Males

	All Sites			Prostate			Colon & Rectum			Lung & Bronchus			Kidney & Renal Pelvis			Non-Hodgkin Lymphoma		
	Count	*Rate*	*S.E.*[3]	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
Delaware																		
Observed	352	421.7	25.75	103	139.1	14.98	28	31.1	6.60	28	37.6	7.85	12	12.4	4.15	13	13.4	4.09
Delay Adjusted	376	450.8	27.53	108	145.4	15.66	29	31.8	6.75	29	38.5	8.05	12	12.9	4.30	14	14.1	4.32
District of Columbia																		
Observed	267	316.7	23.08	58	78.4	11.59	32	31.7	6.81	18	32.1	8.33	12	16.7	5.87	15	17.8	5.56
Delay Adjusted	277	327.3	23.83	59	80.6	12.01	33	32.6	7.00	18	32.8	8.53	12	17.1	6.01	16	18.6	5.81
Florida																		
Observed	37,432	416.0	2.19	9,859	110.0	1.12	4,052	45.7	0.73	4,584	53.8	0.81	1,656	17.8	0.45	1,903	20.4	0.48
Delay Adjusted	39,407	437.7	2.31	10,522	117.4	1.20	4,205	47.4	0.76	4,619	54.2	0.81	1,734	18.7	0.47	2,034	21.8	0.51
Georgia																		
Observed	2,616	328.4	8.30	578	89.8	4.33	233	30.8	2.60	225	39.3	3.15	124	14.8	1.65	195	18.9	1.87
Delay Adjusted	2,726	342.5	8.66	619	96.2	4.64	236	31.3	2.63	227	39.7	3.19	127	15.1	1.68	200	19.4	1.93
Hawaii																		
Observed	640	438.0	19.46	108	75.1	7.91	67	44.5	6.10	93	79.8	8.98	35	20.0	3.73	33	23.8	4.69
Delay Adjusted	749	512.1	22.74	140	97.4	10.23	67	44.7	6.13	102	87.6	9.87	35	20.3	3.78	33	24.1	4.75
Idaho																		
Observed	676	352.9	15.99	143	81.7	7.65	69	37.7	5.11	65	43.8	6.13	47	22.3	3.81	37	16.4	3.24
Delay Adjusted	694	362.4	16.43	150	85.5	8.01	70	38.2	5.18	66	44.7	6.25	49	23.1	3.94	38	16.8	3.32
Illinois																		
Observed	9,462	369.0	4.38	2,403	99.8	2.25	944	37.7	1.41	713	37.2	1.53	611	21.5	1.00	503	18.5	0.97
Delay Adjusted	9,649	376.4	4.47	2,480	103.1	2.33	961	38.4	1.43	723	37.7	1.55	619	21.8	1.02	513	18.8	0.99
Indiana																		
Observed	1,406	316.8	9.85	291	75.3	4.86	157	36.6	3.35	137	38.0	3.58	96	19.4	2.32	71	15.7	2.20
Delay Adjusted	1,453	327.5	10.18	303	78.4	5.07	161	37.5	3.44	140	38.7	3.65	98	19.9	2.38	74	16.4	2.29
Iowa																		
Observed	452	284.3	16.38	81	62.1	7.86	46	29.6	5.12	34	33.8	6.37	21	17.0	4.28	27	15.1	3.67
Delay Adjusted	459	288.9	16.65	82	63.3	8.03	46	29.9	5.17	34	34.3	6.46	21	17.4	4.37	28	15.3	3.74
Kansas																		
Observed	1,247	384.2	12.55	290	100.4	6.41	136	45.9	4.42	100	40.3	4.41	91	27.3	3.31	63	17.2	2.56
Delay Adjusted	^	^	^	^	^	^	^	^	^	^	^	^	^	^	^	^	^	^

1. Rates are per 100,000 population and are age-adjusted by five-year age groups to the 2000 U.S. standard population based on single years of age.
2. See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.
3. Abbreviations: S.E. - Standard Error

^ There are insufficient historical registry data to calculate delay factors necessary for delay adjusted rates.

\- Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for All Sites combined.

Note: Delay adjusted counts are rounded to the nearest whole number. A small registry and/or rare cancer site may have the same delay adjusted count and observed count, but this is just due to rounding to whole counts. For more information about cancer incidence rates adjusted for reporting delay see https://surveillance.cancer.gov/delay/.

Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Delay Adjusted Counts and Rates[1] for the Five Most Commonly Diagnosed Cancers
Selected Areas in the NAACCR United States Combined[2], 2010-2014, Hispanic/Latino, All Races, Males

	All Sites			Prostate			Colon & Rectum			Lung & Bronchus			Kidney & Renal Pelvis			Non-Hodgkin Lymphoma		
	Count	*Rate*	*S.E.*[3]	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
Kentucky																		
Observed	376	241.2	14.82	76	55.8	7.08	42	26.9	4.86	54	42.0	6.46	25	13.3	3.17	26	13.7	3.59
Delay Adjusted	382	245.4	15.08	78	57.4	7.27	43	27.4	4.95	54	42.3	6.51	26	13.7	3.26	27	14.0	3.67
Louisiana																		
Observed	1,143	361.0	11.58	309	107.5	6.46	102	31.5	3.37	125	44.1	4.20	66	21.3	2.83	64	17.1	2.37
Delay Adjusted	1,210	382.3	12.27	332	115.4	6.93	103	31.7	3.39	131	46.3	4.41	67	21.5	2.86	65	17.3	2.39
Maine																		
Observed	77	289.2	35.62	16	63.9	16.97	-	-	-	7	34.1	13.38	8	32.4	12.01	-	-	-
Delay Adjusted	78	292.8	36.05	16	65.0	17.24	-	-	-	7	34.3	13.46	8	33.1	12.26	-	-	-
Maryland																		
Observed	1,645	304.6	9.17	460	96.5	5.11	160	29.2	2.83	100	26.7	2.99	78	16.2	2.22	96	15.3	1.96
Delay Adjusted	1,743	322.7	9.72	512	107.4	5.69	163	29.7	2.88	102	27.3	3.06	81	16.7	2.29	99	15.7	2.01
Massachusetts																		
Observed	3,183	383.1	7.80	953	120.0	4.29	262	34.9	2.51	263	38.6	2.65	166	17.4	1.55	175	19.1	1.69
Delay Adjusted	3,197	384.7	7.83	985	124.0	4.43	268	35.6	2.57	267	39.2	2.70	170	17.8	1.59	177	19.4	1.71
Michigan																		
Observed	2,061	373.2	9.13	481	89.8	4.42	223	42.5	3.13	207	45.3	3.40	136	23.8	2.26	121	22.1	2.25
Delay Adjusted	2,169	393.0	9.62	496	92.4	4.55	233	44.4	3.27	211	46.2	3.47	139	24.4	2.31	124	22.7	2.32
Detroit																		
Observed	1,022	506.4	17.25	250	127.8	8.61	114	57.6	5.86	109	62.3	6.37	65	32.5	4.36	59	29.0	4.12
Delay Adjusted	1,115	552.7	18.83	252	128.8	8.67	124	62.7	6.37	110	62.6	6.40	66	32.9	4.42	60	29.3	4.18
Mississippi																		
Observed	215	211.1	17.05	55	64.5	9.77	25	24.9	5.77	22	27.8	6.50	12	11.0	3.85	11	9.6	3.41
Delay Adjusted	223	219.5	17.74	56	66.5	10.09	25	25.2	5.84	22	28.2	6.59	12	11.4	4.02	11	10.0	3.55
Missouri																		
Observed	745	277.9	11.43	140	59.6	5.39	78	29.9	3.77	87	41.8	4.79	47	14.7	2.35	51	18.5	3.01
Delay Adjusted	803	299.4	12.32	145	61.9	5.59	81	30.9	3.88	88	42.4	4.86	48	15.0	2.40	53	19.1	3.11
Montana																		
Observed	181	456.6	37.61	41	96.4	16.11	23	53.6	12.42	21	73.8	17.10	10	29.0	9.79	-	-	-
Delay Adjusted	188	474.9	39.12	43	101.3	16.96	24	55.5	12.83	21	75.1	17.39	10	30.4	10.29	-	-	-
Nebraska																		
Observed	634	351.8	17.11	138	96.3	9.23	55	30.3	4.99	49	35.8	5.72	52	27.4	4.58	41	25.0	4.79
Delay Adjusted	642	356.7	17.37	145	101.4	9.71	56	31.0	5.10	50	36.6	5.88	55	28.8	4.83	43	26.0	4.99

Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Delay Adjusted Counts and Rates[1] for the Five Most Commonly Diagnosed Cancers
Selected Areas in the NAACCR United States Combined[2], 2010-2014, Hispanic/Latino, All Races, Males

	All Sites			Prostate			Colon & Rectum			Lung & Bronchus			Kidney & Renal Pelvis			Non-Hodgkin Lymphoma		
	Count	*Rate*	*S.E.[3]*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
New Hampshire																		
Observed	174	364.9	31.58	50	109.3	16.88	14	27.9	8.80	16	48.5	12.96	7	17.8	7.45	10	16.4	6.24
Delay Adjusted	178	373.1	32.28	51	112.1	17.30	14	28.4	8.95	16	49.4	13.19	7	18.4	7.71	10	17.0	6.48
New Jersey																		
Observed	10,484	435.6	4.69	2,989	129.4	2.55	1,056	44.3	1.50	931	45.0	1.59	501	17.8	0.87	631	24.5	1.09
Delay Adjusted	10,929	454.1	4.89	3,136	135.9	2.68	1,101	46.2	1.57	955	46.2	1.63	526	18.6	0.91	649	25.2	1.12
New York																		
Observed	25,859	459.9	3.08	7,906	141.3	1.68	2,496	46.0	0.99	2,319	46.8	1.03	1,073	17.3	0.56	1,414	23.4	0.68
Delay Adjusted	26,061	463.6	3.10	8,040	143.9	1.71	2,510	46.3	1.00	2,339	47.2	1.04	1,092	17.6	0.57	1,442	23.9	0.70
North Carolina																		
Observed	2,205	295.9	8.20	486	81.2	4.27	186	25.8	2.48	168	31.2	2.90	131	15.7	1.82	143	15.6	1.80
Delay Adjusted	2,275	305.5	8.47	513	85.7	4.51	190	26.4	2.54	170	31.6	2.94	134	16.1	1.87	148	16.2	1.87
North Dakota																		
Observed	38	279.9	55.28	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Delay Adjusted	39	287.9	56.81	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Ohio																		
Observed	1,317	292.4	8.98	322	76.0	4.57	143	30.3	2.82	133	37.1	3.45	66	12.0	1.65	66	14.6	2.02
Delay Adjusted	1,335	296.6	9.11	325	76.6	4.61	145	30.8	2.86	136	38.0	3.54	68	12.4	1.70	69	15.2	2.10
Oklahoma																		
Observed	1,135	316.6	11.41	202	70.5	5.61	129	34.3	3.59	112	43.7	4.63	81	19.9	2.71	59	15.5	2.49
Delay Adjusted	1,146	319.8	11.53	206	72.0	5.73	130	34.6	3.61	114	44.3	4.70	83	20.4	2.78	60	15.9	2.56
Oregon																		
Observed	1,668	362.5	10.83	326	80.1	5.04	175	40.5	3.72	129	36.7	3.74	113	23.9	2.77	107	20.1	2.47
Delay Adjusted	1,696	368.9	11.03	332	81.9	5.17	177	41.1	3.76	132	37.6	3.83	115	24.2	2.80	110	20.6	2.54
Pennsylvania																		
Observed	3,834	414.5	7.69	1,006	116.3	4.05	388	43.3	2.52	376	49.5	2.84	170	16.5	1.45	186	18.1	1.55
Delay Adjusted	4,126	446.1	8.28	1,076	124.4	4.33	411	45.9	2.66	397	52.2	3.00	182	17.7	1.56	202	19.6	1.68
Rhode Island																		
Observed	564	365.6	18.08	150	103.1	9.50	62	39.5	6.02	56	43.5	6.51	22	10.9	2.50	28	18.2	4.05
Delay Adjusted	585	379.2	18.77	156	107.4	9.91	63	40.0	6.08	57	44.2	6.62	22	11.0	2.54	29	18.6	4.16

1. Rates are per 100,000 population and are age-adjusted by five-year age groups to the 2000 U.S. standard population based on single years of age.
2. See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.
3. Abbreviations: S.E. - Standard Error

^ There are insufficient historical registry data to calculate delay factors necessary for delay adjusted rates.

\- Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for All Sites combined.

Note: Delay adjusted counts are rounded to the nearest whole number. A small registry and/or rare cancer site may have the same delay adjusted count and observed count, but this is just due to rounding to whole counts. For more information about cancer incidence rates adjusted for reporting delay see https://surveillance.cancer.gov/delay/.

Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Delay Adjusted Counts and Rates[1] for the Five Most Commonly Diagnosed Cancers
Selected Areas in the NAACCR United States Combined[2], 2010-2014, Hispanic/Latino, All Races, Males

	All Sites			Prostate			Lung & Bronchus			Lung & Bronchus			Kidney & Renal Pelvis			Non-Hodgkin Lymphoma		
	Count	*Rate*	*S.E.*[3]	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
South Carolina																		
Observed	727	289.3	13.32	147	67.6	6.22	72	25.6	3.79	71	34.3	4.80	39	18.2	3.45	37	13.8	2.95
Delay Adjusted	751	299.0	13.78	162	74.6	6.89	74	26.3	3.89	72	34.9	4.89	41	19.0	3.62	39	14.4	3.06
South Dakota																		
Observed	74	255.3	33.44	13	53.6	15.61	8	25.5	9.60	6	33.9	13.96	-	-	-	11	36.9	13.37
Delay Adjusted	78	267.6	35.06	14	57.1	16.66	8	26.5	9.96	6	35.0	14.45	-	-	-	11	38.5	14.08
Tennessee																		
Observed	729	235.0	10.93	142	59.8	5.68	57	19.6	3.16	73	30.0	4.06	37	12.1	2.46	39	10.6	2.28
Delay Adjusted	733	236.2	10.99	145	60.9	5.78	57	19.7	3.19	74	30.3	4.09	38	12.3	2.50	40	10.9	2.34
Texas																		
Observed	50,914	368.9	1.79	10,000	78.2	0.84	6,400	46.5	0.63	4,662	41.0	0.64	3,914	26.1	0.45	2,677	19.0	0.41
Delay Adjusted	52,529	380.6	1.85	10,423	81.5	0.87	6,516	47.3	0.64	4,736	41.6	0.65	4,023	26.8	0.47	2,795	19.8	0.42
Utah																		
Observed	1,432	385.8	12.13	354	108.0	6.41	141	36.4	3.59	92	33.6	3.92	91	25.0	3.05	89	24.0	3.09
Delay Adjusted	1,471	396.8	12.48	372	113.5	6.72	142	36.7	3.62	94	34.4	4.02	94	25.8	3.16	90	24.3	3.12
Vermont																		
Observed	31	192.9	37.12	6	44.9	19.22	-	-	-	-	-	-	-	-	-	-	-	-
Delay Adjusted	31	195.1	37.54	6	45.8	19.59	-	-	-	-	-	-	-	-	-	-	-	-
Virginia																		
Observed	1,939	282.1	8.08	452	73.6	4.06	182	29.1	2.70	168	33.5	3.04	107	14.1	1.71	128	16.4	1.88
Delay Adjusted	2,001	290.7	8.32	471	76.7	4.22	189	30.1	2.80	170	33.9	3.08	110	14.5	1.76	133	17.0	1.94
Washington																		
Observed	2,816	360.9	8.32	594	92.7	4.33	253	31.0	2.29	227	40.8	3.07	156	18.8	1.82	182	22.2	2.06
Delay Adjusted	2,983	381.9	8.80	645	100.4	4.69	262	32.1	2.38	236	42.4	3.20	159	19.2	1.86	191	23.3	2.16
Seattle																		
Observed	1,706	421.4	12.79	358	105.6	6.44	147	36.4	3.67	127	44.3	4.62	102	21.3	2.64	96	21.9	2.85
Delay Adjusted	1,845	455.5	13.82	404	119.0	7.25	153	37.9	3.82	133	46.4	4.85	103	21.5	2.66	102	23.2	3.02
West Virginia																		
Observed	81	195.5	22.95	18	49.4	12.04	11	25.5	8.19	9	24.3	8.46	-	-	-	-	-	-
Delay Adjusted	84	201.6	23.66	18	50.6	12.38	11	26.1	8.36	9	25.0	8.73	-	-	-	-	-	-
Wisconsin																		
Observed	1,317	390.6	12.83	292	92.2	6.07	110	30.8	3.47	104	39.8	4.40	85	22.7	2.89	87	24.7	3.28
Delay Adjusted	1,372	409.1	13.49	307	96.8	6.37	115	32.0	3.59	108	41.1	4.54	90	24.0	3.06	93	26.4	3.50

Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Delay Adjusted Counts and Rates[1] for the Five Most Commonly Diagnosed Cancers
Selected Areas in the NAACCR United States Combined[2], 2010-2014, Hispanic/Latino, All Races, Males

	All Sites			Prostate			Lung & Bronchus			Lung & Bronchus			Kidney & Renal Pelvis			Non-Hodgkin Lymphoma		
	Count	*Rate*	*S.E.*[3]	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
Wyoming																		
Observed	230	267.9	18.83	48	57.3	8.72	33	38.3	7.07	22	32.9	7.36	18	18.9	4.72	9	11.0	3.88
Delay Adjusted	232	270.6	19.02	48	57.8	8.80	33	38.4	7.08	22	33.0	7.38	18	19.2	4.79	9	11.1	3.93

1. Rates are per 100,000 population and are age-adjusted by five-year age groups to the 2000 U.S. standard population based on single years of age.
2. See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.
3. Abbreviations: S.E. - Standard Error

^ There are insufficient historical registry data to calculate delay factors necessary for delay adjusted rates.

\- Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for All Sites combined.

Note: Delay adjusted counts are rounded to the nearest whole number. A small registry and/or rare cancer site may have the same delay adjusted count and observed count, but this is just due to rounding to whole counts. For more information about cancer incidence rates adjusted for reporting delay see https://surveillance.cancer.gov/delay/.

Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Delay Adjusted Counts and Rates[1] for the Five Most Commonly Diagnosed Cancers
Selected Areas in the NAACCR United States Combined[2], 2010-2014, Hispanic/Latina, All Races, Females

	All Sites			Breast			Colon & Rectum			Thyroid			Uterus Corpus & Uterus, NOS[3]			Lung & Bronchus		
	Count	*Rate*	*S.E.*[3]	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
NAACCR U.S. Combined																		
Observed	288,386	324.0	0.63	83,873	91.9	0.33	24,205	29.3	0.19	21,544	20.2	0.14	20,579	22.2	0.16	19,371	25.2	0.19
Delay Adjusted	297,556	334.3	0.65	86,639	94.9	0.34	24,726	29.9	0.20	21,956	20.5	0.14	20,924	22.6	0.16	19,681	25.6	0.19
Alabama																		
Observed	449	203.1	10.72	143	64.3	5.83	44	25.1	4.08	38	12.1	2.25	23	10.8	2.49	33	20.8	3.79
Delay Adjusted	455	205.6	10.84	145	65.2	5.91	44	25.4	4.11	38	12.2	2.27	23	10.8	2.50	33	21.0	3.81
Alaska																		
Observed	166	259.2	22.84	54	80.4	12.10	22	32.3	7.21	13	13.6	3.88	8	9.8	3.51	9	15.4	5.58
Delay Adjusted	179	280.0	24.67	55	81.7	12.30	22	32.9	7.35	13	14.0	3.99	8	10.2	3.63	9	15.9	5.74
Arizona																		
Observed	10,457	319.9	3.29	2,976	88.9	1.69	847	28.2	1.01	901	22.3	0.77	712	20.3	0.79	734	26.8	1.02
Delay Adjusted	11,198	342.6	3.52	3,256	97.3	1.85	891	29.7	1.06	950	23.5	0.81	747	21.3	0.83	752	27.5	1.05
Arkansas																		
Observed	1,310	681.7	22.23	297	148.1	10.10	116	70.7	7.64	82	24.4	3.13	80	39.7	5.10	130	90.6	8.76
Delay Adjusted	1,343	699.8	22.84	300	149.9	10.22	117	71.5	7.73	83	25.0	3.24	81	40.1	5.15	131	91.6	8.86
California																		
Observed	79,013	310.6	1.15	23,487	89.2	0.60	6,633	27.9	0.36	5,433	17.7	0.25	5,684	21.4	0.29	4,552	21.4	0.33
Delay Adjusted	80,455	316.4	1.17	23,802	90.4	0.61	6,710	28.2	0.36	5,489	17.9	0.25	5,725	21.5	0.29	4,619	21.7	0.33
Greater California																		
Observed	40,857	314.6	1.63	12,442	92.2	0.86	3,400	28.3	0.51	2,921	18.2	0.35	2,687	19.9	0.40	2,508	23.5	0.49
Delay Adjusted	41,573	320.3	1.66	12,621	93.5	0.87	3,438	28.6	0.51	2,953	18.4	0.36	2,706	20.0	0.40	2,542	23.9	0.49
Greater Bay																		
Observed	10,359	329.2	3.36	3,065	94.7	1.77	848	28.4	1.01	648	16.4	0.67	746	23.2	0.88	627	23.6	0.96
Delay Adjusted	10,563	335.7	3.43	3,117	96.3	1.80	858	28.8	1.02	653	16.6	0.67	753	23.4	0.88	639	24.0	0.98
Los Angeles																		
Observed	27,783	298.8	1.85	7,976	83.0	0.95	2,384	27.1	0.57	1,864	17.3	0.41	2,251	22.9	0.50	1,416	17.7	0.48
Delay Adjusted	28,304	304.5	1.89	8,061	83.8	0.96	2,413	27.4	0.58	1,883	17.5	0.42	2,265	23.1	0.50	1,438	18.0	0.49
Colorado																		
Observed	6,302	354.9	4.69	1,910	104.2	2.47	516	32.1	1.47	512	23.0	1.05	373	19.5	1.04	528	35.6	1.60
Delay Adjusted	6,433	362.3	4.79	1,919	104.7	2.49	520	32.3	1.48	517	23.2	1.06	375	19.6	1.04	532	35.9	1.61
Connecticut																		
Observed	3,753	439.1	7.75	1,142	127.5	4.03	294	36.7	2.31	344	31.6	1.79	247	28.3	1.93	279	41.3	2.61
Delay Adjusted	3,980	465.8	8.22	1,229	137.3	4.34	307	38.3	2.41	349	32.1	1.82	264	30.2	2.06	285	42.1	2.67

Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Delay Adjusted Counts and Rates[1] for the Five Most Commonly Diagnosed Cancers
Selected Areas in the NAACCR United States Combined[2], 2010-2014, Hispanic/Latina, All Races, Females

	All Sites			Breast			Colon & Rectum			Thyroid			Uterus Corpus & Uterus, NOS[3]			Lung & Bronchus		
	Count	*Rate*	*S.E.[3]*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
Delaware																		
Observed	373	369.1	21.35	102	100.0	11.00	21	26.9	6.29	40	28.2	4.73	28	26.6	5.54	30	38.4	7.39
Delay Adjusted	399	394.7	22.83	104	101.5	11.17	21	27.5	6.44	41	28.9	4.85	28	26.9	5.58	31	39.3	7.56
District of Columbia																		
Observed	300	275.0	16.86	80	70.8	8.22	20	22.8	5.43	29	21.4	4.09	18	15.6	3.76	19	23.7	5.60
Delay Adjusted	312	285.0	17.43	85	74.9	8.69	21	23.3	5.55	31	22.6	4.32	18	15.9	3.84	19	24.1	5.70
Florida																		
Observed	38,723	335.5	1.71	11,343	98.1	0.92	3,823	33.2	0.54	2,716	23.5	0.45	2,778	23.7	0.45	3,136	27.3	0.49
Delay Adjusted	40,818	353.6	1.80	12,053	104.2	0.98	3,965	34.4	0.56	2,733	23.7	0.46	2,867	24.5	0.47	3,161	27.5	0.49
Georgia																		
Observed	3,124	307.5	6.42	1,012	93.2	3.35	209	26.2	2.06	259	17.4	1.24	197	19.2	1.53	173	24.0	1.98
Delay Adjusted	3,256	320.7	6.70	1,038	95.7	3.44	212	26.6	2.09	261	17.5	1.25	198	19.3	1.54	175	24.2	2.00
Hawaii																		
Observed	812	442.6	16.48	282	151.6	9.51	71	42.3	5.29	40	14.9	2.44	45	22.2	3.41	82	51.8	6.00
Delay Adjusted	955	520.2	19.39	346	186.0	11.68	71	42.5	5.31	40	15.0	2.45	45	22.3	3.42	90	56.7	6.57
Idaho																		
Observed	732	323.1	13.47	200	89.1	7.05	55	26.4	3.97	68	21.5	2.86	57	22.5	3.20	43	23.9	3.93
Delay Adjusted	752	332.1	13.86	211	93.8	7.41	56	26.8	4.03	68	21.7	2.89	58	23.0	3.26	44	24.4	4.02
Illinois																		
Observed	10,429	336.8	3.58	2,928	89.5	1.76	795	29.0	1.11	873	21.1	0.76	843	24.9	0.91	663	26.2	1.08
Delay Adjusted	10,639	343.7	3.65	2,994	91.5	1.80	810	29.6	1.13	888	21.5	0.78	851	25.1	0.92	672	26.6	1.09
Indiana																		
Observed	1,588	312.1	8.63	438	83.7	4.33	137	30.2	2.79	114	16.6	1.71	117	21.0	2.08	123	30.9	2.95
Delay Adjusted	1,642	322.7	8.93	449	85.7	4.44	140	30.9	2.85	116	16.8	1.74	119	21.5	2.13	126	31.5	3.01
Iowa																		
Observed	514	264.3	13.41	142	67.1	6.13	36	25.1	4.73	38	13.4	2.41	40	16.9	2.90	35	28.1	5.03
Delay Adjusted	521	268.3	13.62	143	67.5	6.17	36	25.3	4.77	38	13.6	2.44	40	16.9	2.91	35	28.5	5.09
Kansas																		
Observed	1,294	320.6	9.74	347	84.2	4.85	80	22.9	2.74	138	27.5	2.54	87	20.9	2.37	91	28.9	3.19
Delay Adjusted	^	^	^	^	^	^	^	^	^	^	^	^	^	^	^	^	^	^

1. Rates are per 100,000 population and are age-adjusted by five-year age groups to the 2000 U.S. standard population based on single years of age.
2. See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.
3. Abbreviations: NOS - Not Otherwise Specified; S.E. - Standard Error

^ There are insufficient historical registry data to calculate delay factors necessary for delay adjusted rates.

\- Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for All Sites combined.

Note: Delay adjusted counts are rounded to the nearest whole number. A small registry and/or rare cancer site may have the same delay adjusted count and observed count, but this is just due to rounding to whole counts. For more information about cancer incidence rates adjusted for reporting delay see https://surveillance.cancer.gov/delay/.

Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Delay Adjusted Counts and Rates[1] for the Five Most Commonly Diagnosed Cancers
Selected Areas in the NAACCR United States Combined[2], 2010-2014, Hispanic/Latina, All Races, Females

	All Sites			Breast			Colon & Rectum			Thyroid			Uterus Corpus & Uterus, NOS[3]			Lung & Bronchus		
	Count	*Rate*	*S.E.*[3]	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
Kentucky																		
Observed	304	199.7	12.70	82	54.2	6.51	25	18.1	3.98	21	8.6	2.05	15	10.4	2.93	33	29.2	5.27
Delay Adjusted	310	203.4	12.94	83	54.7	6.57	25	18.4	4.05	21	8.7	2.07	15	10.5	2.96	33	29.5	5.34
Louisiana																		
Observed	1,144	322.7	9.73	323	91.5	5.15	108	31.5	3.09	71	17.2	2.09	74	20.2	2.38	105	32.3	3.19
Delay Adjusted	1,210	341.4	10.30	332	94.1	5.30	109	31.7	3.11	72	17.4	2.12	74	20.4	2.40	110	33.9	3.35
Maine																		
Observed	66	222.0	28.30	19	56.6	13.26	-	-	-	-	-	-	6	17.6	7.24	11	44.1	13.62
Delay Adjusted	67	225.3	28.71	19	57.6	13.51	-	-	-	-	-	-	6	17.8	7.30	11	44.5	13.75
Maryland																		
Observed	2,239	299.2	6.94	703	91.8	3.75	157	22.7	1.95	200	19.8	1.50	150	19.1	1.68	137	24.6	2.22
Delay Adjusted	2,372	317.2	7.37	717	93.6	3.82	160	23.1	1.98	207	20.5	1.55	151	19.3	1.70	140	25.2	2.27
Massachusetts																		
Observed	3,564	316.5	5.74	1,022	87.2	2.90	270	27.0	1.77	390	26.9	1.43	237	20.5	1.40	258	28.1	1.86
Delay Adjusted	3,580	318.0	5.77	1,034	88.3	2.94	276	27.6	1.81	394	27.2	1.45	240	20.7	1.42	262	28.6	1.89
Michigan																		
Observed	2,004	299.1	7.07	547	80.0	3.58	150	24.2	2.07	143	16.7	1.49	173	24.5	1.94	178	31.2	2.43
Delay Adjusted	2,116	316.0	7.47	561	82.0	3.67	159	25.6	2.19	145	16.9	1.51	175	24.8	1.96	181	31.8	2.48
Detroit																		
Observed	1,207	477.7	14.44	315	125.3	7.37	96	39.6	4.19	75	23.0	2.78	106	39.1	3.95	123	56.2	5.25
Delay Adjusted	1,318	521.6	15.76	317	126.2	7.42	105	43.3	4.60	76	23.2	2.81	107	39.3	3.97	124	56.5	5.27
Mississippi																		
Observed	197	188.5	14.49	46	41.4	6.44	14	18.5	5.14	15	11.7	3.23	10	8.8	2.93	19	22.1	5.24
Delay Adjusted	204	195.6	15.06	47	42.2	6.56	14	18.7	5.19	15	12.0	3.30	10	8.8	2.94	19	22.4	5.31
Missouri																		
Observed	879	284.4	10.27	250	78.9	5.28	59	22.3	3.04	66	17.4	2.28	64	18.0	2.35	86	34.7	3.88
Delay Adjusted	948	306.8	11.08	280	88.4	5.92	61	22.9	3.13	67	17.6	2.31	65	18.2	2.37	88	35.3	3.95
Montana																		
Observed	211	431.3	31.10	66	134.3	17.13	16	33.0	8.57	12	20.5	6.41	9	20.4	7.03	25	56.8	11.85
Delay Adjusted	219	447.8	32.29	68	138.6	17.67	17	34.2	8.90	12	20.7	6.47	9	21.0	7.23	25	58.0	12.09
Nebraska																		
Observed	690	349.7	15.16	188	93.4	7.67	55	32.5	4.90	51	16.2	2.56	42	20.4	3.51	61	39.6	5.48
Delay Adjusted	699	354.4	15.37	189	93.9	7.71	56	33.3	5.02	52	16.4	2.60	42	20.5	3.52	63	40.6	5.63

Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Delay Adjusted Counts and Rates[1] for the Five Most Commonly Diagnosed Cancers
Selected Areas in the NAACCR United States Combined[2], 2010-2014, Hispanic/Latina, All Races, Females

	All Sites			Breast			Colon & Rectum			Thyroid			Uterus Corpus & Uterus, NOS[3]			Lung & Bronchus		
	Count	*Rate*	*S.E.[3]*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
New Hampshire																		
Observed	220	340.2	25.05	60	92.2	12.88	20	31.7	7.70	28	33.4	6.62	12	18.5	5.51	18	36.5	9.16
Delay Adjusted	225	348.3	25.66	61	93.3	13.02	20	32.0	7.75	28	33.8	6.73	12	18.7	5.57	18	37.3	9.38
New Jersey																		
Observed	11,452	358.6	3.48	3,299	98.7	1.77	1,010	34.6	1.13	953	24.9	0.83	828	25.1	0.90	802	28.4	1.04
Delay Adjusted	11,946	374.1	3.64	3,449	103.2	1.86	1,053	36.1	1.18	978	25.5	0.85	859	26.0	0.93	824	29.2	1.06
New York																		
Observed	27,562	355.6	2.18	7,996	101.3	1.15	2,262	30.6	0.65	2,259	26.4	0.56	2,088	26.1	0.58	2,126	29.3	0.65
Delay Adjusted	27,783	358.5	2.20	8,027	101.7	1.15	2,275	30.7	0.66	2,268	26.5	0.56	2,097	26.2	0.58	2,145	29.5	0.65
North Carolina																		
Observed	2,724	305.7	6.98	755	82.5	3.49	172	22.5	2.00	251	17.1	1.24	184	20.8	1.76	177	29.3	2.44
Delay Adjusted	2,811	315.7	7.21	807	88.2	3.73	176	23.0	2.05	259	17.6	1.28	186	21.1	1.78	179	29.7	2.47
North Dakota																		
Observed	37	200.5	36.43	12	58.5	18.17	-	-	-	-	-	-	-	-	-	-	-	-
Delay Adjusted	38	206.4	37.46	12	59.1	18.33	-	-	-	-	-	-	-	-	-	-	-	-
Ohio																		
Observed	1,271	232.0	6.87	357	64.3	3.55	115	22.4	2.19	91	12.7	1.41	91	16.4	1.78	115	24.1	2.34
Delay Adjusted	1,289	235.4	6.98	365	65.8	3.63	117	22.7	2.22	93	13.0	1.44	92	16.6	1.81	118	24.7	2.39
Oklahoma																		
Observed	1,413	353.4	10.71	395	96.9	5.43	115	32.9	3.43	82	14.8	1.90	77	17.2	2.14	104	34.6	3.67
Delay Adjusted	1,428	357.3	10.83	402	98.7	5.53	116	33.1	3.46	83	15.1	1.92	78	17.5	2.17	106	35.1	3.73
Oregon																		
Observed	1,958	362.5	9.38	532	97.0	4.72	140	28.2	2.67	156	18.2	1.64	147	24.6	2.26	170	44.3	3.66
Delay Adjusted	1,991	369.0	9.55	539	98.2	4.78	142	28.6	2.70	157	18.3	1.65	148	24.9	2.28	174	45.4	3.75
Pennsylvania																		
Observed	3,880	330.1	5.74	1,040	86.5	2.87	298	27.4	1.71	351	22.7	1.28	297	25.5	1.57	291	29.9	1.86
Delay Adjusted	4,182	355.8	6.19	1,116	92.8	3.08	316	29.0	1.81	380	24.6	1.39	304	26.1	1.61	307	31.5	1.96
Rhode Island																		
Observed	592	281.1	12.63	191	85.2	6.68	43	22.5	3.73	43	15.6	2.52	42	20.6	3.47	36	21.2	3.77
Delay Adjusted	614	291.8	13.12	203	90.6	7.11	44	22.8	3.78	43	15.7	2.54	43	21.0	3.53	37	21.6	3.83

1. Rates are per 100,000 population and are age-adjusted by five-year age groups to the 2000 U.S. standard population based on single years of age.
2. See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.
3. Abbreviations: NOS - Not Otherwise Specified; S.E. - Standard Error

^ There are insufficient historical registry data to calculate delay factors necessary for delay adjusted rates.

\- Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for All Sites combined.

Note: Delay adjusted counts are rounded to the nearest whole number. A small registry and/or rare cancer site may have the same delay adjusted count and observed count, but this is just due to rounding to whole counts. For more information about cancer incidence rates adjusted for reporting delay see https://surveillance.cancer.gov/delay/.

Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Delay Adjusted Counts and Rates[1] for the Five Most Commonly Diagnosed Cancers
Selected Areas in the NAACCR United States Combined[2], 2010-2014, Hispanic/Latina, All Races, Females

	All Sites			Prostate			Lung & Bronchus			Thyroid			Uterus Corpus & Uterus, NOS[3]			Lung & Bronchus		
	Count	*Rate*	*S.E.*[3]	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
South Carolina																		
Observed	809	282.1	11.27	266	89.6	6.10	76	28.3	3.57	62	15.1	2.07	38	13.4	2.41	60	29.8	4.17
Delay Adjusted	836	291.4	11.64	272	91.7	6.25	79	29.3	3.68	64	15.5	2.13	39	13.7	2.46	61	30.3	4.25
South Dakota																		
Observed	60	182.5	25.19	16	46.8	12.24	11	36.1	11.52	-	-	-	-	-	-	-	-	-
Delay Adjusted	63	190.7	26.31	16	47.6	12.44	11	37.3	11.87	-	-	-	-	-	-	-	-	-
Tennessee																		
Observed	830	236.6	9.50	228	66.5	4.91	55	18.9	2.81	64	11.8	1.68	55	15.1	2.27	61	25.8	3.54
Delay Adjusted	835	238.1	9.56	231	67.3	4.96	55	19.0	2.84	65	12.0	1.70	55	15.2	2.28	62	26.1	3.58
Texas																		
Observed	55,396	315.4	1.39	15,884	88.0	0.72	4,624	27.8	0.42	3,623	17.5	0.30	3,982	21.3	0.35	3,196	21.4	0.39
Delay Adjusted	57,173	325.5	1.44	16,474	91.3	0.75	4,709	28.4	0.43	3,766	18.2	0.31	4,055	21.7	0.35	3,248	21.7	0.39
Utah																		
Observed	1,729	367.4	9.98	479	101.8	5.14	124	30.8	3.05	207	31.0	2.37	125	25.7	2.52	110	33.0	3.37
Delay Adjusted	1,776	377.8	10.27	481	102.2	5.16	125	31.1	3.08	208	31.2	2.38	126	25.8	2.53	112	33.7	3.44
Vermont																		
Observed	39	231.8	39.48	15	88.1	23.96	-	-	-	-	-	-	7	45.7	18.62	-	-	-
Delay Adjusted	40	235.5	40.19	15	88.5	24.08	-	-	-	-	-	-	7	45.9	18.72	-	-	-
Virginia																		
Observed	2,788	290.6	6.27	823	80.6	3.13	209	25.1	1.95	314	21.9	1.34	161	16.2	1.41	166	23.0	1.96
Delay Adjusted	2,879	299.8	6.46	853	83.6	3.24	217	26.0	2.01	317	22.1	1.35	163	16.4	1.43	168	23.3	1.98
Washington																		
Observed	3,272	345.8	6.96	930	93.6	3.41	227	27.9	2.07	290	20.4	1.37	195	20.4	1.62	220	32.0	2.36
Delay Adjusted	3,472	367.2	7.39	1,017	102.5	3.74	235	28.8	2.14	294	20.7	1.39	197	20.5	1.63	229	33.4	2.46
Seattle																		
Observed	2,224	456.9	11.18	697	137.3	5.84	143	34.4	3.24	195	25.6	2.10	133	27.9	2.70	172	47.9	4.00
Delay Adjusted	2,405	494.1	12.10	790	155.7	6.62	149	35.9	3.37	196	25.7	2.11	134	28.1	2.72	180	50.2	4.19
West Virginia																		
Observed	60	140.1	18.45	14	34.5	9.40	-	-	-	-	-	-	-	-	-	9	21.7	7.30
Delay Adjusted	62	144.0	18.95	14	35.5	9.66	-	-	-	-	-	-	-	-	-	9	21.9	7.39
Wisconsin																		
Observed	1,385	327.4	9.93	376	90.2	5.14	101	28.4	3.11	131	20.1	2.00	111	25.1	2.57	104	32.4	3.41
Delay Adjusted	1,439	341.7	10.42	390	93.7	5.34	105	29.5	3.23	135	20.9	2.08	117	26.5	2.74	107	33.4	3.51

Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Delay Adjusted Counts and Rates[1] for the Five Most Commonly Diagnosed Cancers
Selected Areas in the NAACCR United States Combined[2], 2010-2014, Hispanic/Latina, All Races, Females

	All Sites			Prostate			Lung & Bronchus			Thyroid			Uterus Corpus & Uterus, NOS[3]			Lung & Bronchus		
	Count	*Rate*	*S.E.*[3]	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
Wyoming																		
Observed	272	304.9	19.01	76	85.5	10.00	16	19.7	5.03	15	13.7	3.68	14	14.8	4.03	22	28.4	6.18
Delay Adjusted	275	307.9	19.20	77	86.1	10.08	16	19.7	5.05	15	13.8	3.71	14	15.0	4.09	22	28.5	6.21

1. Rates are per 100,000 population and are age-adjusted by five-year age groups to the 2000 U.S. standard population based on single years of age.
2. See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.
3. Abbreviations: NOS - Not Otherwise Specified; S.E. - Standard Error

^ There are insufficient historical registry data to calculate delay factors necessary for delay adjusted rates.

\- Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for All Sites combined.

Note: Delay adjusted counts are rounded to the nearest whole number. A small registry and/or rare cancer site may have the same delay adjusted count and observed count, but this is just due to rounding to whole counts. For more information about cancer incidence rates adjusted for reporting delay see https://surveillance.cancer.gov/delay/.

NAACCR U.S. Combined Delay Adjusted Incidence 2010-2014, Non-Hispanic White Combined

UNITED STATES

V-75 Combined Delay Adjusted Incidence – Non-Hispanic White

Average Annual Age-adjusted Cancer Incidence Delay Adjusted Counts and Rates*, Selected Sites
NAACCR United States Combined[1], 2010-2014, Non-Hispanic White

	MALES (ALL RACES)						FEMALES (ALL RACES)					
	Observed			Delay Adjusted			Observed			Delay Adjusted		
SITES	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
All Sites (invasive & in situ bladder)	2,961,072	500.2	0.30	3,037,937	513.2	0.31	2,849,465	429.1	0.26	2,923,925	440.3	0.27
Oral Cavity & Pharynx	115,165	18.8	0.06	118,226	19.3	0.06	45,991	6.8	0.03	47,175	7.0	0.03
Digestive												
Esophagus	52,340	8.6	0.04	53,198	8.8	0.04	13,113	1.8	0.02	13,328	1.8	0.02
Stomach	46,073	7.8	0.04	46,761	8.0	0.04	24,856	3.5	0.02	25,221	3.6	0.02
Colon & Rectum	263,005	45.2	0.09	267,433	45.9	0.09	241,669	34.5	0.07	245,691	35.1	0.07
Liver & IBD[2]	62,694	10.0	0.04	64,260	10.2	0.04	24,221	3.4	0.02	24,841	3.5	0.02
Pancreas	85,017	14.3	0.05	86,376	14.5	0.05	78,507	10.8	0.04	79,737	10.9	0.04
Respiratory												
Larynx	37,392	6.1	0.03	38,101	6.2	0.03	10,060	1.5	0.01	10,250	1.5	0.02
Lung & bronchus	450,901	75.9	0.12	457,728	77.0	0.12	408,779	57.5	0.09	414,999	58.4	0.09
Skin												
Melanomas of the skin	187,736	32.8	0.08	195,582	34.1	0.08	128,264	21.3	0.06	133,435	22.1	0.06
Breast (female cases only)	-	-	-	-	-	-	841,438	128.7	0.15	856,201	130.9	0.15
Female Genital												
Cervix uteri	-	-	-	-	-	-	36,848	7.0	0.04	37,481	7.1	0.04
Corpus & uterus, NOS[2]	-	-	-	-	-	-	182,636	26.6	0.06	185,236	26.9	0.07
Ovary	-	-	-	-	-	-	79,140	12.0	0.04	81,049	12.3	0.05
Male Genital												
Prostate	679,454	107.1	0.13	707,793	111.6	0.14	-	-	-	-	-	-
Testis	30,642	6.8	0.04	31,155	7.0	0.04	-	-	-	-	-	-
Urinary												
Bladder (incl. *in situ*)	229,998	39.8	0.08	236,465	40.9	0.09	70,592	9.7	0.04	72,566	10.0	0.04
Kidney & renal pelvis	130,539	22.1	0.06	133,802	22.6	0.06	75,325	11.3	0.04	77,192	11.6	0.04
Brain & Other Nervous System	47,107	8.7	0.04	48,325	8.9	0.04	37,072	6.3	0.03	38,013	6.4	0.04
Endocrine												
Thyroid gland	43,776	8.0	0.04	44,623	8.2	0.04	120,802	22.7	0.07	123,122	23.1	0.07
Lymphomas												
Hodgkin Lymphoma	16,069	3.3	0.03	16,372	3.3	0.03	12,933	2.6	0.02	13,170	2.6	0.02
Non-Hodgkin Lymphoma	136,941	23.9	0.07	141,681	24.8	0.07	112,218	16.4	0.05	116,053	16.9	0.05
Multiple Myeloma	42,996	7.3	0.04	46,315	7.9	0.04	32,044	4.5	0.03	34,504	4.8	0.03
Leukemias	101,723	18.2	0.06	112,232	20.1	0.06	72,000	11.0	0.04	79,338	12.1	0.05

* Rates are per 100,000 population and are age-adjusted by five-year age groups to the 2000 U.S. standard population based on single years of age and to standard.

\- Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for All Sites combined.

[1] NAACCR United States Combined includes registries for which a U.S. average delay adjustment factor was calculated. See technical notes.

[2] Abbreviations: IBD – Intrahepatic Bile Duct; NOS - Not Otherwise Specified

Note: Delay adjusted counts are rounded to the nearest whole number. A small registry and/or rare cancer site may have the same delay adjusted count and observed count, but this is just due to rounding to whole counts. For more information about cancer incidence rates adjusted for reporting delay see https://surveillance.cancer.gov/delay/.

Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Delay Adjusted Counts and Rates[1] for the Five Most Commonly Diagnosed Cancers
Selected Areas in the NAACCR United States Combined[2], 2010-2014, Non-Hispanic White, Males

	All Sites			Prostate			Lung & Bronchus			Colon & Rectum			Bladder			Melanoma of the Skin		
	Count	*Rate*	*S.E.*[3]	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
NAACCR U.S. Combined																		
Observed	2,961,072	500.2	0.30	679,454	107.1	0.13	450,901	75.9	0.12	263,005	45.2	0.09	229,998	39.8	0.08	187,736	32.8	0.08
Delay Adjusted	3,037,937	513.2	0.31	707,793	111.6	0.14	457,728	77.0	0.12	267,433	45.9	0.09	236,465	40.9	0.09	195,582	34.1	0.08
Alabama																		
Observed	50,731	517.4	2.36	11,375	108.3	1.04	9,248	93.1	0.99	4,837	50.2	0.74	3,567	37.6	0.65	3,269	34.8	0.63
Delay Adjusted	51,439	524.6	2.39	11,695	111.4	1.07	9,297	93.6	1.00	4,874	50.6	0.75	3,607	38.0	0.65	3,356	35.7	0.64
Alaska																		
Observed	5,219	437.6	6.85	1,267	95.4	3.03	678	61.5	2.66	442	39.0	2.11	388	37.3	2.12	238	19.4	1.39
Delay Adjusted	5,426	455.1	7.13	1,340	101.0	3.21	696	63.2	2.74	455	40.1	2.18	395	38.0	2.16	262	21.4	1.53
Arizona																		
Observed	57,067	419.7	1.81	11,759	80.6	0.76	8,519	60.1	0.66	5,142	38.4	0.55	5,030	36.0	0.52	3,994	30.5	0.50
Delay Adjusted	59,746	439.4	1.89	12,684	86.9	0.82	8,720	61.5	0.68	5,396	40.3	0.58	5,258	37.7	0.54	4,339	33.1	0.54
Arkansas																		
Observed	34,375	504.2	2.78	7,865	108.3	1.24	6,749	97.0	1.20	3,246	48.4	0.87	2,491	37.3	0.76	1,821	28.0	0.67
Delay Adjusted	34,756	509.8	2.81	8,018	110.4	1.26	6,794	97.7	1.21	3,268	48.8	0.88	2,515	37.7	0.77	1,872	28.7	0.69
California																		
Observed	250,684	493.7	1.01	59,883	110.1	0.46	28,555	56.3	0.34	21,396	42.5	0.30	19,685	39.4	0.29	22,302	45.1	0.31
Delay Adjusted	255,331	502.9	1.03	61,962	114.0	0.48	28,904	57.0	0.34	21,609	43.0	0.30	20,056	40.1	0.29	23,035	46.5	0.32
Greater California																		
Observed	157,767	493.9	1.28	37,199	108.4	0.57	19,127	59.6	0.44	13,518	42.8	0.38	12,277	39.1	0.36	13,945	45.0	0.39
Delay Adjusted	161,624	505.9	1.31	38,761	113.0	0.60	19,418	60.5	0.45	13,715	43.4	0.38	12,598	40.1	0.37	14,523	46.8	0.41
Greater Bay																		
Observed	47,919	503.2	2.37	12,314	120.0	1.11	4,470	48.1	0.74	3,752	39.8	0.67	3,570	38.4	0.66	4,968	53.5	0.78
Delay Adjusted	48,086	505.0	2.38	12,388	120.8	1.12	4,489	48.3	0.74	3,756	39.8	0.67	3,587	38.6	0.66	5,068	54.6	0.80
Los Angeles																		
Observed	44,873	482.6	2.33	10,328	105.2	1.05	4,942	53.0	0.76	4,118	44.4	0.70	3,833	41.4	0.68	3,386	37.0	0.65
Delay Adjusted	45,490	489.2	2.36	10,767	109.7	1.10	4,979	53.4	0.77	4,129	44.5	0.71	3,865	41.7	0.68	3,441	37.6	0.66
Colorado																		
Observed	43,690	431.1	2.14	10,993	100.1	0.99	4,717	48.7	0.73	3,592	36.6	0.63	3,231	34.2	0.62	3,324	33.3	0.60
Delay Adjusted	44,927	443.3	2.20	11,766	107.5	1.07	4,756	49.1	0.74	3,634	37.0	0.64	3,353	35.5	0.65	3,416	34.2	0.61
Connecticut																		
Observed	41,355	514.5	2.60	9,638	111.6	1.16	5,668	70.2	0.95	3,425	43.3	0.76	3,959	49.5	0.80	2,420	31.2	0.66
Delay Adjusted	42,251	525.6	2.66	9,915	114.9	1.20	5,744	71.1	0.96	3,462	43.8	0.77	4,035	50.4	0.82	2,485	32.1	0.67

Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Delay Adjusted Counts and Rates[1] for the Five Most Commonly Diagnosed Cancers
Selected Areas in the NAACCR United States Combined[2], 2010-2014, Non-Hispanic White, Males

	All Sites			Prostate			Lung & Bronchus			Colon & Rectum			Bladder			Melanoma of the Skin		
	Count	*Rate*	*S.E.*[3]	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
Delaware																		
Observed	11,189	565.8	5.50	2,755	128.3	2.49	1,671	83.4	2.08	839	43.4	1.54	911	46.4	1.57	949	49.6	1.66
Delay Adjusted	11,502	581.6	5.66	2,886	134.4	2.61	1,707	85.2	2.12	849	43.9	1.56	943	48.0	1.62	982	51.4	1.72
District of Columbia																		
Observed	1,892	387.0	9.29	504	101.6	4.66	164	37.0	2.98	127	25.9	2.41	121	26.3	2.47	119	24.3	2.33
Delay Adjusted	1,990	407.2	9.78	523	105.5	4.84	167	37.7	3.04	131	26.7	2.48	124	26.9	2.53	125	25.6	2.47
Florida																		
Observed	208,899	479.8	1.09	43,335	93.1	0.45	34,836	76.2	0.42	17,716	41.2	0.32	16,827	36.6	0.29	15,825	37.8	0.32
Delay Adjusted	216,742	498.0	1.13	45,788	98.3	0.48	35,099	76.8	0.42	18,178	42.3	0.33	17,424	37.9	0.30	16,953	40.5	0.34
Georgia																		
Observed	81,616	524.8	1.90	18,487	110.2	0.84	13,373	87.1	0.78	7,240	47.2	0.57	5,573	38.0	0.53	7,041	46.4	0.57
Delay Adjusted	83,057	534.1	1.94	18,884	112.6	0.85	13,515	88.0	0.79	7,312	47.6	0.58	5,669	38.6	0.54	7,273	48.0	0.59
Hawaii																		
Observed	5,635	512.3	7.31	1,072	87.7	2.85	586	52.9	2.33	459	41.2	2.05	401	38.7	2.03	859	81.6	2.96
Delay Adjusted	5,747	522.5	7.46	1,104	90.3	2.94	593	53.5	2.35	463	41.6	2.07	412	39.7	2.09	895	85.0	3.08
Idaho																		
Observed	17,781	467.0	3.59	4,691	115.9	1.73	2,101	56.0	1.25	1,482	39.4	1.05	1,411	38.5	1.04	1,287	34.4	0.99
Delay Adjusted	18,271	479.9	3.69	4,929	121.8	1.82	2,144	57.1	1.27	1,504	40.0	1.07	1,489	40.6	1.10	1,338	35.8	1.02
Illinois																		
Observed	125,155	521.1	1.51	28,724	112.1	0.68	19,537	81.8	0.60	12,361	52.1	0.48	9,833	42.1	0.43	6,648	28.5	0.36
Delay Adjusted	127,507	530.9	1.53	29,737	116.2	0.70	19,752	82.7	0.60	12,467	52.5	0.48	10,039	43.0	0.44	6,886	29.5	0.37
Indiana																		
Observed	71,727	485.5	1.86	14,283	90.7	0.78	13,240	90.1	0.80	7,090	48.7	0.59	5,384	37.8	0.53	3,311	22.9	0.41
Delay Adjusted	74,197	502.3	1.92	15,086	95.8	0.82	13,532	92.1	0.82	7,278	50.0	0.61	5,625	39.5	0.55	3,523	24.4	0.44
Iowa																		
Observed	41,938	518.4	2.58	9,538	110.8	1.15	6,403	78.8	1.00	4,132	52.0	0.82	3,095	38.6	0.70	2,316	29.7	0.63
Delay Adjusted	42,736	528.3	2.63	9,819	114.1	1.19	6,485	79.8	1.01	4,162	52.4	0.83	3,164	39.5	0.72	2,383	30.6	0.65
Kansas																		
Observed	32,993	505.5	2.84	8,015	116.6	1.33	4,776	73.1	1.08	3,010	46.8	0.87	2,558	39.9	0.80	1,964	31.3	0.72
Delay Adjusted	^	^	^	^	^	^	^	^	^	^	^	^	^	^	^	^	^	^

1. Rates are per 100,000 population and are age-adjusted by five-year age groups to the 2000 U.S. standard population based on single years of age.
2. See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.
3. Abbreviations: S.E. - Standard Error

^ There are insufficient historical registry data to calculate delay factors necessary for delay adjusted rates.

\- Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for All Sites combined.

Note: Delay adjusted counts are rounded to the nearest whole number. A small registry and/or rare cancer site may have the same delay adjusted count and observed count, but this is just due to rounding to whole counts. For more information about cancer incidence rates adjusted for reporting delay see https://surveillance.cancer.gov/delay/.

Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Delay Adjusted Counts and Rates[1] for the Five Most Commonly Diagnosed Cancers
Selected Areas in the NAACCR United States Combined[2], 2010-2014, Non-Hispanic White, Males

	All Sites			Prostate			Lung & Bronchus			Colon & Rectum			Bladder			Melanoma of the Skin		
	Count	*Rate*	*S.E.*[3]	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
Kentucky																		
Observed	60,771	578.0	2.42	11,920	106.1	1.00	12,387	117.7	1.09	6,166	59.5	0.78	4,109	41.3	0.66	3,333	32.8	0.59
Delay Adjusted	62,145	591.0	2.47	12,509	111.2	1.05	12,532	119.0	1.10	6,281	60.6	0.80	4,218	42.4	0.68	3,425	33.7	0.60
Louisiana																		
Observed	44,556	558.8	2.72	10,920	128.2	1.26	6,880	86.9	1.07	4,203	53.2	0.84	2,965	38.7	0.73	2,420	31.7	0.66
Delay Adjusted	44,955	563.9	2.74	11,021	129.3	1.27	6,920	87.4	1.08	4,215	53.4	0.85	2,995	39.0	0.73	2,454	32.2	0.67
Maine																		
Observed	20,375	505.6	3.65	4,332	98.3	1.53	3,435	84.9	1.48	1,689	42.9	1.07	1,867	47.7	1.13	1,105	28.1	0.87
Delay Adjusted	20,707	513.8	3.71	4,450	101.0	1.57	3,472	85.9	1.50	1,700	43.2	1.08	1,902	48.6	1.15	1,139	28.9	0.90
Maryland																		
Observed	48,752	502.4	2.34	12,000	114.3	1.07	6,557	68.4	0.86	3,957	41.6	0.68	4,016	42.8	0.69	3,887	41.0	0.68
Delay Adjusted	49,972	515.0	2.40	12,498	119.1	1.11	6,702	69.9	0.88	4,012	42.2	0.69	4,211	44.9	0.72	3,907	41.2	0.68
Massachusetts																		
Observed	73,973	492.3	1.86	17,028	105.0	0.82	10,665	71.8	0.71	6,301	42.7	0.55	6,238	42.6	0.55	3,998	27.4	0.44
Delay Adjusted	75,655	503.6	1.90	17,737	109.3	0.86	10,875	73.2	0.72	6,432	43.6	0.56	6,395	43.6	0.56	4,190	28.7	0.46
Michigan																		
Observed	111,897	490.3	1.50	27,507	112.2	0.69	17,273	75.8	0.59	9,369	42.1	0.45	9,033	41.0	0.44	5,446	24.8	0.35
Delay Adjusted	114,620	502.3	1.54	28,318	115.5	0.71	17,740	77.9	0.61	9,579	43.1	0.46	9,253	42.0	0.45	5,601	25.5	0.36
Detroit																		
Observed	41,313	561.2	2.84	10,874	138.9	1.37	5,996	82.9	1.10	3,292	45.4	0.81	3,412	47.9	0.84	1,963	27.4	0.63
Delay Adjusted	41,777	567.5	2.87	11,134	142.2	1.40	6,030	83.4	1.10	3,313	45.7	0.82	3,471	48.7	0.85	1,981	27.6	0.64
Mississippi																		
Observed	28,000	535.4	3.28	6,230	111.0	1.43	5,147	97.8	1.39	2,766	53.7	1.05	1,822	36.0	0.86	1,628	32.3	0.82
Delay Adjusted	28,671	548.2	3.36	6,455	115.0	1.49	5,222	99.2	1.41	2,801	54.3	1.06	1,893	37.4	0.90	1,657	32.9	0.84
Missouri																		
Observed	70,143	490.3	1.89	14,759	96.1	0.80	12,643	88.5	0.80	6,868	49.1	0.60	4,904	35.4	0.51	3,755	27.1	0.45
Delay Adjusted	71,840	502.2	1.94	15,407	100.4	0.84	12,846	89.9	0.81	6,971	49.8	0.61	5,027	36.3	0.53	3,863	27.8	0.46
Montana																		
Observed	13,378	465.3	4.16	3,560	113.7	1.95	1,660	58.7	1.48	1,226	43.0	1.27	1,008	35.9	1.16	845	30.3	1.09
Delay Adjusted	13,944	484.9	4.33	3,772	120.5	2.07	1,696	60.0	1.51	1,267	44.4	1.31	1,061	37.8	1.22	896	32.2	1.16
Nebraska																		
Observed	21,853	497.3	3.43	5,380	116.4	1.61	3,152	72.1	1.30	2,170	49.6	1.09	1,640	38.3	0.96	1,051	24.3	0.77
Delay Adjusted	22,700	516.6	3.57	5,751	124.4	1.73	3,218	73.6	1.33	2,223	50.9	1.11	1,727	40.3	1.01	1,131	26.1	0.83

Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Delay Adjusted Counts and Rates[1] for the Five Most Commonly Diagnosed Cancers
Selected Areas in the NAACCR United States Combined[2], 2010-2014, Non-Hispanic White, Males

	All Sites			Prostate			Lung & Bronchus			Colon & Rectum			Bladder			Melanoma of the Skin		
	Count	*Rate*	*S.E.*[3]	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
New Hampshire																		
Observed	18,749	516.7	3.91	4,864	121.7	1.80	2,525	71.8	1.47	1,449	40.9	1.11	1,672	48.4	1.22	1,145	32.6	1.00
Delay Adjusted	19,221	529.7	4.01	5,007	125.2	1.85	2,584	73.5	1.51	1,471	41.5	1.13	1,702	49.3	1.24	1,187	33.8	1.03
New Jersey																		
Observed	89,861	551.6	1.89	22,800	131.7	0.89	11,354	69.6	0.66	7,865	48.8	0.56	7,550	46.9	0.55	5,942	37.2	0.50
Delay Adjusted	92,229	566.1	1.93	23,989	138.5	0.93	11,493	70.4	0.67	7,963	49.4	0.57	7,801	48.4	0.57	6,196	38.8	0.52
New York																		
Observed	190,578	551.0	1.29	46,407	126.5	0.60	25,886	74.4	0.47	15,846	46.5	0.38	16,558	48.4	0.38	10,489	31.1	0.31
Delay Adjusted	194,295	561.7	1.31	47,689	130.1	0.61	26,040	74.9	0.47	16,034	47.0	0.38	16,879	49.3	0.39	11,087	32.9	0.33
North Carolina																		
Observed	97,567	512.0	1.68	22,424	109.1	0.75	16,779	87.8	0.69	7,847	42.1	0.49	7,161	38.8	0.47	7,156	39.1	0.47
Delay Adjusted	100,352	526.6	1.73	23,491	114.3	0.78	17,045	89.2	0.70	7,998	42.9	0.50	7,356	39.9	0.48	7,397	40.4	0.49
North Dakota																		
Observed	8,993	498.6	5.38	2,354	123.9	2.61	1,236	69.5	2.01	947	53.8	1.79	664	37.3	1.47	461	26.2	1.26
Delay Adjusted	9,247	512.7	5.53	2,432	128.0	2.69	1,267	71.2	2.06	973	55.2	1.84	690	38.8	1.53	473	26.9	1.29
Ohio																		
Observed	133,505	488.5	1.37	28,942	98.8	0.59	22,859	83.6	0.56	12,706	47.4	0.43	10,543	39.8	0.40	6,992	26.5	0.32
Delay Adjusted	138,496	506.8	1.42	30,699	104.8	0.63	23,388	85.6	0.58	13,047	48.7	0.44	10,980	41.5	0.41	7,398	28.0	0.34
Oklahoma																		
Observed	39,270	481.1	2.48	8,557	98.8	1.09	7,060	85.4	1.03	3,739	46.6	0.78	2,803	35.0	0.67	2,063	26.2	0.59
Delay Adjusted	39,792	487.5	2.51	8,713	100.6	1.11	7,138	86.4	1.05	3,769	47.0	0.79	2,832	35.4	0.68	2,074	26.3	0.59
Oregon																		
Observed	45,567	465.0	2.25	10,680	100.4	1.00	6,175	63.4	0.83	3,848	40.4	0.67	3,706	38.7	0.65	3,241	34.2	0.62
Delay Adjusted	46,607	475.7	2.30	10,993	103.3	1.03	6,292	64.6	0.84	3,926	41.2	0.68	3,799	39.7	0.67	3,291	34.7	0.63
Pennsylvania																		
Observed	165,634	522.6	1.31	36,108	106.4	0.57	24,940	77.9	0.50	15,486	49.5	0.41	14,418	45.9	0.39	9,119	29.7	0.32
Delay Adjusted	168,604	532.0	1.34	37,359	110.0	0.59	25,203	78.7	0.51	15,629	50.0	0.41	14,646	46.6	0.39	9,417	30.6	0.33
Rhode Island																		
Observed	12,620	510.5	4.67	2,650	100.0	1.99	1,961	79.3	1.83	984	40.3	1.31	1,163	47.0	1.41	736	30.9	1.18
Delay Adjusted	12,893	521.6	4.77	2,762	104.2	2.07	1,994	80.7	1.86	995	40.7	1.33	1,188	48.0	1.44	757	31.8	1.21

1. Rates are per 100,000 population and are age-adjusted by five-year age groups to the 2000 U.S. standard population based on single years of age.
2. See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.
3. Abbreviations: S.E. - Standard Error

^ There are insufficient historical registry data to calculate delay factors necessary for delay adjusted rates.

\- Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for All Sites combined.

Note: Delay adjusted counts are rounded to the nearest whole number. A small registry and/or rare cancer site may have the same delay adjusted count and observed count, but this is just due to rounding to whole counts. For more information about cancer incidence rates adjusted for reporting delay see https://surveillance.cancer.gov/delay/.

Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Delay Adjusted Counts and Rates[1] for the Five Most Commonly Diagnosed Cancers
Selected Areas in the NAACCR United States Combined[2], 2010-2014, Non-Hispanic White, Males

	All Sites			Prostate			Lung & Bronchus			Colon & Rectum			Bladder			Melanoma of the Skin		
	Count	*Rate*	*S.E.*[3]	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
South Carolina																		
Observed	47,655	503.8	2.39	10,510	101.8	1.02	8,222	85.8	0.97	3,933	42.5	0.70	3,605	39.8	0.68	3,377	36.8	0.66
Delay Adjusted	49,381	522.1	2.47	11,034	106.9	1.07	8,429	87.9	1.00	4,050	43.8	0.72	3,736	41.3	0.71	3,506	38.2	0.68
South Dakota																		
Observed	10,467	491.2	4.91	2,667	117.9	2.33	1,454	68.2	1.82	1,072	51.0	1.59	754	36.2	1.34	552	26.8	1.17
Delay Adjusted	10,949	513.9	5.14	2,857	126.4	2.49	1,504	70.5	1.88	1,108	52.7	1.65	815	39.2	1.45	591	28.6	1.26
Tennessee																		
Observed	75,330	521.9	1.96	17,392	111.8	0.87	13,953	96.2	0.84	6,475	46.1	0.59	5,131	37.0	0.53	4,157	30.1	0.48
Delay Adjusted	76,680	531.3	1.99	17,753	114.1	0.89	14,078	97.1	0.84	6,531	46.5	0.60	5,204	37.5	0.54	4,274	30.9	0.49
Texas																		
Observed	169,156	483.9	1.21	37,569	100.6	0.53	26,304	75.6	0.48	15,679	45.5	0.37	11,165	33.4	0.32	8,923	26.3	0.29
Delay Adjusted	175,547	502.2	1.25	40,479	108.6	0.57	26,802	77.0	0.49	15,907	46.2	0.38	11,612	34.7	0.34	9,724	28.7	0.31
Utah																		
Observed	22,671	455.1	3.09	6,737	131.5	1.63	1,544	32.3	0.84	1,709	35.2	0.87	1,438	31.1	0.83	2,554	51.9	1.05
Delay Adjusted	23,025	462.2	3.14	6,895	134.6	1.67	1,562	32.7	0.85	1,719	35.5	0.87	1,458	31.5	0.85	2,592	52.7	1.06
Vermont																		
Observed	8,774	485.1	5.37	1,935	97.6	2.29	1,289	71.4	2.05	705	39.4	1.54	682	39.1	1.54	702	39.7	1.56
Delay Adjusted	8,885	491.2	5.44	1,961	98.9	2.32	1,303	72.2	2.08	714	39.9	1.56	695	39.9	1.57	713	40.3	1.58
Virginia																		
Observed	68,461	446.7	1.76	15,501	93.3	0.77	11,081	73.2	0.71	6,065	40.1	0.53	4,929	33.9	0.50	4,247	28.4	0.45
Delay Adjusted	69,910	456.2	1.79	15,853	95.4	0.79	11,218	74.1	0.72	6,134	40.6	0.54	5,029	34.6	0.51	4,353	29.1	0.46
Washington																		
Observed	74,979	499.2	1.88	18,744	115.5	0.87	9,793	66.7	0.69	6,005	40.6	0.54	5,662	39.5	0.54	5,068	34.5	0.50
Delay Adjusted	76,866	511.8	1.93	19,493	120.2	0.90	9,947	67.8	0.70	6,125	41.4	0.55	5,863	40.9	0.56	5,183	35.3	0.51
Seattle																		
Observed	51,964	525.2	2.38	13,323	125.3	1.12	6,422	67.2	0.86	4,049	41.4	0.67	3,867	41.4	0.68	3,814	39.2	0.65
Delay Adjusted	52,340	529.0	2.39	13,509	127.0	1.14	6,430	67.3	0.86	4,063	41.5	0.67	3,911	41.9	0.69	3,877	39.9	0.67
West Virginia																		
Observed	27,941	516.6	3.19	5,708	97.9	1.33	5,467	99.6	1.38	2,813	53.0	1.03	2,131	40.6	0.90	1,338	26.2	0.74
Delay Adjusted	28,582	528.5	3.26	5,842	100.2	1.36	5,558	101.2	1.41	2,861	53.9	1.05	2,175	41.5	0.92	1,381	27.0	0.76
Wisconsin																		
Observed	71,156	495.2	1.90	17,341	112.4	0.87	9,699	67.7	0.70	5,999	42.6	0.56	5,692	41.0	0.55	3,899	27.8	0.46
Delay Adjusted	75,134	523.5	2.01	18,334	118.9	0.92	10,203	71.3	0.74	6,310	44.8	0.59	6,078	43.7	0.59	4,125	29.4	0.48

Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Delay Adjusted Counts and Rates[1] for the Five Most Commonly Diagnosed Cancers
Selected Areas in the NAACCR United States Combined[2], 2010-2014, Non-Hispanic White, Males

	All Sites			Prostate			Lung & Bronchus			Colon & Rectum			Bladder			Melanoma of the Skin		
	Count	*Rate*	*S.E.*[3]	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
Wyoming																		
Observed	6,494	455.0	5.87	1,744	112.7	2.80	700	50.6	1.98	582	41.9	1.80	504	37.5	1.73	420	31.0	1.57
Delay Adjusted	6,567	460.1	5.93	1,762	113.8	2.83	704	50.8	1.99	586	42.2	1.82	508	37.8	1.74	434	32.0	1.62

1. Rates are per 100,000 population and are age-adjusted by five-year age groups to the 2000 U.S. standard population based on single years of age.
2. See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.
3. Abbreviations: S.E. - Standard Error

^ There are insufficient historical registry data to calculate delay factors necessary for delay adjusted rates.

\- Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for All Sites combined.

Note: Delay adjusted counts are rounded to the nearest whole number. A small registry and/or rare cancer site may have the same delay adjusted count and observed count, but this is just due to rounding to whole counts. For more information about cancer incidence rates adjusted for reporting delay see https://surveillance.cancer.gov/delay/.

Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Delay Adjusted Counts and Rates[1] for the Five Most Commonly Diagnosed Cancers
Selected Areas in the NAACCR United States Combined[2], 2010-2014, Non-Hispanic White, Females

	All Sites			Breast			Lung & Bronchus			Colon & Rectum			Uterus Corpus & Uterus, NOS[3]			Melanoma of the Skin		
	Count	*Rate*	*S.E.*[3]	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
NAACCR U.S. Combined																		
Observed	2,849,465	429.1	0.26	841,438	128.7	0.15	408,779	57.5	0.09	241,669	34.5	0.07	182,636	26.6	0.06	128,264	21.3	0.06
Delay Adjusted	2,923,925	440.3	0.27	856,201	130.9	0.15	414,999	58.4	0.09	245,691	35.1	0.07	185,236	26.9	0.07	133,435	22.1	0.06
Alabama																		
Observed	44,548	399.6	1.96	13,076	118.2	1.07	6,820	56.7	0.70	4,016	35.0	0.57	2,168	18.9	0.42	2,161	21.6	0.49
Delay Adjusted	45,180	405.3	1.99	13,273	120.0	1.09	6,856	57.0	0.70	4,047	35.2	0.57	2,173	19.0	0.42	2,218	22.1	0.50
Alaska																		
Observed	4,622	402.4	6.30	1,508	125.9	3.45	564	53.8	2.42	375	33.4	1.84	341	27.0	1.56	193	16.4	1.24
Delay Adjusted	4,808	418.5	6.55	1,542	128.6	3.53	578	55.2	2.48	386	34.4	1.89	353	28.0	1.62	211	18.0	1.36
Arizona																		
Observed	55,092	385.3	1.75	16,896	120.5	0.99	7,994	49.7	0.57	4,566	30.0	0.47	3,121	21.0	0.40	2,203	17.2	0.40
Delay Adjusted	57,685	403.4	1.83	17,628	125.7	1.03	8,183	50.8	0.58	4,790	31.5	0.49	3,252	21.8	0.41	2,391	18.6	0.43
Arkansas																		
Observed	29,420	388.5	2.35	8,288	110.5	1.26	5,088	62.2	0.89	2,780	35.2	0.69	1,641	21.5	0.55	1,128	16.6	0.52
Delay Adjusted	29,751	392.9	2.38	8,361	111.5	1.27	5,124	62.6	0.89	2,799	35.4	0.69	1,653	21.7	0.55	1,155	17.0	0.53
California																		
Observed	239,215	430.2	0.92	75,781	139.0	0.53	29,335	48.7	0.29	19,897	33.7	0.25	15,188	26.1	0.22	13,601	26.7	0.24
Delay Adjusted	243,708	438.2	0.94	76,271	139.9	0.53	29,699	49.3	0.29	20,090	34.0	0.25	15,266	26.3	0.22	14,028	27.5	0.25
Greater California																		
Observed	149,726	426.8	1.16	47,034	136.1	0.66	19,486	51.2	0.38	12,414	33.5	0.31	9,210	25.2	0.27	8,395	26.2	0.31
Delay Adjusted	153,446	437.4	1.19	47,501	137.3	0.67	19,791	52.0	0.38	12,593	34.0	0.32	9,282	25.4	0.27	8,726	27.2	0.32
Greater Bay																		
Observed	45,461	436.4	2.15	14,552	143.1	1.24	5,055	45.4	0.66	3,675	33.3	0.58	3,033	27.4	0.51	3,191	33.3	0.62
Delay Adjusted	45,624	437.9	2.16	14,552	143.1	1.24	5,077	45.6	0.66	3,679	33.4	0.58	3,035	27.5	0.51	3,256	34.0	0.64
Los Angeles																		
Observed	43,939	434.9	2.18	14,173	144.8	1.27	4,785	43.3	0.64	3,800	34.7	0.59	2,944	28.4	0.54	2,011	21.3	0.50
Delay Adjusted	44,544	440.8	2.21	14,196	145.0	1.27	4,821	43.6	0.65	3,810	34.8	0.59	2,948	28.5	0.54	2,043	21.6	0.51
Colorado																		
Observed	44,193	391.8	1.92	14,484	127.1	1.09	4,883	42.5	0.62	3,450	30.4	0.53	2,746	22.7	0.45	2,299	21.4	0.46
Delay Adjusted	45,418	402.4	1.97	14,658	128.6	1.10	4,923	42.8	0.63	3,488	30.8	0.54	2,774	22.9	0.45	2,364	22.0	0.47
Connecticut																		
Observed	42,968	459.4	2.34	13,019	143.3	1.32	5,998	59.7	0.79	3,409	33.5	0.61	2,889	29.3	0.56	1,738	20.0	0.52
Delay Adjusted	43,910	469.5	2.40	13,201	145.3	1.34	6,079	60.5	0.80	3,447	33.9	0.61	2,907	29.5	0.56	1,785	20.6	0.53

Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Delay Adjusted Counts and Rates[1] for the Five Most Commonly Diagnosed Cancers
Selected Areas in the NAACCR United States Combined[2], 2010-2014, Non-Hispanic White, Females

	All Sites			Breast			Lung & Bronchus			Colon & Rectum			Uterus Corpus & Uterus, NOS[3]			Melanoma of the Skin		
	Count	*Rate*	*S.E.[3]*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
Delaware																		
Observed	10,364	467.0	4.84	3,010	135.8	2.61	1,599	66.2	1.69	771	33.1	1.24	692	29.4	1.16	631	31.2	1.33
Delay Adjusted	10,657	480.1	4.97	3,040	137.2	2.63	1,635	67.7	1.73	781	33.5	1.26	700	29.8	1.18	653	32.3	1.37
District of Columbia																		
Observed	2,002	398.0	9.41	763	157.6	5.95	181	37.8	2.94	131	26.7	2.46	120	24.0	2.28	81	14.0	1.70
Delay Adjusted	2,107	418.9	9.90	804	166.1	6.27	185	38.7	3.00	135	27.5	2.53	125	24.9	2.36	85	14.6	1.78
Florida																		
Observed	189,126	411.1	1.03	53,659	121.3	0.57	32,399	62.2	0.36	15,804	31.5	0.27	10,643	22.8	0.24	9,092	22.3	0.26
Delay Adjusted	196,358	427.1	1.07	55,555	125.6	0.59	32,643	62.7	0.36	16,212	32.3	0.28	10,974	23.5	0.24	9,736	23.9	0.28
Georgia																		
Observed	74,467	421.3	1.58	22,256	125.2	0.86	10,795	57.9	0.57	6,097	33.8	0.44	3,944	21.3	0.35	4,774	29.6	0.44
Delay Adjusted	75,771	428.6	1.61	22,459	126.4	0.87	10,909	58.5	0.57	6,156	34.1	0.45	3,967	21.4	0.35	4,918	30.5	0.45
Hawaii																		
Observed	4,353	452.9	7.41	1,375	143.3	4.17	450	44.4	2.21	343	35.4	2.05	252	24.0	1.62	457	49.8	2.53
Delay Adjusted	4,442	462.0	7.56	1,379	143.7	4.18	456	45.0	2.24	346	35.8	2.07	259	24.6	1.66	478	52.0	2.64
Idaho																		
Observed	16,445	406.6	3.26	4,936	121.9	1.79	2,031	47.7	1.08	1,323	32.3	0.91	1,014	24.0	0.78	894	23.9	0.83
Delay Adjusted	16,905	417.9	3.35	5,004	123.6	1.81	2,072	48.7	1.10	1,343	32.7	0.92	1,032	24.4	0.79	931	24.9	0.86
Illinois																		
Observed	125,192	448.7	1.32	37,050	135.7	0.73	18,092	61.4	0.47	11,169	37.4	0.37	8,568	29.7	0.33	4,920	19.8	0.29
Delay Adjusted	127,571	457.2	1.34	37,422	137.1	0.74	18,295	62.0	0.47	11,264	37.7	0.37	8,645	30.0	0.33	5,097	20.5	0.31
Indiana																		
Observed	72,489	423.8	1.62	20,471	120.9	0.87	11,090	61.9	0.60	6,800	38.2	0.48	4,873	27.6	0.41	2,636	17.0	0.34
Delay Adjusted	75,006	438.5	1.68	20,998	124.0	0.89	11,339	63.3	0.61	6,978	39.2	0.49	4,973	28.2	0.41	2,804	18.1	0.37
Iowa																		
Observed	39,835	436.1	2.29	11,095	124.7	1.24	5,200	53.4	0.76	3,975	40.0	0.66	2,914	31.3	0.60	1,866	23.7	0.58
Delay Adjusted	40,588	444.2	2.33	11,186	125.7	1.25	5,270	54.1	0.77	4,004	40.3	0.67	2,933	31.5	0.60	1,919	24.4	0.60
Kansas																		
Observed	30,906	421.5	2.50	9,020	124.4	1.36	4,154	53.1	0.84	2,791	36.0	0.71	1,854	24.8	0.60	1,340	20.3	0.58
Delay Adjusted	^	^	^	^	^	^	^	^	^	^	^	^	^	^	^	^	^	^

1. Rates are per 100,000 population and are age-adjusted by five-year age groups to the 2000 U.S. standard population based on single years of age.
2. See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.
3. Abbreviations: NOS - Not Otherwise Specified; S.E. - Standard Error

^ There are insufficient historical registry data to calculate delay factors necessary for delay adjusted rates.

\- Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for All Sites combined.

Note: Delay adjusted counts are rounded to the nearest whole number. A small registry and/or rare cancer site may have the same delay adjusted count and observed count, but this is just due to rounding to whole counts. For more information about cancer incidence rates adjusted for reporting delay see https://surveillance.cancer.gov/delay/.

Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Delay Adjusted Counts and Rates[1] for the Five Most Commonly Diagnosed Cancers
Selected Areas in the NAACCR United States Combined[2], 2010-2014, Non-Hispanic White, Females

	All Sites			Breast			Lung & Bronchus			Colon & Rectum			Uterus Corpus & Uterus, NOS[3]			Melanoma of the Skin		
	Count	*Rate*	*S.E.*[3]	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
Kentucky																		
Observed	57,182	470.4	2.02	15,099	124.3	1.04	10,334	80.9	0.81	5,272	42.2	0.59	3,265	26.3	0.47	2,562	22.8	0.47
Delay Adjusted	58,500	481.3	2.06	15,260	125.7	1.05	10,455	81.8	0.82	5,369	43.0	0.60	3,305	26.6	0.48	2,632	23.4	0.48
Louisiana																		
Observed	37,711	421.1	2.23	10,881	121.8	1.20	5,499	58.0	0.79	3,534	38.5	0.66	1,656	18.1	0.46	1,366	16.8	0.47
Delay Adjusted	38,049	424.9	2.25	10,938	122.5	1.21	5,532	58.4	0.80	3,544	38.6	0.67	1,667	18.2	0.46	1,385	17.1	0.48
Maine																		
Observed	20,284	448.6	3.29	5,641	126.5	1.76	3,135	65.4	1.19	1,610	34.4	0.89	1,484	31.1	0.84	830	20.0	0.73
Delay Adjusted	20,621	456.0	3.34	5,736	128.7	1.79	3,169	66.1	1.20	1,621	34.6	0.89	1,496	31.4	0.84	852	20.5	0.75
Maryland																		
Observed	48,447	440.2	2.08	14,755	135.2	1.16	6,679	57.2	0.72	3,719	32.3	0.55	3,186	27.6	0.50	2,661	26.4	0.54
Delay Adjusted	49,670	451.3	2.14	15,006	137.5	1.18	6,828	58.4	0.73	3,771	32.8	0.56	3,224	27.9	0.51	2,674	26.6	0.54
Massachusetts																		
Observed	80,786	453.8	1.67	24,646	141.8	0.94	11,934	63.5	0.60	6,330	33.5	0.44	5,491	29.8	0.41	3,181	19.0	0.35
Delay Adjusted	82,631	464.2	1.70	24,994	143.8	0.95	12,169	64.8	0.61	6,459	34.2	0.45	5,564	30.2	0.42	3,324	19.8	0.37
Michigan																		
Observed	107,371	419.9	1.33	30,809	122.0	0.72	15,975	58.8	0.47	8,784	32.7	0.36	7,798	29.4	0.34	4,020	17.7	0.29
Delay Adjusted	110,027	430.3	1.37	31,334	124.0	0.73	16,390	60.4	0.49	8,978	33.4	0.37	7,913	29.9	0.35	4,133	18.2	0.30
Detroit																		
Observed	39,662	460.0	2.40	11,314	132.8	1.29	6,018	66.3	0.87	3,236	35.2	0.64	2,801	31.6	0.62	1,465	19.3	0.53
Delay Adjusted	40,117	465.3	2.43	11,415	134.0	1.30	6,052	66.6	0.88	3,257	35.4	0.65	2,817	31.8	0.62	1,478	19.4	0.53
Mississippi																		
Observed	23,962	409.6	2.75	6,582	113.4	1.45	3,849	61.0	1.00	2,307	37.6	0.81	1,109	18.8	0.59	1,072	20.7	0.66
Delay Adjusted	24,541	419.5	2.81	6,694	115.4	1.48	3,904	61.9	1.01	2,336	38.1	0.82	1,119	19.0	0.59	1,091	21.1	0.67
Missouri																		
Observed	68,764	424.0	1.67	20,169	126.0	0.92	11,102	64.7	0.62	6,135	36.2	0.48	4,370	26.3	0.41	2,347	16.1	0.35
Delay Adjusted	70,418	434.1	1.71	20,462	127.9	0.93	11,282	65.7	0.63	6,226	36.7	0.48	4,401	26.5	0.41	2,413	16.6	0.36
Montana																		
Observed	12,132	409.7	3.89	3,613	122.6	2.14	1,634	51.7	1.31	1,012	32.9	1.07	796	25.4	0.94	592	22.3	0.98
Delay Adjusted	12,653	427.3	4.06	3,740	126.8	2.21	1,670	52.9	1.34	1,046	34.0	1.11	818	26.1	0.97	628	23.7	1.03
Nebraska																		
Observed	20,375	412.2	3.01	6,026	123.7	1.66	2,623	50.0	1.00	2,002	38.0	0.88	1,369	27.3	0.76	845	19.3	0.70
Delay Adjusted	21,165	428.1	3.12	6,127	125.8	1.68	2,677	51.1	1.02	2,049	38.9	0.90	1,377	27.4	0.77	902	20.6	0.74

Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Delay Adjusted Counts and Rates[1] for the Five Most Commonly Diagnosed Cancers
Selected Areas in the NAACCR United States Combined[2], 2010-2014, Non-Hispanic White, Females

	All Sites			Breast			Lung & Bronchus			Colon & Rectum			Uterus Corpus & Uterus, NOS[3]			Melanoma of the Skin		
	Count	*Rate*	*S.E.[3]*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
New Hampshire																		
Observed	18,571	458.1	3.48	5,757	142.3	1.94	2,673	64.1	1.27	1,429	34.1	0.93	1,420	33.1	0.90	915	23.8	0.82
Delay Adjusted	19,053	469.9	3.57	5,823	143.9	1.96	2,735	65.5	1.30	1,450	34.6	0.94	1,436	33.5	0.91	947	24.6	0.85
New Jersey																		
Observed	91,497	478.0	1.67	26,345	142.3	0.92	12,211	58.9	0.55	8,013	38.1	0.45	6,657	33.6	0.42	4,171	23.8	0.39
Delay Adjusted	93,929	490.6	1.71	26,693	144.1	0.93	12,363	59.7	0.55	8,106	38.5	0.45	6,734	34.0	0.43	4,347	24.8	0.41
New York																		
Observed	191,107	478.0	1.14	54,058	139.6	0.63	26,970	62.8	0.39	15,687	36.4	0.30	13,416	32.7	0.29	7,854	21.4	0.26
Delay Adjusted	194,852	487.3	1.17	54,739	141.3	0.63	27,132	63.2	0.39	15,870	36.8	0.31	13,531	33.0	0.29	8,300	22.6	0.27
North Carolina																		
Observed	92,640	423.4	1.44	28,293	130.3	0.80	13,897	59.4	0.51	7,200	32.0	0.39	5,548	24.4	0.34	4,800	24.0	0.36
Delay Adjusted	95,299	435.4	1.48	28,938	133.2	0.82	14,124	60.3	0.52	7,338	32.6	0.39	5,601	24.6	0.34	4,959	24.8	0.37
North Dakota																		
Observed	7,869	405.4	4.79	2,274	122.2	2.69	968	47.0	1.55	820	39.3	1.43	500	25.3	1.17	393	22.6	1.20
Delay Adjusted	8,095	417.0	4.92	2,295	123.2	2.71	993	48.2	1.59	842	40.4	1.47	511	25.9	1.20	402	23.1	1.22
Ohio																		
Observed	133,602	424.3	1.20	38,491	123.7	0.65	19,725	59.4	0.43	11,766	35.3	0.34	9,536	29.4	0.31	5,211	18.4	0.27
Delay Adjusted	138,625	440.2	1.25	39,488	126.9	0.67	20,185	60.8	0.44	12,082	36.3	0.34	9,722	29.9	0.32	5,506	19.5	0.28
Oklahoma																		
Observed	36,437	397.1	2.16	10,453	114.8	1.17	5,669	57.7	0.78	3,343	34.9	0.62	1,955	20.9	0.49	1,246	15.0	0.45
Delay Adjusted	36,925	402.4	2.19	10,546	115.9	1.18	5,732	58.4	0.79	3,369	35.2	0.63	1,974	21.1	0.50	1,252	15.1	0.45
Oregon																		
Observed	45,236	420.5	2.07	13,795	128.5	1.15	6,213	54.1	0.70	3,487	30.9	0.55	3,004	26.6	0.51	2,594	27.0	0.56
Delay Adjusted	46,261	429.9	2.12	14,035	130.7	1.17	6,329	55.1	0.72	3,558	31.5	0.56	3,037	26.9	0.51	2,632	27.4	0.57
Pennsylvania																		
Observed	166,496	456.5	1.18	46,936	131.8	0.64	22,068	55.9	0.39	14,759	37.1	0.32	12,343	33.0	0.31	6,552	20.2	0.27
Delay Adjusted	169,525	464.8	1.20	47,387	133.0	0.64	22,305	56.5	0.39	14,893	37.4	0.32	12,400	33.1	0.31	6,751	20.8	0.27
Rhode Island																		
Observed	13,802	467.9	4.21	3,868	135.6	2.29	2,095	66.2	1.49	1,133	34.8	1.10	1,010	33.2	1.08	584	21.5	0.95
Delay Adjusted	14,103	478.0	4.30	3,916	137.3	2.32	2,131	67.3	1.52	1,146	35.2	1.11	1,028	33.8	1.10	600	22.0	0.97

1. Rates are per 100,000 population and are age-adjusted by five-year age groups to the 2000 U.S. standard population based on single years of age.
2. See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.
3. Abbreviations: NOS - Not Otherwise Specified; S.E. - Standard Error

^ There are insufficient historical registry data to calculate delay factors necessary for delay adjusted rates.

\- Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for All Sites combined.

Note: Delay adjusted counts are rounded to the nearest whole number. A small registry and/or rare cancer site may have the same delay adjusted count and observed count, but this is just due to rounding to whole counts. For more information about cancer incidence rates adjusted for reporting delay see https://surveillance.cancer.gov/delay/.

Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Delay Adjusted Counts and Rates[1] for the Five Most Commonly Diagnosed Cancers
Selected Areas in the NAACCR United States Combined[2], 2010-2014, Non-Hispanic White, Females

	All Sites			Prostate			Lung & Bronchus			Colon & Rectum			Uterus Corpus & Uterus, NOS[3]			Melanoma of the Skin		
	Count	*Rate*	*S.E.*[3]	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
South Carolina																		
Observed	44,077	418.7	2.07	13,544	128.8	1.15	6,614	58.3	0.73	3,542	32.6	0.56	2,416	22.2	0.47	2,292	24.0	0.53
Delay Adjusted	45,675	433.8	2.15	13,956	132.7	1.19	6,781	59.8	0.75	3,646	33.6	0.58	2,477	22.7	0.48	2,378	24.9	0.55
South Dakota																		
Observed	9,718	424.1	4.52	2,937	132.1	2.56	1,218	50.0	1.47	960	38.0	1.29	675	29.1	1.16	468	23.6	1.16
Delay Adjusted	10,163	443.6	4.72	2,992	134.6	2.60	1,259	51.7	1.52	992	39.3	1.33	683	29.4	1.17	501	25.2	1.24
Tennessee																		
Observed	69,170	422.0	1.65	19,895	121.8	0.89	11,041	63.3	0.61	6,044	35.8	0.47	3,919	23.4	0.38	2,962	19.7	0.38
Delay Adjusted	70,438	429.9	1.69	20,156	123.4	0.90	11,140	63.9	0.62	6,096	36.1	0.48	3,950	23.6	0.39	3,046	20.2	0.39
Texas																		
Observed	155,363	401.9	1.05	47,036	122.5	0.58	22,116	53.8	0.37	13,062	32.6	0.29	8,313	20.8	0.23	5,449	15.4	0.22
Delay Adjusted	161,279	417.2	1.09	48,159	125.4	0.60	22,540	54.9	0.37	13,251	33.1	0.30	8,454	21.1	0.24	5,932	16.8	0.24
Utah																		
Observed	20,185	371.4	2.66	6,265	116.8	1.50	1,271	23.3	0.66	1,467	27.2	0.72	1,426	25.5	0.69	1,646	30.6	0.77
Delay Adjusted	20,509	377.4	2.70	6,279	117.1	1.50	1,286	23.6	0.67	1,476	27.4	0.72	1,429	25.6	0.69	1,671	31.0	0.78
Vermont																		
Observed	8,711	433.6	4.84	2,599	130.5	2.67	1,247	59.2	1.72	699	33.2	1.30	701	33.2	1.30	495	25.6	1.21
Delay Adjusted	8,822	439.0	4.90	2,613	131.2	2.68	1,261	59.9	1.74	709	33.6	1.31	703	33.3	1.30	501	25.9	1.22
Virginia																		
Observed	69,911	404.7	1.58	22,334	130.0	0.90	9,778	54.2	0.56	5,672	32.0	0.44	4,531	24.9	0.38	2,968	18.3	0.35
Delay Adjusted	71,407	413.3	1.61	22,745	132.4	0.91	9,901	54.9	0.56	5,737	32.3	0.44	4,588	25.2	0.38	3,043	18.7	0.36
Washington																		
Observed	72,824	442.0	1.70	22,925	139.3	0.95	9,566	55.7	0.58	5,568	33.1	0.46	4,603	26.3	0.40	4,033	26.5	0.44
Delay Adjusted	74,682	453.3	1.74	23,352	141.9	0.97	9,713	56.5	0.59	5,680	33.7	0.47	4,706	26.8	0.41	4,123	27.1	0.45
Seattle																		
Observed	49,905	454.9	2.11	15,861	144.5	1.19	6,317	55.9	0.72	3,731	33.4	0.57	3,167	27.0	0.50	3,010	29.4	0.56
Delay Adjusted	50,271	458.3	2.12	15,888	144.7	1.19	6,324	55.9	0.72	3,744	33.5	0.57	3,192	27.2	0.50	3,058	29.8	0.56
West Virginia																		
Observed	26,370	441.8	2.83	6,838	115.1	1.45	4,297	67.1	1.04	2,575	41.2	0.84	1,865	30.7	0.74	968	18.3	0.62
Delay Adjusted	27,013	453.1	2.91	7,038	118.5	1.49	4,371	68.2	1.06	2,621	41.9	0.85	1,885	31.0	0.75	1,002	18.9	0.64
Wisconsin																		
Observed	68,035	426.6	1.70	20,182	129.1	0.94	9,021	53.8	0.58	5,583	32.9	0.46	4,977	29.8	0.43	2,878	20.2	0.40
Delay Adjusted	71,765	449.5	1.79	21,083	134.8	0.98	9,493	56.6	0.61	5,867	34.6	0.48	5,305	31.8	0.46	3,037	21.3	0.42

Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Delay Adjusted Counts and Rates[1] for the Five Most Commonly Diagnosed Cancers
Selected Areas in the NAACCR United States Combined[2], 2010-2014, Non-Hispanic White, Females

	All Sites			Prostate			Lung & Bronchus			Colon & Rectum			Uterus Corpus & Uterus, NOS[3]			Melanoma of the Skin		
	Count	*Rate*	*S.E.*[3]	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
Wyoming																		
Observed	5,593	384.7	5.33	1,705	116.1	2.92	684	45.4	1.77	458	31.2	1.50	329	21.7	1.24	295	21.8	1.32
Delay Adjusted	5,657	389.1	5.39	1,722	117.3	2.95	688	45.6	1.78	461	31.4	1.51	332	21.9	1.25	305	22.5	1.37

1. Rates are per 100,000 population and are age-adjusted by five-year age groups to the 2000 U.S. standard population based on single years of age.
2. See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.
3. Abbreviations: NOS - Not Otherwise Specified; S.E. - Standard Error

^ There are insufficient historical registry data to calculate delay factors necessary for delay adjusted rates.

\- Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for All Sites combined.

Note: Delay adjusted counts are rounded to the nearest whole number. A small registry and/or rare cancer site may have the same delay adjusted count and observed count, but this is just due to rounding to whole counts. For more information about cancer incidence rates adjusted for reporting delay see https://surveillance.cancer.gov/delay/.

NAACCR U.S. Combined Delay Adjusted Incidence 2010-2014, Non-Hispanic Black Combined

UNITED STATES

V-89 Combined Delay Adjusted Incidence – Non-Hispanic Black

Average Annual Age-adjusted Cancer Incidence Delay Adjusted Counts and Rates*, Selected Sites
NAACCR United States Combined[1], 2010-2014, Non-Hispanic Black

	MALES (ALL RACES)						FEMALES (ALL RACES)					
	Observed			Delay Adjusted			Observed			Delay Adjusted		
SITES	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
All Sites (invasive & in situ bladder)	419,479	553.1	0.91	432,724	570.7	0.94	403,126	401.6	0.65	416,080	414.5	0.67
Oral Cavity & Pharynx	12,247	14.8	0.14	12,560	15.2	0.15	5,377	5.2	0.07	5,513	5.3	0.07
Digestive												
Esophagus	5,478	7.2	0.10	5,551	7.3	0.11	2,339	2.3	0.05	2,371	2.4	0.05
Stomach	9,880	14.3	0.15	10,055	14.5	0.16	7,547	7.8	0.09	7,678	8.0	0.09
Colon & Rectum	41,389	56.4	0.30	42,354	57.7	0.30	41,429	41.7	0.21	42,392	42.6	0.21
Liver & IBD[2]	15,084	17.2	0.15	15,452	17.6	0.15	5,316	5.1	0.07	5,443	5.2	0.07
Pancreas	12,381	17.3	0.17	12,615	17.6	0.17	14,183	14.7	0.13	14,444	14.9	0.13
Respiratory												
Larynx	6,771	8.7	0.11	6,883	8.9	0.11	1,802	1.7	0.04	1,832	1.7	0.04
Lung & bronchus	62,068	88.0	0.38	63,035	89.4	0.38	49,342	50.1	0.23	50,135	50.9	0.23
Skin												
Melanomas of the skin	795	1.1	0.04	831	1.2	0.04	955	1.0	0.03	996	1.0	0.03
Breast (female cases only)	-	-	-	-	-	-	127,722	125.5	0.36	130,953	128.7	0.37
Female Genital												
Cervix uteri	-	-	-	-	-	-	9,517	9.5	0.10	9,675	9.6	0.10
Corpus & uterus, NOS[2]	-	-	-	-	-	-	26,498	25.6	0.16	26,999	26.1	0.16
Ovary	-	-	-	-	-	-	9,430	9.4	0.10	9,739	9.7	0.10
Male Genital												
Prostate	147,464	186.8	0.52	155,050	196.6	0.55	-	-	-	-	-	-
Testis	1,375	1.5	0.04	1,400	1.5	0.04	-	-	-	-	-	-
Urinary												
Bladder (incl. *in situ*)	12,822	19.8	0.19	13,193	20.4	0.19	6,327	6.7	0.09	6,506	6.9	0.09
Kidney & renal pelvis	19,663	24.8	0.19	20,249	25.6	0.19	12,994	12.9	0.12	13,373	13.3	0.12
Brain & Other Nervous System	4,111	4.9	0.08	4,248	5.0	0.08	3,635	3.6	0.06	3,754	3.7	0.06
Endocrine												
Thyroid gland	3,179	3.8	0.07	3,259	3.9	0.07	14,196	13.8	0.12	14,543	14.2	0.12
Lymphomas												
Hodgkin Lymphoma	2,944	3.2	0.06	3,010	3.3	0.06	2,461	2.4	0.05	2,517	2.4	0.05
Non-Hodgkin Lymphoma	13,417	17.2	0.16	13,935	17.8	0.16	12,103	12.1	0.11	12,571	12.6	0.12
Multiple Myeloma	11,087	15.7	0.16	12,057	17.1	0.17	11,306	11.5	0.11	12,275	12.5	0.12
Leukemias	10,010	13.6	0.15	10,975	14.9	0.16	8,578	8.7	0.10	9,399	9.5	0.10

* Rates are per 100,000 population and are age-adjusted by five-year age groups to the 2000 U.S. standard population based on single years of age and to standard.

\- Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for All Sites combined.

[1] NAACCR United States Combined includes registries for which a U.S. average delay adjustment factor was calculated. See technical notes.

[2] Abbreviations: IBD – Intrahepatic Bile Duct; NOS - Not Otherwise Specified

Note: Delay adjusted counts are rounded to the nearest whole number. A small registry and/or rare cancer site may have the same delay adjusted count and observed count, but this is just due to rounding to whole counts. For more information about cancer incidence rates adjusted for reporting delay see https://surveillance.cancer.gov/delay/.

Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Delay Adjusted Counts and Rates[1] for the Five Most Commonly Diagnosed Cancers
Selected Areas in the NAACCR United States Combined[2], 2010-2014, Non-Hispanic Black, Males

	All Sites			Prostate			Lung & Bronchus			Colon & Rectum			Kidney & Renal Pelvis			Liver & IBD[3]		
	Count	*Rate*	*S.E.*[3]	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
NAACCR U.S. Combined																		
Observed	419,479	553.1	0.91	147,464	186.8	0.52	62,068	88.0	0.38	41,389	56.4	0.30	19,663	24.8	0.19	15,084	17.2	0.15
Delay Adjusted	432,724	570.7	0.94	155,050	196.6	0.55	63,035	89.4	0.38	42,354	57.7	0.30	20,249	25.6	0.19	15,452	17.6	0.15
Alabama																		
Observed	14,317	566.6	5.09	5,168	197.6	2.95	2,325	95.8	2.14	1,541	63.8	1.75	619	23.9	1.02	365	12.5	0.70
Delay Adjusted	14,583	577.1	5.18	5,361	205.5	3.07	2,340	96.4	2.16	1,556	64.4	1.76	626	24.2	1.03	370	12.6	0.71
Alaska																		
Observed	213	471.3	40.08	87	167.2	21.69	27	82.6	18.87	21	45.5	12.17	8	12.9	5.22	9	16.6	6.49
Delay Adjusted	221	488.4	41.52	92	176.7	23.01	28	84.5	19.29	21	46.5	12.46	8	13.3	5.32	9	17.0	6.61
Arizona																		
Observed	2,068	405.4	9.91	654	120.5	5.16	293	65.9	4.26	194	40.3	3.28	126	23.5	2.33	84	14.9	1.75
Delay Adjusted	2,242	439.8	10.76	730	134.7	5.78	300	67.6	4.37	203	42.3	3.44	137	25.5	2.53	89	15.7	1.85
Arkansas																		
Observed	4,939	566.7	8.70	1,658	178.8	4.71	951	114.9	4.03	498	59.1	2.85	198	22.7	1.73	132	13.0	1.21
Delay Adjusted	5,062	580.9	8.92	1,730	186.5	4.91	969	117.0	4.10	501	59.5	2.87	201	23.1	1.76	133	13.1	1.22
California																		
Observed	27,017	521.6	3.35	9,571	174.1	1.87	3,389	71.2	1.29	2,721	54.7	1.11	1,320	24.4	0.71	1,074	18.5	0.59
Delay Adjusted	28,289	546.4	3.51	10,247	186.7	2.00	3,489	73.3	1.33	2,810	56.5	1.15	1,367	25.3	0.73	1,107	19.1	0.61
Greater California																		
Observed	10,723	507.5	5.32	3,856	172.8	2.99	1,350	71.5	2.11	1,090	54.3	1.80	546	24.1	1.11	435	18.3	0.95
Delay Adjusted	11,038	522.5	5.48	4,070	183.0	3.18	1,371	72.7	2.15	1,106	55.1	1.82	557	24.5	1.13	449	18.9	0.98
Greater Bay																		
Observed	5,904	553.5	7.60	2,049	179.3	4.16	767	77.1	2.95	525	51.5	2.38	289	27.1	1.68	288	23.3	1.43
Delay Adjusted	6,258	587.0	8.06	2,205	192.9	4.48	798	80.3	3.07	554	54.3	2.51	297	27.9	1.73	298	24.1	1.49
Los Angeles																		
Observed	10,379	520.6	5.29	3,660	173.1	2.95	1,269	67.8	1.97	1,105	57.3	1.78	484	23.4	1.10	351	16.1	0.89
Delay Adjusted	10,981	550.9	5.60	3,966	187.5	3.19	1,316	70.3	2.04	1,149	59.5	1.85	512	24.8	1.16	359	16.5	0.91
Colorado																		
Observed	1,970	462.7	11.59	689	157.1	6.69	220	60.1	4.46	198	48.9	3.87	104	23.2	2.43	98	18.9	2.08
Delay Adjusted	2,024	475.7	11.92	733	168.0	7.19	222	60.6	4.49	201	49.4	3.92	106	23.7	2.48	98	19.0	2.09
Connecticut																		
Observed	3,628	541.9	9.61	1,312	187.0	5.47	458	75.6	3.78	338	52.9	3.10	169	21.8	1.75	110	14.9	1.50
Delay Adjusted	3,727	556.6	9.88	1,380	197.2	5.78	464	76.6	3.83	342	53.6	3.14	172	22.3	1.79	112	15.2	1.53

Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Delay Adjusted Counts and Rates[1] for the Five Most Commonly Diagnosed Cancers
Selected Areas in the NAACCR United States Combined[2], 2010-2014, Non-Hispanic Black, Males

	All Sites			Prostate			Lung & Bronchus			Colon & Rectum			Kidney & Renal Pelvis			Liver & IBD[3]		
	Count	*Rate*	*S.E.[3]*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
Delaware																		
Observed	2,314	590.5	13.14	916	217.0	7.56	294	81.0	5.06	184	50.9	4.05	124	31.5	3.02	84	18.9	2.20
Delay Adjusted	2,419	617.5	13.74	960	227.6	7.93	300	82.8	5.18	194	53.7	4.27	129	32.8	3.14	88	19.7	2.27
District of Columbia																		
Observed	4,562	604.7	9.23	1,453	188.5	5.10	679	90.1	3.56	449	61.9	3.01	198	26.1	1.91	254	30.1	1.94
Delay Adjusted	4,697	622.6	9.51	1,505	195.2	5.28	693	91.9	3.64	461	63.6	3.10	204	26.8	1.97	260	30.8	1.98
Florida																		
Observed	27,764	482.3	3.07	10,261	173.6	1.80	3,655	67.5	1.18	2,857	51.3	1.02	1,070	17.8	0.57	787	12.0	0.45
Delay Adjusted	28,887	501.3	3.19	10,913	184.8	1.92	3,682	67.9	1.19	2,956	53.1	1.06	1,107	18.4	0.59	803	12.3	0.46
Georgia																		
Observed	29,302	562.8	3.70	10,844	202.3	2.16	4,063	85.5	1.51	2,996	58.4	1.21	1,297	23.2	0.72	807	13.0	0.50
Delay Adjusted	29,845	573.4	3.77	11,194	208.8	2.23	4,109	86.5	1.53	3,036	59.1	1.22	1,328	23.8	0.74	816	13.1	0.51
Hawaii																		
Observed	252	482.5	35.72	103	171.9	18.99	20	47.9	12.81	25	49.6	12.12	10	16.3	5.79	12	18.2	5.56
Delay Adjusted	264	505.8	37.53	112	186.7	20.67	20	48.4	12.97	25	50.0	12.21	10	16.9	5.95	12	18.6	5.71
Idaho																		
Observed	66	399.6	61.00	25	132.8	31.33	-	-	-	-	-	-	-	-	-	8	59.5	28.79
Delay Adjusted	68	412.5	63.17	26	139.6	33.01	-	-	-	-	-	-	-	-	-	8	62.9	30.57
Illinois																		
Observed	21,334	581.7	4.18	6,960	181.5	2.28	3,464	99.6	1.78	2,394	67.6	1.45	1,039	27.2	0.88	654	16.0	0.65
Delay Adjusted	21,749	593.0	4.27	7,256	189.7	2.38	3,505	100.8	1.80	2,422	68.4	1.47	1,060	27.8	0.90	667	16.3	0.67
Indiana																		
Observed	5,620	516.9	7.37	1,613	141.6	3.75	980	97.7	3.33	575	54.1	2.41	321	26.9	1.59	223	17.1	1.20
Delay Adjusted	5,858	539.1	7.69	1,705	149.7	3.96	1,002	99.9	3.40	590	55.5	2.47	331	27.8	1.64	232	17.8	1.25
Iowa																		
Observed	744	542.1	22.89	220	162.4	12.29	124	96.5	9.86	74	50.6	6.95	45	32.3	5.58	35	21.1	4.03
Delay Adjusted	770	561.4	23.73	234	172.9	13.12	125	97.6	9.97	78	53.3	7.31	46	32.9	5.71	36	21.6	4.15
Kansas																		
Observed	1,860	572.4	14.37	627	189.3	8.13	316	101.7	6.18	190	62.8	4.94	83	23.9	2.80	65	16.8	2.21
Delay Adjusted	^	^	^	^	^	^	^	^	^	^	^	^	^	^	^	^	^	^

1. Rates are per 100,000 population and are age-adjusted by five-year age groups to the 2000 U.S. standard population based on single years of age.
2. See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.
3. Abbreviations: IBD - Intrahepatic Bile Duct; S.E. - Standard Error

^ There are insufficient historical registry data to calculate delay factors necessary for delay adjusted rates.

\- Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for All Sites combined.

Note: Delay adjusted counts are rounded to the nearest whole number. A small registry and/or rare cancer site may have the same delay adjusted count and observed count, but this is just due to rounding to whole counts. For more information about cancer incidence rates adjusted for reporting delay see https://surveillance.cancer.gov/delay/.

Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Delay Adjusted Counts and Rates[1] for the Five Most Commonly Diagnosed Cancers
Selected Areas in the NAACCR United States Combined[2], 2010-2014, Non-Hispanic Black, Males

	All Sites			Prostate			Lung & Bronchus			Colon & Rectum			Kidney & Renal Pelvis			Liver & IBD[3]		
	Count	*Rate*	*S.E.*[3]	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
Kentucky																		
Observed	3,980	597.9	10.28	1,150	166.1	5.28	729	119.0	4.78	413	63.2	3.39	235	31.9	2.21	142	18.1	1.62
Delay Adjusted	4,076	612.0	10.52	1,213	174.7	5.54	737	120.3	4.84	421	64.4	3.45	240	32.6	2.26	145	18.4	1.66
Louisiana																		
Observed	17,728	621.1	5.01	5,883	200.8	2.80	2,967	110.1	2.18	1,905	67.0	1.65	795	27.6	1.04	675	19.9	0.81
Delay Adjusted	18,041	632.1	5.10	6,051	206.5	2.88	3,001	111.3	2.20	1,936	68.1	1.68	804	28.0	1.05	685	20.2	0.82
Maine																		
Observed	108	370.1	39.76	35	124.3	23.08	7	30.7	12.73	7	18.7	7.29	-	-	-	7	26.7	11.82
Delay Adjusted	110	377.2	40.52	36	128.0	23.82	7	31.1	12.90	7	18.9	7.34	-	-	-	7	27.4	12.16
Maryland																		
Observed	18,465	534.3	4.23	7,117	194.2	2.46	2,350	74.8	1.67	1,646	50.5	1.35	804	22.5	0.85	753	18.8	0.73
Delay Adjusted	19,100	552.7	4.38	7,415	202.5	2.56	2,403	76.5	1.71	1,708	52.4	1.40	839	23.5	0.88	776	19.3	0.75
Massachusetts																		
Observed	4,419	520.4	8.41	1,720	194.7	5.00	506	67.8	3.24	375	45.3	2.52	182	20.8	1.65	186	19.6	1.54
Delay Adjusted	4,690	552.4	8.93	1,817	205.6	5.28	529	70.8	3.38	398	48.0	2.67	186	21.3	1.69	202	21.3	1.67
Michigan																		
Observed	16,922	587.0	4.76	5,890	196.2	2.69	2,603	96.8	2.00	1,541	55.2	1.48	921	31.3	1.08	655	19.7	0.81
Delay Adjusted	17,175	595.8	4.83	6,044	201.3	2.76	2,633	97.9	2.03	1,559	55.8	1.50	948	32.2	1.11	670	20.2	0.83
Detroit																		
Observed	12,860	631.4	5.84	4,580	215.4	3.33	1,925	101.5	2.43	1,181	60.0	1.83	701	33.8	1.33	508	21.6	1.00
Delay Adjusted	13,041	640.3	5.92	4,688	220.5	3.41	1,940	102.3	2.45	1,188	60.4	1.84	720	34.8	1.37	515	21.9	1.01
Mississippi																		
Observed	12,670	609.8	5.85	4,352	203.4	3.32	2,264	114.6	2.61	1,477	73.0	2.06	600	28.7	1.26	229	9.4	0.67
Delay Adjusted	12,974	624.3	5.99	4,505	210.8	3.45	2,297	116.2	2.64	1,496	73.9	2.08	615	29.4	1.29	238	9.8	0.69
Missouri																		
Observed	7,500	564.3	6.95	2,228	159.2	3.57	1,339	106.0	3.08	745	60.3	2.36	360	25.9	1.45	366	23.6	1.30
Delay Adjusted	7,691	578.8	7.13	2,360	169.1	3.81	1,358	107.5	3.13	755	61.1	2.39	371	26.7	1.49	374	24.1	1.33
Montana																		
Observed	50	588.5	96.16	17	190.9	49.43	9	129.0	47.41	-	-	-	-	-	-	-	-	-
Delay Adjusted	54	636.5	104.48	18	204.3	53.07	9	131.8	48.62	-	-	-	-	-	-	-	-	-
Nebraska																		
Observed	888	621.2	22.94	308	209.4	12.99	125	99.2	9.63	106	80.2	8.66	53	36.2	5.49	35	21.8	3.94
Delay Adjusted	924	646.7	23.89	335	227.6	14.14	127	100.9	9.81	108	81.8	8.83	56	38.1	5.77	37	23.1	4.15

Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Delay Adjusted Counts and Rates[1] for the Five Most Commonly Diagnosed Cancers
Selected Areas in the NAACCR United States Combined[2], 2010-2014, Non-Hispanic Black, Males

	All Sites			Prostate			Lung & Bronchus			Colon & Rectum			Kidney & Renal Pelvis			Liver & IBD[3]		
	Count	*Rate*	*S.E.[3]*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
New Hampshire																		
Observed	129	424.6	41.51	51	164.5	25.39	12	39.6	12.22	15	48.3	13.53	-	-	-	6	23.5	10.48
Delay Adjusted	132	435.4	42.60	53	169.6	26.20	12	40.6	12.61	15	49.1	13.74	-	-	-	6	24.3	10.85
New Jersey																		
Observed	12,850	554.6	5.16	5,024	210.9	3.12	1,581	74.5	1.98	1,217	53.9	1.63	597	24.4	1.04	390	14.8	0.79
Delay Adjusted	13,459	580.9	5.41	5,431	227.9	3.37	1,609	75.9	2.02	1,278	56.5	1.71	611	25.0	1.07	398	15.1	0.80
New York																		
Observed	34,209	572.9	3.25	13,992	226.1	2.00	3,888	69.1	1.17	3,034	53.4	1.02	1,420	22.9	0.63	1,246	18.8	0.56
Delay Adjusted	35,000	586.3	3.33	14,624	236.6	2.09	3,920	69.7	1.18	3,078	54.2	1.04	1,464	23.6	0.65	1,264	19.1	0.56
North Carolina																		
Observed	24,317	567.9	3.92	8,548	191.0	2.21	4,019	100.2	1.70	2,226	53.4	1.22	1,116	25.4	0.81	734	14.6	0.57
Delay Adjusted	25,292	590.6	4.08	9,075	202.7	2.35	4,083	101.8	1.73	2,304	55.2	1.26	1,144	26.0	0.83	746	14.9	0.58
North Dakota																		
Observed	46	431.6	88.49	14	120.7	42.10	7	100.7	47.14	6	59.2	32.38	-	-	-	-	-	-
Delay Adjusted	47	444.9	91.12	15	125.8	44.11	7	102.9	48.04	6	60.8	33.10	-	-	-	-	-	-
Ohio																		
Observed	15,637	532.8	4.46	5,258	171.8	2.48	2,658	96.1	1.95	1,458	51.1	1.41	769	25.2	0.94	607	17.8	0.75
Delay Adjusted	16,210	552.5	4.63	5,579	182.2	2.63	2,720	98.4	1.99	1,497	52.5	1.44	794	26.0	0.97	632	18.6	0.78
Oklahoma																		
Observed	3,118	589.0	11.52	1,073	200.3	6.66	522	103.5	4.91	296	54.8	3.49	144	27.2	2.45	99	15.8	1.73
Delay Adjusted	3,223	608.8	11.90	1,119	208.7	6.93	528	104.6	4.97	298	55.3	3.52	146	27.5	2.49	101	16.1	1.76
Oregon																		
Observed	796	514.3	20.50	260	162.9	11.32	114	81.5	8.63	74	51.8	6.67	42	23.8	4.11	50	30.7	4.86
Delay Adjusted	828	535.8	21.38	282	177.2	12.34	116	82.8	8.76	75	52.7	6.79	43	24.3	4.19	51	31.5	4.97
Pennsylvania																		
Observed	16,184	595.1	4.92	5,152	181.5	2.65	2,476	99.1	2.09	1,457	55.4	1.53	894	31.0	1.08	900	28.9	1.01
Delay Adjusted	16,749	615.9	5.10	5,621	198.1	2.90	2,502	100.1	2.12	1,475	56.1	1.55	925	32.0	1.12	925	29.7	1.04
Rhode Island																		
Observed	493	461.4	22.63	148	126.9	11.27	68	71.9	9.46	38	40.5	7.09	20	18.5	4.56	22	14.6	3.26
Delay Adjusted	511	478.6	23.49	160	137.4	12.20	69	72.8	9.58	38	41.0	7.17	20	18.8	4.64	23	15.0	3.34

1. Rates are per 100,000 population and are age-adjusted by five-year age groups to the 2000 U.S. standard population based on single years of age.
2. See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.
3. Abbreviations: IBD - Intrahepatic Bile Duct; S.E. - Standard Error

^ There are insufficient historical registry data to calculate delay factors necessary for delay adjusted rates.

\- Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for All Sites combined.

Note: Delay adjusted counts are rounded to the nearest whole number. A small registry and/or rare cancer site may have the same delay adjusted count and observed count, but this is just due to rounding to whole counts. For more information about cancer incidence rates adjusted for reporting delay see https://surveillance.cancer.gov/delay/.

Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Delay Adjusted Counts and Rates[1] for the Five Most Commonly Diagnosed Cancers
Selected Areas in the NAACCR United States Combined[2], 2010-2014, Non-Hispanic Black, Males

	All Sites			Prostate			Lung & Bronchus			Colon & Rectum			Kidney & Renal Pelvis			Liver & IBD[3]		
	Count	*Rate*	*S.E.*[3]	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
South Carolina																		
Observed	15,284	559.9	4.87	5,428	189.4	2.75	2,368	90.8	2.01	1,461	55.1	1.55	639	22.8	0.96	387	12.4	0.67
Delay Adjusted	15,819	579.8	5.04	5,722	199.8	2.91	2,427	93.1	2.06	1,505	56.8	1.60	664	23.6	1.00	402	12.9	0.70
South Dakota																		
Observed	61	253.2	38.42	10	33.7	10.92	8	45.0	21.99	7	29.7	11.68	6	22.1	9.29	-	-	-
Delay Adjusted	64	265.9	40.25	11	36.4	11.84	8	46.1	22.46	7	30.8	12.13	6	23.4	9.79	-	-	-
Tennessee																		
Observed	11,089	561.5	5.87	3,734	183.0	3.28	1,897	103.8	2.63	1,091	58.2	1.94	503	24.6	1.20	382	16.1	0.89
Delay Adjusted	11,318	573.1	5.99	3,828	187.6	3.36	1,914	104.7	2.65	1,101	58.7	1.96	511	25.0	1.22	391	16.4	0.91
Texas																		
Observed	29,176	537.3	3.48	8,950	160.1	1.86	4,539	92.8	1.52	3,209	59.6	1.17	1,496	25.9	0.73	1,498	22.7	0.63
Delay Adjusted	30,495	561.6	3.64	9,621	172.9	2.02	4,626	94.6	1.55	3,309	61.4	1.20	1,559	27.0	0.76	1,524	23.1	0.64
Utah																		
Observed	233	542.7	43.54	87	209.8	26.75	23	60.8	15.08	29	66.1	15.26	14	32.0	10.24	16	33.7	9.67
Delay Adjusted	237	552.3	44.29	92	222.3	28.35	23	61.8	15.30	29	66.5	15.36	14	32.6	10.44	17	34.9	10.00
Vermont																		
Observed	33	277.3	54.76	14	115.1	35.02	-	-	-	-	-	-	-	-	-	-	-	-
Delay Adjusted	33	281.8	55.82	14	116.7	35.54	-	-	-	-	-	-	-	-	-	-	-	-
Virginia																		
Observed	17,709	538.3	4.32	6,409	182.5	2.42	2,671	88.9	1.84	1,595	50.1	1.34	900	26.0	0.91	542	13.9	0.63
Delay Adjusted	18,149	551.7	4.43	6,600	188.0	2.49	2,705	90.0	1.86	1,626	51.0	1.37	924	26.7	0.94	561	14.3	0.65
Washington																		
Observed	2,693	522.9	11.50	926	169.2	6.25	314	74.6	4.73	224	44.2	3.35	161	27.8	2.44	138	23.9	2.28
Delay Adjusted	2,777	539.6	11.87	968	176.7	6.53	316	75.2	4.77	232	45.6	3.46	166	28.6	2.52	141	24.4	2.32
Seattle																		
Observed	2,448	536.7	12.38	854	174.2	6.68	284	77.6	5.16	197	44.6	3.61	142	27.7	2.60	132	26.4	2.57
Delay Adjusted	2,515	551.4	12.72	896	182.8	7.01	284	77.7	5.17	204	46.2	3.74	145	28.3	2.66	134	26.8	2.61
West Virginia																		
Observed	846	521.9	19.60	281	160.4	10.33	131	92.1	8.79	91	56.0	6.44	46	25.7	4.02	38	18.9	3.27
Delay Adjusted	869	535.8	20.10	288	164.5	10.59	133	93.6	8.94	93	57.0	6.55	48	26.6	4.18	39	19.4	3.35
Wisconsin																		
Observed	3,845	673.4	12.16	1,235	209.9	6.63	605	119.0	5.39	375	67.8	3.94	192	30.1	2.40	171	24.9	2.02
Delay Adjusted	4,021	708.1	12.86	1,308	222.5	7.02	637	125.2	5.67	392	70.8	4.11	208	32.6	2.60	180	26.2	2.13

Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Delay Adjusted Counts and Rates[1] for the Five Most Commonly Diagnosed Cancers
Selected Areas in the NAACCR United States Combined[2], 2010-2014, Non-Hispanic Black, Males

	All Sites			Prostate			Lung & Bronchus			Colon & Rectum			Kidney & Renal Pelvis			Liver & IBD[3]		
	Count	*Rate*	*S.E.*[3]	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
Wyoming																		
Observed	31	186.4	34.82	9	51.9	17.96	-	-	-	-	-	-	-	-	-	-	-	-
Delay Adjusted	31	188.9	35.28	9	52.6	18.18	-	-	-	-	-	-	-	-	-	-	-	-

1. Rates are per 100,000 population and are age-adjusted by five-year age groups to the 2000 U.S. standard population based on single years of age.
2. See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.
3. Abbreviations: IBD - Intrahepatic Bile Duct; S.E. - Standard Error

^ There are insufficient historical registry data to calculate delay factors necessary for delay adjusted rates.

\- Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for All Sites combined.

Note: Delay adjusted counts are rounded to the nearest whole number. A small registry and/or rare cancer site may have the same delay adjusted count and observed count, but this is just due to rounding to whole counts. For more information about cancer incidence rates adjusted for reporting delay see https://surveillance.cancer.gov/delay/.

Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Delay Adjusted Counts and Rates[1] for the Five Most Commonly Diagnosed Cancers
Selected Areas in the NAACCR United States Combined[2], 2010-2014, Non-Hispanic Black, Females

	All Sites			Breast			Lung & Bronchus			Colon & Rectum			Uterus Corpus & Uterus, NOS[3]			Thyroid		
	Count	*Rate*	*S.E.*[3]	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
NAACCR U.S. Combined																		
Observed	403,126	401.6	0.65	127,722	125.5	0.36	49,342	50.1	0.23	41,429	41.7	0.21	26,498	25.6	0.16	14,196	13.8	0.12
Delay Adjusted	416,080	414.5	0.67	130,953	128.7	0.37	50,135	50.9	0.23	42,392	42.6	0.21	26,999	26.1	0.16	14,543	14.2	0.12
Alabama																		
Observed	12,796	375.6	3.39	4,306	124.9	1.95	1,312	38.6	1.09	1,504	44.5	1.17	750	21.7	0.81	339	9.9	0.55
Delay Adjusted	13,038	382.6	3.46	4,374	126.9	1.98	1,321	38.9	1.10	1,518	44.9	1.18	762	22.1	0.83	341	10.0	0.55
Alaska																		
Observed	153	341.1	29.20	60	133.7	18.13	11	28.5	8.78	12	23.6	7.18	-	-	-	8	15.5	5.81
Delay Adjusted	159	354.8	30.38	61	136.5	18.51	11	29.3	9.02	12	24.0	7.30	-	-	-	8	15.8	5.93
Arizona																		
Observed	1,926	353.4	8.35	607	107.8	4.51	229	44.5	3.05	189	36.0	2.72	132	23.2	2.09	114	18.9	1.80
Delay Adjusted	2,090	383.6	9.06	669	118.7	4.97	234	45.5	3.12	198	37.7	2.85	144	25.4	2.28	120	19.8	1.90
Arkansas																		
Observed	4,177	371.9	5.91	1,300	114.2	3.25	532	48.3	2.15	498	44.6	2.05	239	20.7	1.38	113	9.7	0.93
Delay Adjusted	4,280	381.1	6.05	1,355	119.0	3.38	542	49.2	2.19	501	45.0	2.07	241	20.8	1.39	115	9.9	0.95
California																		
Observed	24,949	401.1	2.59	8,101	129.0	1.46	3,063	49.7	0.92	2,634	42.7	0.85	1,678	25.7	0.64	776	12.7	0.47
Delay Adjusted	26,167	420.7	2.71	8,388	133.6	1.51	3,158	51.3	0.94	2,724	44.2	0.88	1,713	26.3	0.65	786	12.9	0.47
Greater California																		
Observed	9,142	390.2	4.21	3,012	124.9	2.34	1,098	49.3	1.54	942	41.8	1.40	579	23.4	1.00	329	13.1	0.74
Delay Adjusted	9,417	402.0	4.33	3,087	128.1	2.40	1,115	50.1	1.56	956	42.4	1.42	584	23.6	1.01	333	13.3	0.75
Greater Bay																		
Observed	5,226	415.1	5.87	1,660	131.3	3.30	696	55.3	2.15	556	43.9	1.91	352	27.0	1.47	105	8.7	0.87
Delay Adjusted	5,545	440.5	6.24	1,745	138.0	3.46	725	57.6	2.23	586	46.3	2.01	357	27.3	1.49	105	8.8	0.87
Los Angeles																		
Observed	10,578	404.9	4.00	3,429	131.4	2.28	1,268	47.6	1.35	1,136	43.1	1.29	747	27.4	1.02	342	14.1	0.78
Delay Adjusted	11,202	428.7	4.23	3,557	136.3	2.36	1,317	49.5	1.40	1,182	44.8	1.35	771	28.3	1.05	348	14.3	0.79
Colorado																		
Observed	1,602	358.1	9.18	545	119.1	5.22	183	44.1	3.35	147	34.6	2.92	87	19.2	2.12	80	16.2	1.84
Delay Adjusted	1,645	367.7	9.43	550	120.3	5.27	184	44.4	3.37	149	35.0	2.96	88	19.4	2.14	81	16.5	1.87
Connecticut																		
Observed	3,672	400.3	6.73	1,158	122.9	3.67	395	44.4	2.28	339	37.5	2.08	266	27.6	1.72	173	18.4	1.41
Delay Adjusted	3,772	411.2	6.91	1,178	125.0	3.74	400	45.0	2.31	343	38.0	2.10	276	28.7	1.79	176	18.7	1.44

Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Delay Adjusted Counts and Rates[1] for the Five Most Commonly Diagnosed Cancers
Selected Areas in the NAACCR United States Combined[2], 2010-2014, Non-Hispanic Black, Females

	All Sites			Breast			Lung & Bronchus			Colon & Rectum			Uterus Corpus & Uterus, NOS[3]			Thyroid		
	Count	*Rate*	*S.E.[3]*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
Delaware																		
Observed	2,164	422.4	9.33	686	130.6	5.11	276	57.3	3.55	181	35.9	2.74	157	29.9	2.45	94	17.6	1.85
Delay Adjusted	2,264	442.0	9.76	710	135.2	5.29	282	58.6	3.63	190	37.8	2.89	158	30.1	2.47	96	18.1	1.90
District of Columbia																		
Observed	4,853	465.8	6.89	1,438	141.6	3.85	622	57.5	2.35	508	47.9	2.19	385	35.2	1.84	170	18.1	1.44
Delay Adjusted	5,001	480.0	7.10	1,515	149.2	4.06	635	58.7	2.40	521	49.1	2.24	400	36.6	1.91	178	18.9	1.52
Florida																		
Observed	26,161	353.3	2.23	8,145	108.3	1.22	2,595	35.9	0.72	2,673	36.8	0.73	1,953	25.6	0.59	910	11.9	0.40
Delay Adjusted	27,252	367.8	2.32	8,496	113.0	1.27	2,615	36.2	0.72	2,767	38.1	0.75	2,023	26.5	0.61	916	12.0	0.40
Georgia																		
Observed	28,379	387.8	2.39	9,577	126.3	1.33	2,861	41.2	0.80	3,036	42.6	0.80	1,846	24.6	0.59	926	11.5	0.39
Delay Adjusted	28,906	395.0	2.43	9,670	127.5	1.35	2,893	41.7	0.81	3,076	43.2	0.81	1,878	25.0	0.60	938	11.7	0.39
Hawaii																		
Observed	117	320.6	32.90	49	120.5	18.63	9	38.0	13.17	6	21.1	9.83	6	17.4	7.69	-	-	-
Delay Adjusted	122	334.4	34.37	52	128.7	19.95	9	38.5	13.36	6	21.2	9.87	6	17.8	7.90	-	-	-
Idaho																		
Observed	40	359.4	62.51	12	111.2	35.98	-	-	-	-	-	-	-	-	-	-	-	-
Delay Adjusted	41	370.3	64.41	12	113.5	36.67	-	-	-	-	-	-	-	-	-	-	-	-
Illinois																		
Observed	22,129	435.8	2.98	6,725	131.8	1.63	3,215	63.5	1.14	2,469	48.9	1.00	1,457	27.5	0.73	650	12.8	0.51
Delay Adjusted	22,570	444.5	3.03	6,833	133.9	1.66	3,254	64.3	1.15	2,497	49.5	1.01	1,487	28.1	0.75	660	13.0	0.52
Indiana																		
Observed	6,057	420.1	5.52	1,916	130.6	3.05	875	62.3	2.15	619	43.8	1.80	341	23.1	1.28	152	10.0	0.83
Delay Adjusted	6,317	438.2	5.76	1,977	134.8	3.14	895	63.7	2.20	635	45.0	1.85	349	23.6	1.31	154	10.2	0.84
Iowa																		
Observed	657	443.0	18.13	162	105.1	8.60	102	74.4	7.68	65	47.3	6.11	45	30.1	4.65	17	9.5	2.37
Delay Adjusted	679	458.4	18.78	163	106.0	8.67	103	75.2	7.75	69	49.9	6.44	45	30.3	4.70	17	9.6	2.40
Kansas																		
Observed	1,616	425.8	10.79	482	126.2	5.85	235	63.7	4.22	152	40.5	3.34	95	25.1	2.62	77	19.6	2.27
Delay Adjusted	^	^	^	^	^	^	^	^	^	^	^	^	^	^	^	^	^	^

1. Rates are per 100,000 population and are age-adjusted by five-year age groups to the 2000 U.S. standard population based on single years of age.
2. See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.
3. Abbreviations: NOS - Not Otherwise Specified; S.E. - Standard Error

^ There are insufficient historical registry data to calculate delay factors necessary for delay adjusted rates.

\- Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for All Sites combined.

Note: Delay adjusted counts are rounded to the nearest whole number. A small registry and/or rare cancer site may have the same delay adjusted count and observed count, but this is just due to rounding to whole counts. For more information about cancer incidence rates adjusted for reporting delay see https://surveillance.cancer.gov/delay/.

Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Delay Adjusted Counts and Rates[1] for the Five Most Commonly Diagnosed Cancers
Selected Areas in the NAACCR United States Combined[2], 2010-2014, Non-Hispanic Black, Females

	All Sites			Breast			Lung & Bronchus			Colon & Rectum			Uterus Corpus & Uterus, NOS[3]			Thyroid		
	Count	*Rate*	*S.E.[3]*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
Kentucky																		
Observed	3,739	441.6	7.40	1,090	127.0	3.94	640	78.1	3.17	421	49.6	2.48	200	23.5	1.70	133	15.4	1.36
Delay Adjusted	3,831	452.3	7.58	1,105	128.7	3.99	647	79.0	3.21	429	50.5	2.52	201	23.6	1.71	135	15.6	1.38
Louisiana																		
Observed	15,505	418.6	3.44	4,964	132.5	1.93	1,848	50.4	1.20	1,852	50.5	1.20	840	21.9	0.78	526	14.0	0.62
Delay Adjusted	15,783	426.1	3.51	5,036	134.4	1.95	1,869	51.0	1.22	1,882	51.3	1.22	848	22.1	0.78	545	14.5	0.65
Maine																		
Observed	72	306.0	38.60	15	61.7	16.51	10	48.9	16.56	-	-	-	6	26.3	11.50	-	-	-
Delay Adjusted	74	312.5	39.42	15	62.8	16.84	10	49.5	16.75	-	-	-	6	26.4	11.58	-	-	-
Maryland																		
Observed	19,161	400.1	2.97	6,513	132.4	1.68	2,213	47.4	1.04	1,763	37.5	0.92	1,310	26.4	0.75	760	15.2	0.56
Delay Adjusted	19,826	414.0	3.08	6,697	136.2	1.73	2,264	48.5	1.06	1,829	38.9	0.96	1,341	27.0	0.77	780	15.6	0.58
Massachusetts																		
Observed	4,234	384.1	6.01	1,281	114.7	3.26	481	45.9	2.13	401	36.3	1.85	276	24.5	1.50	268	22.9	1.42
Delay Adjusted	4,502	408.4	6.39	1,319	118.2	3.36	504	48.0	2.23	425	38.5	1.96	280	24.9	1.53	273	23.4	1.45
Michigan																		
Observed	16,089	425.1	3.42	4,828	127.1	1.87	2,395	63.4	1.33	1,610	42.6	1.08	1,058	26.8	0.84	529	14.1	0.63
Delay Adjusted	16,341	431.8	3.48	4,887	128.6	1.89	2,424	64.1	1.34	1,629	43.1	1.10	1,070	27.1	0.85	536	14.3	0.63
Detroit																		
Observed	12,364	441.0	4.05	3,722	132.4	2.22	1,822	64.6	1.55	1,275	45.6	1.30	794	27.1	0.99	433	15.7	0.77
Delay Adjusted	12,545	447.5	4.11	3,755	133.6	2.24	1,837	65.2	1.56	1,283	45.9	1.31	800	27.3	0.99	438	15.9	0.78
Mississippi																		
Observed	11,035	401.2	3.91	3,396	121.0	2.12	1,271	47.8	1.37	1,434	52.6	1.42	639	22.8	0.93	350	12.2	0.66
Delay Adjusted	11,300	410.8	4.00	3,472	123.8	2.17	1,289	48.5	1.39	1,452	53.3	1.44	641	22.9	0.93	358	12.5	0.68
Missouri																		
Observed	7,934	440.2	5.03	2,431	133.4	2.75	1,214	68.2	2.00	785	44.0	1.60	497	26.9	1.23	227	12.3	0.83
Delay Adjusted	8,135	451.3	5.16	2,471	135.5	2.80	1,232	69.2	2.03	796	44.6	1.62	507	27.4	1.26	236	12.8	0.86
Montana																		
Observed	17	309.8	80.22	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Delay Adjusted	18	330.5	85.35	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Nebraska																		
Observed	723	431.3	16.54	213	127.8	8.99	107	65.4	6.49	70	43.5	5.34	34	19.1	3.37	31	15.9	2.93
Delay Adjusted	752	448.4	17.20	216	129.8	9.13	109	66.8	6.63	71	44.4	5.46	34	19.2	3.40	32	16.2	2.98

Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Delay Adjusted Counts and Rates[1] for the Five Most Commonly Diagnosed Cancers
Selected Areas in the NAACCR United States Combined[2], 2010-2014, Non-Hispanic Black, Females

	All Sites			Breast			Lung & Bronchus			Colon & Rectum			Uterus Corpus & Uterus, NOS[3]			Thyroid		
	Count	*Rate*	*S.E.[3]*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
New Hampshire																		
Observed	63	227.0	29.82	21	77.9	17.61	7	26.2	10.05	6	22.6	9.65	6	21.9	9.47	-	-	-
Delay Adjusted	65	233.1	30.62	21	78.8	17.82	7	26.8	10.27	6	23.0	9.81	6	22.0	9.54	-	-	-
New Jersey																		
Observed	13,386	410.0	3.59	4,176	125.9	1.97	1,595	49.3	1.25	1,394	43.3	1.18	949	28.3	0.93	459	14.1	0.66
Delay Adjusted	14,038	429.9	3.77	4,327	130.5	2.05	1,624	50.2	1.28	1,463	45.4	1.23	967	28.8	0.95	472	14.5	0.68
New York																		
Observed	33,983	397.2	2.18	10,215	119.5	1.20	3,722	43.1	0.72	3,302	38.6	0.68	2,851	32.1	0.61	1,356	16.0	0.44
Delay Adjusted	34,783	406.5	2.24	10,374	121.3	1.22	3,753	43.5	0.72	3,350	39.2	0.69	2,880	32.4	0.62	1,387	16.4	0.45
North Carolina																		
Observed	23,393	404.7	2.70	7,837	134.1	1.55	2,725	47.6	0.93	2,209	38.4	0.83	1,519	25.5	0.67	817	14.1	0.50
Delay Adjusted	24,338	421.0	2.81	8,121	139.0	1.60	2,769	48.3	0.95	2,286	39.7	0.86	1,536	25.8	0.68	839	14.5	0.52
North Dakota																		
Observed	26	307.7	77.30	6	55.5	22.84	-	-	-	-	-	-	-	-	-	-	-	-
Delay Adjusted	27	318.2	80.04	6	56.0	23.07	-	-	-	-	-	-	-	-	-	-	-	-
Ohio																		
Observed	15,334	402.0	3.30	4,740	123.8	1.83	2,424	63.8	1.31	1,445	37.9	1.01	919	23.3	0.78	602	16.3	0.68
Delay Adjusted	15,909	417.1	3.42	4,864	127.0	1.88	2,481	65.3	1.35	1,484	38.9	1.04	937	23.7	0.80	619	16.8	0.70
Oklahoma																		
Observed	2,685	401.0	7.95	839	122.9	4.35	372	57.8	3.09	271	40.2	2.51	164	23.6	1.89	84	11.8	1.32
Delay Adjusted	2,774	414.4	8.22	864	126.6	4.48	376	58.4	3.12	273	40.5	2.53	166	23.8	1.91	85	11.9	1.33
Oregon																		
Observed	603	407.0	17.07	192	127.1	9.40	77	55.4	6.51	54	38.8	5.45	37	24.0	4.05	28	17.4	3.35
Delay Adjusted	628	423.8	17.78	196	129.4	9.57	78	56.3	6.62	55	39.5	5.55	37	24.3	4.11	29	17.8	3.41
Pennsylvania																		
Observed	17,088	463.4	3.60	4,810	130.8	1.92	2,573	70.0	1.40	1,581	42.6	1.09	1,121	29.4	0.89	819	22.7	0.81
Delay Adjusted	17,687	479.6	3.73	4,898	133.2	1.95	2,600	70.8	1.42	1,602	43.1	1.10	1,134	29.7	0.90	860	23.8	0.85
Rhode Island																		
Observed	502	369.1	16.81	156	113.4	9.25	66	49.6	6.22	41	30.7	4.88	32	23.9	4.32	36	24.4	4.13
Delay Adjusted	520	382.4	17.41	165	119.8	9.78	67	50.1	6.28	41	31.0	4.94	33	24.4	4.40	36	24.6	4.17

1. Rates are per 100,000 population and are age-adjusted by five-year age groups to the 2000 U.S. standard population based on single years of age.
2. See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.
3. Abbreviations: NOS - Not Otherwise Specified; S.E. - Standard Error

^ There are insufficient historical registry data to calculate delay factors necessary for delay adjusted rates.

\- Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for All Sites combined.

Note: Delay adjusted counts are rounded to the nearest whole number. A small registry and/or rare cancer site may have the same delay adjusted count and observed count, but this is just due to rounding to whole counts. For more information about cancer incidence rates adjusted for reporting delay see https://surveillance.cancer.gov/delay/.

Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Delay Adjusted Counts and Rates[1] for the Five Most Commonly Diagnosed Cancers
Selected Areas in the NAACCR United States Combined[2], 2010-2014, Non-Hispanic Black, Females

	All Sites			Prostate			Lung & Bronchus			Colon & Rectum			Uterus Corpus & Uterus, NOS[3]			Thyroid		
	Count	*Rate*	*S.E.*[3]	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
South Carolina																		
Observed	13,912	385.1	3.35	4,573	125.7	1.91	1,536	42.3	1.11	1,364	37.2	1.03	992	26.5	0.86	461	13.0	0.62
Delay Adjusted	14,410	399.0	3.47	4,713	129.6	1.96	1,574	43.3	1.13	1,404	38.3	1.06	1,020	27.3	0.89	474	13.3	0.63
South Dakota																		
Observed	38	212.5	35.46	16	93.4	23.62	-	-	-	-	-	-	-	-	-	-	-	-
Delay Adjusted	40	221.9	37.01	16	95.1	24.03	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee																		
Observed	10,804	400.0	3.96	3,506	126.3	2.19	1,357	52.3	1.47	1,123	42.8	1.31	617	22.0	0.92	292	10.5	0.62
Delay Adjusted	11,034	408.5	4.04	3,552	128.0	2.22	1,369	52.8	1.48	1,134	43.2	1.33	622	22.2	0.92	296	10.6	0.63
Texas																		
Observed	28,396	396.0	2.42	8,919	120.3	1.31	3,335	49.4	0.89	3,021	42.3	0.80	1,682	22.8	0.58	1,004	12.9	0.42
Delay Adjusted	29,682	413.9	2.53	9,226	124.4	1.35	3,398	50.3	0.90	3,114	43.6	0.82	1,729	23.5	0.59	1,053	13.5	0.44
Utah																		
Observed	126	340.9	32.82	34	90.3	16.21	13	39.8	11.72	16	50.6	13.28	-	-	-	7	14.0	5.81
Delay Adjusted	128	346.8	33.41	35	93.4	16.77	13	40.8	12.02	16	50.8	13.34	-	-	-	7	14.0	5.82
Vermont																		
Observed	23	253.0	58.19	8	75.1	28.27	-	-	-	-	-	-	-	-	-	-	-	-
Delay Adjusted	23	256.5	59.10	8	75.5	28.40	-	-	-	-	-	-	-	-	-	-	-	-
Virginia																		
Observed	16,752	393.6	3.10	5,807	134.3	1.80	2,005	48.0	1.09	1,696	40.2	1.00	950	21.8	0.72	536	12.5	0.55
Delay Adjusted	17,174	403.5	3.18	5,920	136.9	1.83	2,030	48.6	1.11	1,728	40.9	1.01	962	22.1	0.73	545	12.7	0.56
Washington																		
Observed	1,937	382.8	9.19	656	125.7	5.15	216	48.1	3.44	152	30.0	2.57	98	19.5	2.07	96	15.9	1.69
Delay Adjusted	1,996	394.4	9.47	663	127.0	5.21	217	48.3	3.45	157	31.0	2.66	99	19.7	2.10	98	16.3	1.72
Seattle																		
Observed	1,800	398.4	9.94	612	130.3	5.54	203	51.5	3.80	141	31.5	2.80	92	20.6	2.27	91	16.9	1.84
Delay Adjusted	1,850	409.3	10.21	616	131.2	5.58	203	51.6	3.80	146	32.7	2.90	93	20.7	2.29	92	17.2	1.87
West Virginia																		
Observed	641	376.7	15.26	200	120.1	8.74	90	51.3	5.51	64	37.5	4.81	36	19.3	3.32	25	15.6	3.21
Delay Adjusted	659	387.4	15.70	211	126.7	9.21	92	52.3	5.62	65	38.1	4.89	36	19.6	3.36	26	16.0	3.29
Wisconsin																		
Observed	3,448	476.5	8.47	1,001	133.5	4.37	517	76.9	3.53	308	43.4	2.58	205	28.0	2.03	127	15.4	1.42
Delay Adjusted	3,607	500.2	8.93	1,047	139.6	4.57	544	81.0	3.71	322	45.4	2.70	219	29.9	2.17	133	16.1	1.48

Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Delay Adjusted Counts and Rates[1] for the Five Most Commonly Diagnosed Cancers
Selected Areas in the NAACCR United States Combined[2], 2010-2014, Non-Hispanic Black, Females

	All Sites			Prostate			Lung & Bronchus			Colon & Rectum			Uterus Corpus & Uterus, NOS[3]			Thyroid		
	Count	*Rate*	*S.E.[3]*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
Wyoming																		
Observed	29	182.4	35.04	7	48.8	19.06	-	-	-	-	-	-	-	-	-	-	-	-
Delay Adjusted	29	184.9	35.57	7	49.2	19.22	-	-	-	-	-	-	-	-	-	-	-	-

1. Rates are per 100,000 population and are age-adjusted by five-year age groups to the 2000 U.S. standard population based on single years of age.
2. See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.
3. Abbreviations: NOS - Not Otherwise Specified; S.E. - Standard Error

^ There are insufficient historical registry data to calculate delay factors necessary for delay adjusted rates.

\- Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for All Sites combined.

Note: Delay adjusted counts are rounded to the nearest whole number. A small registry and/or rare cancer site may have the same delay adjusted count and observed count, but this is just due to rounding to whole counts. For more information about cancer incidence rates adjusted for reporting delay see https://surveillance.cancer.gov/delay/.

NAACCR Canadian Combined Delay Adjusted Incidence 2010-2014, All Races Combined

CANADA

V-103 Combined Delay Adjusted Incidence – All Races

Average Annual Age-adjusted Cancer Incidence Delay Adjusted Counts and Rates*, Selected Sites
NAACCR Canadian Combined[1], 2010-2014, All Races

	MALES (ALL RACES)						FEMALES (ALL RACES)					
	Observed			Delay Adjusted			Observed			Delay Adjusted		
SITES	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
All Sites (invasive & in situ bladder)	173,695	480.4	1.18	175,448	485.4	1.19	162,514	404.8	1.02	164,125	408.8	1.03
Oral Cavity & Pharynx	5,512	14.4	0.20	5,567	14.5	0.20	2,425	6.0	0.12	2,450	6.1	0.13
Digestive												
Esophagus	2,835	7.7	0.15	2,852	7.7	0.15	830	2.0	0.07	835	2.0	0.07
Stomach	3,686	10.4	0.18	3,701	10.5	0.18	1,846	4.5	0.11	1,853	4.5	0.11
Colon & Rectum	24,708	69.0	0.45	24,950	69.7	0.45	19,829	48.1	0.35	20,023	48.5	0.35
Liver & IBD[2]	3,238	8.5	0.15	3,390	8.9	0.16	1,447	3.5	0.09	1,515	3.7	0.10
Pancreas	4,378	12.3	0.19	4,452	12.5	0.19	4,175	10.0	0.16	4,244	10.2	0.16
Respiratory												
Larynx	1,524	4.1	0.11	1,539	4.1	0.11	273	0.7	0.04	275	0.7	0.04
Lung & bronchus	22,994	64.9	0.44	23,264	65.7	0.44	22,021	53.6	0.37	22,280	54.3	0.37
Skin												
Melanomas of the skin	6,951	19.3	0.24	7,102	19.7	0.24	6,104	15.8	0.21	6,229	16.1	0.21
Breast (female cases only)	-	-	-	-	-	-	46,172	115.6	0.55	46,437	116.3	0.55
Female Genital												
Cervix uteri	-	-	-	-	-	-	2,679	7.6	0.15	2,719	7.7	0.15
Corpus & uterus, NOS[2]	-	-	-	-	-	-	10,414	25.1	0.25	10,490	25.3	0.25
Ovary	-	-	-	-	-	-	4,563	11.4	0.17	4,649	11.6	0.18
Male Genital												
Prostate	42,600	112.5	0.56	42,887	113.3	0.56	-	-	-	-	-	-
Testis	2,008	6.0	0.14	2,030	6.1	0.14	-	-	-	-	-	-
Urinary												
Bladder (incl. *in situ*)	12,665	36.7	0.33	12,917	37.4	0.34	3,994	9.6	0.15	4,071	9.8	0.16
Kidney & renal pelvis	6,996	18.9	0.23	7,235	19.5	0.24	3,867	9.6	0.16	3,995	9.9	0.16
Brain & Other Nervous System	2,679	7.6	0.15	2,736	7.7	0.15	1,903	5.1	0.12	1,944	5.2	0.12
Endocrine												
Thyroid gland	2,034	5.6	0.13	2,052	5.7	0.13	5,637	15.7	0.21	5,685	15.8	0.22
Lymphomas												
Hodgkin Lymphoma	890	2.6	0.09	896	2.6	0.09	677	2.0	0.08	682	2.0	0.08
Non-Hodgkin Lymphoma	8,023	22.6	0.26	8,087	22.8	0.26	6,447	16.0	0.20	6,498	16.1	0.20
Multiple Myeloma	2,560	7.1	0.14	2,620	7.3	0.15	1,875	4.6	0.11	1,921	4.7	0.11
Leukemias	6,188	17.7	0.23	6,498	18.6	0.24	4,056	10.4	0.17	4,258	11.0	0.18

* Rates are per 100,000 population and are age-adjusted by five-year age groups to the 2000 U.S. standard population based on single years of age standard.

\- Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for All Sites combined.

[1] NAACCR Canadian Combined includes registries for which a Canada average delay adjustment factor was calculated. See technical notes.

[2] Abbreviations: IBD – Intrahepatic Bile Duct; NOS - Not Otherwise Specified

Note: Delay adjusted counts are rounded to the nearest whole number. A small registry and/or rare cancer site may have the same delay adjusted count and observed count, but this is just due to rounding to whole counts. For more information about cancer incidence rates adjusted for reporting delay see https://surveillance.cancer.gov/delay/.

Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Delay Adjusted Counts and Rates[1] for the Five Most Commonly Diagnosed Cancers
Selected Areas in the NAACCR Canadian Combined[2], 2010-2014, All Races, Males

	All Sites			Prostate			Colon & Rectum			Lung & Bronchus			Bladder			Non-Hodgkin Lymphoma		
	Count	*Rate*	*S.E.*[3]	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
NAACCR Canadian Combined																		
Observed	173,695	480.4	1.18	42,600	112.5	0.56	24,708	69.0	0.45	22,994	64.9	0.44	12,665	36.7	0.33	8,023	22.6	0.26
Delay Adjusted	175,448	485.4	1.19	42,887	113.3	0.56	24,950	69.7	0.45	23,264	65.7	0.44	12,917	37.4	0.34	8,087	22.8	0.26
Alberta																		
Observed	42,908	480.3	2.41	11,240	121.1	1.19	5,606	64.1	0.89	5,063	60.6	0.88	3,048	37.0	0.69	1,992	22.4	0.52
Delay Adjusted	43,354	485.7	2.44	11,333	122.3	1.20	5,641	64.5	0.90	5,128	61.5	0.89	3,083	37.4	0.70	2,005	22.6	0.52
British Columbia																		
Observed	61,349	452.5	1.86	15,257	107.5	0.89	8,552	63.4	0.70	7,595	56.2	0.65	4,820	36.7	0.54	2,875	21.7	0.41
Delay Adjusted	61,937	456.9	1.88	15,280	107.7	0.89	8,678	64.4	0.71	7,683	56.8	0.66	4,992	38.0	0.56	2,888	21.8	0.42
Manitoba																		
Observed	15,967	485.8	3.92	3,399	100.1	1.75	2,366	72.6	1.52	2,203	67.9	1.47	1,018	31.7	1.01	771	23.6	0.87
Delay Adjusted	16,132	491.1	3.96	3,470	102.3	1.79	2,380	73.0	1.53	2,221	68.5	1.49	1,022	31.9	1.02	783	23.9	0.88
New Brunswick																		
Observed	12,396	523.2	4.85	3,011	116.8	2.19	1,627	69.3	1.77	2,064	88.5	2.00	943	41.3	1.38	554	24.8	1.09
Delay Adjusted	12,571	530.8	4.92	3,060	118.8	2.23	1,642	69.9	1.79	2,099	90.1	2.04	952	41.7	1.40	562	25.2	1.10
Newfoundland and Labrador																		
Observed	9,031	544.4	6.01	2,231	123.1	2.72	1,612	99.2	2.59	1,341	82.1	2.34	565	36.1	1.60	380	23.3	1.25
Delay Adjusted	9,096	548.3	6.05	2,238	123.6	2.73	1,626	100.1	2.61	1,355	82.9	2.37	566	36.2	1.60	389	23.9	1.28
Northwest Territories																		
Observed	350	474.2	29.89	55	74.5	11.57	68	96.0	13.58	53	87.8	13.70	7	7.6	3.15	19	25.0	6.92
Delay Adjusted	^	^	^	^	^	^	^	^	^	^	^	^	^	^	^	^	^	^
Nova Scotia																		
Observed	15,645	539.0	4.44	3,492	112.5	1.95	2,300	79.9	1.72	2,453	85.5	1.77	1,117	39.8	1.22	738	26.4	1.00
Delay Adjusted	15,793	544.1	4.48	3,506	112.9	1.96	2,314	80.4	1.73	2,478	86.4	1.79	1,126	40.1	1.23	741	26.5	1.01
Prince Edward Island																		
Observed	2,330	534.7	11.45	594	126.8	5.37	335	77.4	4.36	359	82.0	4.46	142	33.5	2.90	93	22.3	2.39
Delay Adjusted	2,352	539.9	11.57	601	128.5	5.44	336	77.6	4.38	362	82.6	4.50	144	34.0	2.94	95	22.7	2.43
Saskatchewan																		
Observed	13,719	467.1	4.05	3,321	109.1	1.92	2,242	76.3	1.64	1,863	64.4	1.51	1,005	35.2	1.12	601	21.0	0.87
Delay Adjusted	^	^	^	^	^	^	^	^	^	^	^	^	^	^	^	^	^	^

1. Rates are per 100,000 population and are age-adjusted by five-year age groups to the 2000 U.S. standard population based on single years of age.
2. See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.
3. Abbreviations: S.E. - Standard Error

^ There are insufficient historical registry data to calculate delay factors necessary for delay adjusted rates.

\- Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for All Sites combined.

Note: Delay adjusted counts are rounded to the nearest whole number. A small registry and/or rare cancer site may have the same delay adjusted count and observed count, but this is just due to rounding to whole counts. For more information about cancer incidence rates adjusted for reporting delay see https://surveillance.cancer.gov/delay/.

Average Annual Age-adjusted (2000 U.S. Standard) Cancer Incidence Delay Adjusted Counts and Rates[1] for the Five Most Commonly Diagnosed Cancers
Selected Areas in the NAACCR Canadian Combined[2], 2010-2014, All Races, Females

	All Sites			Breast			Lung & Bronchus			Colon & Rectum			Uterus Corpus & Uterus, NOS[3]			Non-Hodgkin Lymphoma		
	Count	*Rate*	*S.E.[3]*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
NAACCR Canadian Combined																		
Observed	162,514	404.8	1.02	46,172	115.6	0.55	22,021	53.6	0.37	19,829	48.1	0.35	10,414	25.1	0.25	6,447	16.0	0.20
Delay Adjusted	164,125	408.8	1.03	46,437	116.3	0.55	22,280	54.3	0.37	20,023	48.5	0.35	10,490	25.3	0.25	6,498	16.1	0.20
Alberta																		
Observed	39,914	400.4	2.03	11,844	117.8	1.10	5,104	52.6	0.75	4,317	43.5	0.67	2,597	24.9	0.50	1,513	15.4	0.40
Delay Adjusted	40,318	404.6	2.06	11,884	118.3	1.10	5,170	53.3	0.76	4,344	43.8	0.68	2,605	25.0	0.50	1,523	15.5	0.40
British Columbia																		
Observed	57,166	386.8	1.65	16,780	114.7	0.90	7,566	49.4	0.57	6,916	45.1	0.55	3,635	23.9	0.40	2,332	15.5	0.33
Delay Adjusted	57,722	390.6	1.67	16,934	115.8	0.91	7,658	50.0	0.58	7,019	45.8	0.56	3,683	24.2	0.41	2,342	15.6	0.33
Manitoba																		
Observed	16,094	426.4	3.44	4,342	117.0	1.82	2,310	59.3	1.26	1,940	49.3	1.15	1,199	31.6	0.93	637	16.6	0.67
Delay Adjusted	16,250	430.4	3.47	4,347	117.2	1.82	2,328	59.8	1.27	1,951	49.6	1.15	1,204	31.7	0.93	647	16.8	0.68
New Brunswick																		
Observed	10,931	412.4	4.07	2,889	109.4	2.10	1,594	57.3	1.46	1,287	46.6	1.32	669	24.0	0.95	474	17.7	0.84
Delay Adjusted	11,083	418.0	4.12	2,910	110.1	2.11	1,621	58.3	1.48	1,299	47.0	1.34	673	24.1	0.95	481	18.0	0.85
Newfoundland and Labrador																		
Observed	7,998	446.8	5.15	2,052	113.0	2.57	939	50.0	1.66	1,280	70.4	2.01	545	29.3	1.29	311	17.3	1.01
Delay Adjusted	8,055	449.9	5.18	2,060	113.5	2.58	949	50.5	1.68	1,291	71.0	2.03	549	29.5	1.30	318	17.7	1.03
Northwest Territories																		
Observed	348	431.5	25.51	112	123.7	12.69	34	58.8	10.78	63	79.9	11.10	22	25.2	5.90	10	11.3	3.86
Delay Adjusted	^	^	^	^	^	^	^	^	^	^	^	^	^	^	^	^	^	^
Nova Scotia																		
Observed	14,828	443.9	3.76	3,835	116.2	1.94	2,303	66.0	1.40	1,977	56.7	1.31	847	24.2	0.86	565	16.8	0.73
Delay Adjusted	14,961	447.7	3.79	3,848	116.6	1.95	2,325	66.6	1.41	1,989	57.1	1.31	849	24.3	0.86	568	16.8	0.73
Prince Edward Island																		
Observed	2,089	415.4	9.36	605	118.4	4.94	287	55.1	3.29	278	54.6	3.35	99	19.3	2.01	80	16.2	1.87
Delay Adjusted	2,109	419.3	9.45	606	118.6	4.95	290	55.6	3.33	279	54.7	3.36	99	19.4	2.01	81	16.4	1.90
Saskatchewan																		
Observed	13,146	401.7	3.60	3,713	115.6	1.95	1,884	56.2	1.32	1,771	51.6	1.26	801	24.4	0.88	525	15.8	0.71
Delay Adjusted	^	^	^	^	^	^	^	^	^	^	^	^	^	^	^	^	^	^

1. Rates are per 100,000 population and are age-adjusted by five-year age groups to the 2000 U.S. standard population based on single years of age.
2. See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.
3. Abbreviations: NOS - Not Otherwise Specified; S.E. - Standard Error

^ There are insufficient historical registry data to calculate delay factors necessary for delay adjusted rates.

\- Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for All Sites combined.

Note: Delay adjusted counts are rounded to the nearest whole number. A small registry and/or rare cancer site may have the same delay adjusted count and observed count, but this is just due to rounding to whole counts. For more information about cancer incidence rates adjusted for reporting delay see https://surveillance.cancer.gov/delay/.

NAACCR North American Combined Delay Adjusted Incidence 2010-2014, All Races Combined

NORTH AMERICA

V-107 Combined Delay Adjusted Incidence – All Races

Average Annual Age-adjusted Cancer Incidence Delay Adjusted Counts and Rates*, Selected Sites
NAACCR North American Combined[1], 2010-2014, All Races

	MALES (ALL RACES)						FEMALES (ALL RACES)					
	Observed			Delay Adjusted			Observed			Delay Adjusted		
SITES	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E*	*Count*	*Rate*	*S.E.*	*Count*	*Rate*	*S.E.*
All Sites (invasive & in situ bladder)	3,995,878	491.5	0.25	4,079,479	501.7	0.26	3,882,395	411.1	0.21	3,964,211	419.7	0.22
Oral Cavity & Pharynx	147,015	17.2	0.05	150,232	17.6	0.05	60,660	6.3	0.03	61,934	6.5	0.03
Digestive												
Esophagus	65,650	8.0	0.03	66,472	8.1	0.03	17,711	1.8	0.01	17,929	1.8	0.01
Stomach	73,503	9.3	0.03	74,390	9.4	0.04	44,667	4.6	0.02	45,207	4.7	0.02
Colon & Rectum	375,540	46.9	0.08	380,844	47.6	0.08	343,101	35.4	0.06	347,869	35.9	0.06
Liver & IBD[2]	104,130	11.9	0.04	106,503	12.2	0.04	40,795	4.2	0.02	41,728	4.2	0.02
Pancreas	113,477	14.2	0.04	115,053	14.4	0.04	109,137	11.0	0.03	110,619	11.2	0.03
Respiratory												
Larynx	50,182	5.9	0.03	50,927	6.0	0.03	12,912	1.3	0.01	13,102	1.3	0.01
Lung & bronchus	578,086	72.7	0.10	585,660	73.6	0.10	515,011	52.8	0.07	521,797	53.5	0.08
Skin												
Melanomas of the skin	208,393	26.2	0.06	214,278	27.0	0.06	147,410	16.3	0.04	151,391	16.7	0.04
Breast (female cases only)	-	-	-	-	-	-	1,153,821	123.3	0.12	1,172,132	125.2	0.12
Female Genital												
Cervix uteri	-	-	-	-	-	-	63,013	7.6	0.03	63,954	7.7	0.03
Corpus & uterus, NOS[2]	-	-	-	-	-	-	251,922	25.8	0.05	254,636	26.1	0.05
Ovary	-	-	-	-	-	-	107,897	11.5	0.04	110,553	11.7	0.04
Male Genital												
Prostate	986,455	114.9	0.12	1,015,018	118.3	0.12	-	-	-	-	-	-
Testis	42,384	5.5	0.03	42,976	5.6	0.03	-	-	-	-	-	-
Urinary												
Bladder (incl. *in situ*)	275,044	35.9	0.07	280,741	36.6	0.07	87,596	8.9	0.03	89,390	9.0	0.03
Kidney & renal pelvis	178,896	21.7	0.05	183,177	22.2	0.05	106,317	11.2	0.04	108,831	11.5	0.04
Brain & Other Nervous System	61,785	7.7	0.03	63,258	7.9	0.03	49,208	5.5	0.03	50,361	5.7	0.03
Endocrine												
Thyroid gland	57,702	7.1	0.03	58,595	7.2	0.03	175,611	20.9	0.05	178,334	21.2	0.05
Lymphomas												
Hodgkin Lymphoma	23,845	3.0	0.02	24,258	3.1	0.02	19,123	2.3	0.02	19,450	2.4	0.02
Non-Hodgkin Lymphoma	181,114	22.9	0.05	186,203	23.5	0.06	150,485	15.8	0.04	154,693	16.2	0.04
Multiple Myeloma	64,121	8.0	0.03	68,813	8.6	0.03	51,423	5.3	0.02	55,186	5.7	0.03
Leukemias	135,235	17.4	0.05	147,553	18.9	0.05	98,124	10.5	0.03	106,932	11.5	0.04

* Rates are per 100,000 population and are age-adjusted by five-year age groups to the 2000 U.S. standard population based on single years of age standard.

- Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for All Sites combined.

[1] NAACCR North American Combined includes registries for which a U.S. and Canada average delay adjustment factor was calculated. See technical notes.

[2] Abbreviations: IBD – Intrahepatic Bile Duct; NOS - Not Otherwise Specified

Note: Delay adjusted counts are rounded to the nearest whole number. A small registry and/or rare cancer site may have the same delay adjusted count and observed count, but this is just due to rounding to whole counts. For more information about cancer incidence rates adjusted for reporting delay see https://surveillance.cancer.gov/delay/.