

Average Annual Age-adjusted (2000 U.S. Standard) NAACCR (U.S.) Combined² Cancer Incidence Rates¹ and Rate Ratios for Selected Cancer Sites³ by Race/Ethnicity and Gender, 2011-2015

Males

Cancer Site	Rates			Rate Ratios		
	Hispanic	Non-Hispanic White	Non-Hispanic Black	Hispanic:Non-Hispanic White	Hispanic:Non-Hispanic Black	Hispanic:Non-Hispanic Black
All Sites	381.1	505.4	549.1	0.75		0.69
Prostate	99.1	101.9	179.2	0.97		0.55
Colon and Rectum	43.0	44.6	55.1	0.96		0.78
Lung and Bronchus	36.9	74.1	85.4	0.50		0.43
Kidney and Renal Pelvis	19.8	22.5	25.4	0.88		0.78
NH ⁴ Lymphoma	19.1	23.9	17.0	0.80		1.12
Liver and IBD ⁴	18.9	10.3	17.7	1.83		1.07
Urinary Bladder	18.7	39.5	19.9	0.47		0.94
Leukemia	12.9	18.6	13.7	0.69		0.94
Stomach	12.3	7.8	14.1	1.58		0.87
Pancreas	11.7	14.6	17.3	0.80		0.68
Oral Cavity and Pharynx	11.2	19.2	14.6	0.58		0.77

Females

Cancer Site	Rates			Rate Ratios		
	Hispanic	Non-Hispanic White	Non-Hispanic Black	Hispanic:Non-Hispanic White	Hispanic:Non-Hispanic Black	Hispanic:Non-Hispanic Black
All Sites	327.6	438.3	407.0	0.75		0.80
Breast	93.0	130.1	126.4	0.71		0.74
Colon and Rectum	29.6	34.2	40.7	0.87		0.73
Corpus and Uterus, NOS ⁴	23.1	26.9	26.2	0.86		0.88
Lung and Bronchus	22.8	57.2	49.2	0.40		0.46
Thyroid	22.4	22.9	14.0	0.98		1.60
NH ⁴ Lymphoma	14.6	16.3	12.1	0.90		1.21
Kidney and Renal Pelvis	11.3	11.4	13.1	0.99		0.86
Ovary	9.9	11.8	9.3	0.84		1.06
Cervix	9.9	7.0	9.2	1.41		1.08
Pancreas	9.7	10.9	14.8	0.89		0.66
Leukemia	8.9	11.2	8.9	0.79		1.00

1. Rates are per 100,000 population and are age-adjusted by five-year age groups to the 2000 U.S. standard population based on single years of age. Counts and rates are suppressed when fewer than 6 cases were reported for the specific cancer. The suppressed cases, however, are included in the counts and rates for the U.S. combined.

2. See Technical Notes, Section I, for NAACCR area inclusion criteria for the combined cancer statistics.

3. Selected cancer sites by gender among the U.S. total population and the U.S. Hispanic/Latino population.

4. Abbreviations: NH - Non-Hodgkin; IBD - Intrahepatic Bile Duct NOS - Not Otherwise Specified

N.B. In areas with small Latino populations, methods to indirectly identify Latinos (like NHIAv2) can overestimate the ethnicity-specific counts of cancer cases. Also, even small errors in Latino population estimates can affect the magnitude of the cancer rates. Data Item 190 used in place of NHIA for MA and RI, see technical notes.